

THE UNIVERSITY OF MELBOURNE

**Department of Mathematics and
Statistics**

620-311

Metric Spaces

Lecture Notes

Semester 1, 2004

Notes by: Kris Wysocki

This compilation has been made in accordance with the provisions of Part VB of the copyright act for the teaching purposes of the University. For the use of students of the University of Melbourne enrolled in the subject 620-311.

Metric and Topological Spaces

Contents

1	Introduction	4
2	Metric Spaces	5
3	Continuity	18
4	Complete Spaces	22
5	Compact Metric Spaces	34
6	Topological Spaces	40
7	Compact Topological Spaces	44
8	Connected Spaces	46
9	Product Spaces	50
10	Uryshon's and Thietze's Theorems	56
11	Appendix	59
12	Problem Sheets	69
12.1	Problem Sheet 1	69
12.2	Problem Sheet 2	71
12.3	Problem Sheet 3	73
12.4	Problem Sheet 4	74
12.5	Problem Sheet 5	76
12.6	Problem Sheet 6	77
12.7	Problem Sheet 7	79

12.8 Problem Sheet 8	80
12.9 Problem Sheet 9	81
12.10 Problem Sheet 10	83
12.11 Problem Sheet 11	84

1 Introduction

The ideas of limit and continuity which we encounter in Euclidean spaces occur in various other contexts e.g in function spaces. Set topology is the study of limits and continuity in a general setting. The notion of limit is based on the idea of nearness. These concepts are easier to perceive when the notion of nearness is given by distance. The corresponding spaces are called metric spaces. These are introduced in Chapter 2 and applications to function spaces are discussed early. The desirability of finding limits leads to the notion of completeness and compactness. As we go on, we find that many of the arguments do not really need the notion of distance. This leads to the concept of topological spaces which are discussed from Chapter 6 onward. The idea of compactness is discussed in general setting in Chapter 7 and the notion of connectedness (which is related to the Intermediate Value Theorem) is discussed in Chapter 8. Under mild assumptions we can study abstract topological spaces by constructing continuous functions to the real line; the results known as Uryshon and Thitze's theorem are discussed in Chapter 10. The concepts of completeness and compactness come again in the guise of the important Ascoli-Arzela theorem are discussed in Chapter 9. The necessary preliminary material is collected in Chapter 11. There are 11 problem sheets which do not exactly correspond to the chapters of the notes. This notes is only a brief introduction to the subject and we refer to Munkre's Topology [Mu] for comprehensive treatment. More elementary introductions are the books by Mendelson [M] and Croom [C].

2 Metric Spaces

Basic Concepts

By a **metric space** we mean a set X together with a function $d : X \times X \rightarrow [0, \infty)$ which satisfies the following axioms:

M1 $d(x, y) = 0$ if and only if $x = y$;

M2 $d(x, y) = d(y, x)$ for every $x, y \in X$;

M3 $d(x, z) \leq d(x, y) + d(y, z)$ for every x, y and $z \in X$.

Elements of X are called **points**, a function d is called a **metric** on X , and the value $d(x, y)$ is called a **distance** between x and y . The axiom M2 says that a metric is symmetric, and the axiom M3 is called the **triangle inequality** since it reflects the geometrical fact that the length of one side of a triangle is less or equal to the sum of the lengths of the other two sides.

Examples

Example 2.1. The most important example of a metric space is the set of all real numbers \mathbb{R} with the metric $d(x, y) = |x - y|$. In the following we will call this metric the **usual metric** in \mathbb{R} .

Example 2.2. Let X be any set and let

$$d(x, y) = \begin{cases} 0 & x = y, \\ 1 & x \neq y. \end{cases}$$

Then d is a metric on X called the **discrete metric**.

Example 2.3. Any subset Y of a metric space (X, d) becomes a metric space with the metric

$$d_Y(x, y) = d(x, y) \quad \text{for all } x, y \in Y.$$

The pair (Y, d_Y) is called a **metric subspace** of (X, d) . We will refer to Y as a subspace of X , rather than (Y, d_Y) as a subspace of (X, d) .

Example 2.4. [Cartesian product of finite number of metric spaces]. Consider a finite collection of metric spaces (X_i, d_i) , $1 \leq i \leq n$, and let X be the cartesian product $\prod_{i=1}^n X_i$. For $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n) \in \prod_{i=1}^n X_i$, set

$$d(x, y) = \sum_{i=1}^n d_i(x_i, y_i).$$

Then d is a metric on X . Clearly, axioms M1-M2 are satisfied. To see that d satisfies M3 take $x = (x_1, \dots, x_n)$, $y = (y_1, \dots, y_n)$ and $z = (z_1, \dots, z_n) \in X$. Then

$$d(x, z) = \sum_{i=1}^n d_i(x_i, z_i) \leq \sum_{i=1}^n [d_i(x_i, y_i) + d_i(y_i, z_i)] = d(x, y) + d(y, z)$$

as required. The pair (X, d) defined above is called a **metric product** (or just a product) of (X_i, d_i) , $1 \leq i \leq n$, and the metric d is called a **product metric**. (Other metrics are also used on $\prod_{i=1}^n X_i$).

Norms and normed vector spaces

We next define the class of metric spaces which are the most interesting in analysis. Let X be a vector space over \mathbb{R} (or \mathbb{C}).

Definition 2.5. A **norm** is a function $\|\cdot\| : X \rightarrow \mathbb{R}$ having the following properties

N1 $\|x\| \geq 0$ and $\|x\| = 0$ if and only if $x = 0$.

N2 $\|\alpha x\| = |\alpha| \cdot \|x\|$ for all $x \in X$ and $\alpha \in \mathbb{R}$.

N3 $\|x + y\| \leq \|x\| + \|y\|$ for all $x, y \in X$.

The pair $(X, \|\cdot\|)$ is called a **normed vector space**.

Proposition 2.6. Let X be a normed space. Then

$$d(x, y) = \|x - y\|$$

is a metric on X .

Proof. The axioms M1 and M2 are clear. If x, y and $z \in X$, then, in view of N3,

$$\begin{aligned} d(x, z) &= \|x - z\| = \|(x - y) + (y - z)\| \\ &\leq \|x - y\| + \|y - z\| = d(x, y) + d(y, z), \end{aligned}$$

and so the triangle inequality follows. ■

Examples of normed spaces

Example 2.7. [Euclidean Space] Consider \mathbb{R}^n and let

$$\|x\| = \left[\sum_{i=1}^n x_i^2 \right]^{1/2}$$

for $x = (x_1, \dots, x_n) \in \mathbb{R}^n$. Clearly, N1 and N2 are satisfied. To see that N3 holds we need the **Cauchy inequality**.

Lemma 2.8. For $x, y \in \mathbb{R}^n$,

$$\left| \sum_{i=1}^n x_i y_i \right| \leq \left[\sum_{i=1}^n |x_i|^2 \right]^{1/2} \cdot \left[\sum_{i=1}^n |y_i|^2 \right]^{1/2}.$$

Proof.

$$\begin{aligned} 0 &\leq \sum_{i,j=1}^n (x_i y_j - x_j y_i)^2 = \sum_{i,j=1}^n (x_i^2 y_j^2 - 2x_i x_j y_i y_j + x_j^2 y_i^2) \\ &= \sum_{i=1}^n \sum_{j=1}^n x_i^2 y_j^2 + \sum_{i=1}^n \sum_{j=1}^n x_j^2 y_i^2 - 2 \sum_{i=1}^n \sum_{j=1}^n x_i x_j y_i y_j \\ &= \sum_{i=1}^n \|y\|^2 x_i^2 + \sum_{j=1}^n \|x\|^2 y_j^2 - 2 \left[\sum_{i=1}^n x_i y_i \right]^2 \\ &= 2\|x\|^2 \cdot \|y\|^2 - 2 \left[\sum_{i=1}^n x_i y_i \right]^2 \end{aligned}$$

As a corollary we have

Corollary 2.9.

$$\|x + y\| \leq \|x\| + \|y\| \quad \text{for all } x, y \in \mathbb{R}^n.$$

Proof. In view of the Cauchy inequality we have

$$\begin{aligned}\|x + y\|^2 &= \sum_{i=1}^n |x_i + y_i|^2 = \sum_{i=1}^n x_i^2 + 2 \sum_{i=1}^n x_i y_i + \sum_{i=1}^n y_i^2 \\ &= \|x\|^2 + 2 \sum_{i=1}^n x_i y_i + \|y\|^2 \leq \|x\|^2 + 2\|x\| \cdot \|y\| + \|y\|^2 \\ &= (\|x\| + \|y\|)^2.\end{aligned}$$

By taking square roots of both sides we the desired inequality follows. \blacksquare

Consequently,

$$d(x, y) = \|x - y\| = \left[\sum_{i=1}^n (x_i - y_i)^2 \right]^{1/2}$$

defines a metric on \mathbb{R}^n . We shall call this metric the Euclidean metric or the standard metric.

Example 2.10. [Space of bounded functions.] Let X be a non-empty set. Call a function $f : X \rightarrow \mathbb{R}$ bounded if there exists a constant M such that $|f(x)| \leq M$ for all $x \in X$. Denote by $B(X) = B(X, \mathbb{R})$ the set of all bounded functions from X to \mathbb{R} , and define

$$\|f\| = \sup\{|f(x)| \mid x \in X\}.$$

Then $\|\cdot\|$ is a norm on $B(X)$, and, in view of Proposition 2.6, this norm defines a metric on $B(X)$ by

$$d(f, g) = \|f - g\| = \sup\{|f(x) - g(x)| \mid x \in X\},$$

for $f, g \in B(X)$.

Example 2.11. Let X be the set of all continuous functions $f : [0, 1] \rightarrow \mathbb{R}$. For any $f \in X$ we set

$$\|f\| = \int_0^1 |f(x)| dx.$$

Then $\|\cdot\|$ defines a norm on X which induces a metric on X by

$$d(f, g) = \int_0^1 |f(x) - g(x)| dx, \quad f, g \in X$$

Balls and diameter

Let $x_0 \in X$ and $r > 0$. The set

$$B(x_0, r) = \{x \in X \mid d(x_0, x) < r\}$$

is called an **open ball** with centre at x_0 and radius $r > 0$, and the set

$$\overline{B}(x_0, r) = \{x \in X \mid d(x_0, x) \leq r\}$$

is called a **closed ball** with centre at x_0 and radius $r > 0$.

If A is a non-empty subset of X , then we define the **distance between x and A** by

$$d(x, A) = \inf\{d(x, y) \mid y \in A\}$$

and more generally if B is another non-empty subset of X , then the **distance between A and B** is defined as

$$d(A, B) = \inf\{d(x, y) \mid x \in A, y \in B\}.$$

For a non-empty subset A of X we define its **diameter** by setting

$$\text{diam } A = \sup\{d(x, y) \mid x, y \in A\}.$$

Clearly, if $A \subset B$, then $\text{diam } A \leq \text{diam } B$. A subset $A \subset X$ **bounded** if its diameter is finite, that is, $\text{diam } A < \infty$.

Sequences and Convergence

Convergence of a sequence in a metric space is defined as in calculus.

Definition 2.12. Let $\{x_n\}$ be a sequence of points in (X, d) and $x \in X$. The sequence $\{x_n\}$ is said to **converge** to x , if for every $\varepsilon > 0$ there exists a positive integer k such that

$$d(x_n, x) < \varepsilon \quad \text{for all } n \geq k.$$

A sequence $\{x_n\}$ is said to converge if there is $x \in X$ to which it converges. If there is no such x , then $\{x_n\}$ is said to **diverge**. If $\{x_n\}$ converges to x we write $\lim_n x_n = x_0$ or $x_n \rightarrow x$. The point x is called the **limit** of $\{x_n\}$.

The definition can be expressed in terms of the convergence of sequences of real numbers. Namely, a sequence $\{x_n\}$ converges to $x \in X$ if and only if $d(x_n, x) \rightarrow 0$ as $n \rightarrow \infty$. We are justified in referring to *the* limit because of the following proposition.

Proposition 2.13. *Let $\{x_n\}$ be a sequence in a metric space (X, d) . Then there is at most one point $x \in X$ such that $\{x_n\}$ converges to x .*

Proof. Arguing by contradiction we assume that $x_n \rightarrow x$ and $x_n \rightarrow y$ with $x \neq y$. Then $d(x, y) > 0$ and we can apply the above definition of convergence with $\varepsilon = d(x, y)/2$. We find a positive integer k such that

$$d(x_n, x) < \varepsilon \quad \text{and} \quad d(x_n, y) < \varepsilon, \quad \text{for } n \geq k.$$

By the triangle law

$$d(x, y) \leq d(x, x_n) + d(x_n, y) < \varepsilon + \varepsilon = d(x, y)$$

which gives a contradiction. Hence we conclude that it is impossible for a sequence $\{x_n\}$ to converge to two different points. ■

Given a sequence $\{x_n\}$ of points in X , consider a sequence $\{n_k\}$ such that $n_1 < n_2 < n_3 < \dots$. Then $\{x_{n_k}\}$ is called a **subsequence** of $\{x_n\}$.

Proposition 2.14. *If $X = \prod_{i=1}^n X_i$ is the product of metric spaces (X_i, d_i) , $1 \leq i \leq n$, and $x^m = (x_1^m, x_2^m, \dots, x_n^m) \in X$, then $x^m \rightarrow x = (x_1, \dots, x_n) \in X$ if and only if $x_i^m \rightarrow x_i$ in X_i for $i = 1, \dots, n$.*

Proof. Recall that we consider X with the metric

$$d(x, y) = \sum_{i=1}^n d_i(x_i, y_i)$$

for $x = (x_1, \dots, x_n)$, $y = (y_1, \dots, y_n) \in X$. Observe that

$$d_i(x_i, y_i) \leq d(x, y) \leq n \cdot \max\{d_i(x_i, y_i) \mid 1 \leq i \leq n\}, \quad x, y \in X. \quad (1)$$

Let $x^m \rightarrow x$, where $x = (x_1, \dots, x_n)$. Then given $\varepsilon > 0$ there exists $k \in \mathbb{N}$ such that

$$d(x^m, x) < \varepsilon \quad \text{for } m \geq k.$$

In view of the left hand side inequality (1)

$$d_j(x_j^m, x_j) < \varepsilon \quad \text{for } m \geq k \text{ and } j = 1, \dots, n.$$

So $x_j^m \rightarrow x_j$ as required. Conversely, assume that $x_j^m \rightarrow x_j$ for $j = 1, \dots, n$. Hence for a given $\varepsilon > 0$, there exists $k(j) \in \mathbb{N}$ such that

$$d_j(x_j^m, x_j) < \varepsilon/n \quad \text{for } m \geq k(j).$$

In view of the right hand side inequality in (1) we get

$$d(x^m, x) \leq n \max\{d_j(x_j^m, x_j) \mid j = 1, \dots, n\} < \varepsilon$$

for all $m > k := \max\{k(j) \mid j = 1, \dots, n\}$. Hence $x_n \rightarrow x$ as required. ■

Definition 2.15. Two metrics d and d' in X are called **equivalent** if

$$d(x_n, x_0) \rightarrow 0 \quad \text{if and only if} \quad d'(x_n, x_0) \rightarrow 0.$$

Example 2.16. Let d be a metric on X . Define

$$d'(x, y) = \frac{d(x, y)}{1 + d(x, y)}, \quad x, y \in X. \quad (2)$$

Then d' is a metric on X (show this!) which is equivalent to d . Indeed, if $d(x_n, x_0) \rightarrow 0$, then $d'(x_n, x_0) = \frac{d(x_n, x_0)}{1 + d(x_n, x_0)} \rightarrow 0$. Conversely, $d(x, y) = \frac{d'(x, y)}{1 - d'(x, y)}$. So if $d'(x_n, x_0) \rightarrow 0$, then $d(x_n, x_0) \rightarrow 0$. Note that with respect to this equivalent metric, the space X is *bounded* since $d'(x, y) < 1$ for all $x, y \in X$.

Example 2.17. Consider the product (X, d) of metric spaces (X_i, d_i) . Recall that

$$d(x, y) = \sum_{i=1}^n d_i(x_i, y_i), \quad x = (x_1, \dots, x_n), \quad y = (y_1, \dots, y_n) \in X.$$

Set

$$\sigma(x, y) = \max\{d_i(x_i, y_i) \mid 1 \leq i \leq n\}$$

$$\rho(x, y) = \left[\sum_{i=1}^n d_i(x_i, y_i)^2 \right]^{1/2}.$$

Then d is equivalent to σ and ρ .

Open and closed sets

Definition 2.18. Let $A \subset X$. A point $x \in A$ is called an **interior point** of A , if $B(x, r) \subset A$ for some $r > 0$. The collection of all interior points of a set A is called the **interior** of A , and is denoted by A° . A set A is called **open** if $A = A^\circ$.

Obviously, the interior of any set is an open set. Hence open sets A are characterized by equality $A^\circ = A$.

Example 2.19. The empty set \emptyset and the whole space X are open in any metric space X . If X is equipped with the discrete metric d , then any subset of X is open.

Example 2.20. The set \mathbb{Q} is not open in \mathbb{R} with the usual metric but it is open in (\mathbb{R}, d) , where d is the discrete metric in \mathbb{R} . Indeed, if $x \in \mathbb{Q}$ and $r > 0$, then for large $n \in \mathbb{N}$, we have

$$x < x + \frac{\sqrt{2}}{n} < x + r$$

so that $x + \sqrt{2}/n \in B(x, r)$ but $x + \sqrt{2}/n \notin \mathbb{Q}$. In the case of the discrete metric, for every $x \in \mathbb{Q}$, $B(x, 1/2) = \{x\} \subset \mathbb{Q}$, so \mathbb{Q} is open in (\mathbb{R}, d) .

Example 2.21. Let $B(x, R)$ be an open ball in a metric space X . Then $B(x, R)$ is an open set. Indeed, let $y \in B(x, R)$. We have to show that y is an interior point of $B(x, R)$, that is, $B(y, r) \subset B(x, R)$ for some $r > 0$. Set $r = R - d(x, y)$. Then for any $z \in B(y, r)$,

$$d(x, z) \leq d(x, y) + d(y, z) < d(x, y) + r = d(x, y) + [R - d(x, y)] = R.$$

Thus $B(y, r) \subset B(x, R)$ as required.




Figure 1: An open ball is an open set

More terminology: if $x \in X$, then a set $A \subset X$ is called a **neighbourhood** of x , if $x \in A^\circ$.

Definition 2.22. A point $x \in X$ is **adherent** to A provided that $B(x, r) \cap A \neq \emptyset$ for all $r > 0$. The set of all the adherent points of A is called the **closure** of A and is denoted by \overline{A} . If $A = \overline{A}$, then A is called **closed**.

Proposition 2.23. A point x is adherent to A if and only if there exists a sequence in A converging to x .

Proof. Suppose that $x \in \overline{A}$. In view of the definition, for each positive integer n , there exists a point $x_n \in B(x, 1/n) \cap A$. Obviously, $\{x_n\}$ is the sequence of points of A converging to x . Conversely, suppose that $\{x_n\} \subset A$ and $x_n \rightarrow x$. Let $r > 0$. Then $d(x_n, x) < r$ for n greater than some k . Hence $x_n \in B(x, r) \cap A$ and x is adherent to A . ■

Example 2.24. Let $\overline{B}(x, r)$ be a closed ball in X . Then it is a closed set in X . To see this we have to show that all adherent points of $\overline{B}(x, r)$ are contained in $\overline{B}(x, r)$. If y is adherent to $\overline{B}(x, r)$, then $y_n \rightarrow y$ for some sequence $\{y_n\} \subset \overline{B}(x, r)$. Since

$$d(y, x) \leq d(y, y_n) + d(y_n, x) \leq d(y, y_n) + r \rightarrow r,$$

it follows that $y \in \overline{B}(x, r)$ as required.

Example 2.25. The closure of an open ball $B(x, r)$ does not have to coincide with a closed ball $\overline{B}(x, r)$. Indeed, consider $X = \mathbb{R} \setminus (0, 1)$ with the usual metric, $d(x, y) = |x - y|$. Then

$$\overline{B(0, 1)} = [-1, 0] \quad \text{but} \quad \overline{B}(0, 1) = [-1, 0] \cap \{1\}.$$

Example 2.26. A subset of metric space may be neither open nor closed. For instance, $[0, 1)$ is neither open nor closed in \mathbb{R} . The same is true for \mathbb{Q} . On the other hand, a subset may be open and at the same time closed. In a metric space equipped with the discrete metric any subset is both open and closed.

The relation between interior and adherent points is given in the next proposition.

Proposition 2.27. *A point $x \in X$ is an adherent point of A if and only if x is not an interior point of A^c .*

Proof. Assume that x is adherent to A . Then for every open ball at x , $B(x, r) \cap A \neq \emptyset$. Hence there is no open ball $B(x, r)$ contained in A^c which means that $x \notin (A^c)^\circ$. Conversely, assume that $x \notin (A^c)^\circ$. Hence there is no open ball $B(x, r)$ contained in A^c . Hence $B(x, r) \cap A \neq \emptyset$, for all $r > 0$, which means that x is adherent to A . ■

As a corollary we obtain.

Corollary 2.28. *If $A \subset X$, then*

$$X \setminus \overline{A} = (X \setminus A)^\circ \quad \text{and} \quad X \setminus A^\circ = \overline{X \setminus A}.$$

A set A is closed if and only if $X \setminus A$ is open, and A is open if and only if $X \setminus A$ is closed.

Theorem 2.29 (Properties of Interiors and Closures).

$$\begin{array}{ll} (a) A^\circ \subset A & (a') A \subset \overline{A} \\ (b) (A^\circ)^\circ = A^\circ & (b') \overline{\overline{A}} = \overline{A} \\ (c) A \subset B \implies A^\circ \subset B^\circ & (c') A \subset B \implies \overline{A} \subset \overline{B} \\ (d) (A \cap B)^\circ = A^\circ \cap B^\circ & (d') \overline{A \cup B} = \overline{A} \cup \overline{B} \\ (e) \bigcup_{i \in I} A_i^\circ \subset \left(\bigcup_{i \in I} A_i \right)^\circ & (e') \overline{\bigcap_{i \in I} A_i} \subset \bigcap_{i \in I} \overline{A_i} \\ (f) \left(\bigcap_{i \in I} A_i \right)^\circ \subset \bigcap_{i \in I} A_i^\circ & (f') \bigcup_{i \in I} \overline{A_i} \subset \overline{\bigcup_{i \in I} A_i} \end{array}$$

Proof. (a) follows immediately from the definition of the interior point. To see (b) note that A° is an open. (c): If $A \subset B$ and $x \in A^\circ$, then $B(x, r) \subset A \subset B$. So $x \in B^\circ$. (d): Note that $(A \cap B)^\circ \subset A^\circ$ and $(A \cap B)^\circ \subset B^\circ$ so $(A \cap B)^\circ \subset A^\circ \cap B^\circ$. On the other hand $A^\circ \cap B^\circ$ is open and contained in $A \cap B$, so by (b), $A^\circ \cap B^\circ \subset (A \cap B)^\circ$. Proofs of (e) and (f) are left as exercises. Proofs of (a')-(f') follows from the corresponding statements for interiors by taking complements. ■

Theorem 2.30 (Properties of open and closed sets).

- | | |
|--|---|
| <p>(a) \emptyset and X are open</p> <p>(b) $\{A_i\}_{i \in I}$ open $\Rightarrow \bigcup_{i \in I} A_i$ open</p> <p>(c) $\{A_i\}_{i=1}^m$ open $\Rightarrow \bigcap_{i=1}^m A_i$ open</p> <p>(d) A° = the largest open set contained in A</p> | <p>(a') \emptyset and X are closed</p> <p>(b') $\{A_i\}_{i \in I}$ closed $\Rightarrow \bigcap_{i \in I} A_i$ closed</p> <p>(c') $\{A_i\}_{i=1}^m$ closed $\Rightarrow \bigcup_{i=1}^m A_i$ is closed</p> <p>(d') \overline{A} = the smallest closed set containing A</p> |
|--|---|

Proof. The parts (a) and (a') are obvious.

(b) Let $x \in \bigcup_{i \in I} A_i$. Choose an index $j \in I$ so that $x \in A_j$. Since A_j is open, $B(x, r) \subset A_j \subset \bigcup_{i \in I} A_i$. Hence any point $x \in \bigcup_{i \in I} A_i$ is an interior point, and so $\bigcup_{i \in I} A_i$ is open.

(c) Assume $A_i \subset X, i = 1, \dots, m$ are open subsets of X , and let $x \in \bigcap_{i=1}^m A_i$. Then $x \in A_i$ for $i = 1, \dots, m$. Since the sets A_i are open, $B(x, r_i) \subset A_i$ for some $r_i > 0$. Take $r = \min\{r_1, \dots, r_m\}$. Then $B(x, r) \subset \bigcap_{i=1}^m A_i$, and the sets $\bigcap_{i=1}^m A_i$ is open.

(d) is left as an exercise. (a')-(d') are obtained from corresponding statements for open sets by taking complements and applying Corollary 2.28 ■

Theorem 2.31. *Let Y be a subspace of X .*

- (a) $B \subset Y$ is open in Y if and only if $B = Y \cap A$ for some open set A in X .
- (b) $B \subset Y$ is closed in Y if and only if $B = Y \cap F$, where F is closed in X .

Proof.

(a) Assume first that $B = Y \cap A$ for some open set A in X . Take $x \in B$. Then there exists an open ball $B(x, r)$ in X such that $B(x, r) \subset A$. But then $Y \cap B(x, r) \subset Y \cap A = B$. Since the open ball in the subspace Y with centre $x \in X$ and radius $r > 0$ is the intersection $Y \cap B(x, r)$, the set B is open in Y . Conversely, suppose that B is an open subset of the subspace Y . Then for every $x \in B$ there exists r_x such that the open ball $B(x, r_x) \cap Y$ in Y is contained in B . Then the open subset $A = \bigcup_{x \in B} B(x, r_x)$ of X satisfies $Y \cap A \subset B$. Since any $x \in B$ also belongs to A , $Y \cap A = B$ as required.

(b) A set B is closed in Y if and only if $Y \setminus B$ is open in Y . Hence if and only if $Y \setminus B = Y \cap A$ for some open subset A of X . Let $F = X \setminus A$. Then

F is closed in X and $B = Y \setminus [Y \cap A] = Y \setminus A = Y \cap [X \setminus A] = X \cap F$ as required. ■

Theorem 2.32. *Let X be the product of metric spaces (X_i, d_i) , $1 \leq i \leq m$.*

- (a) *If A_i is open in X_i , $1 \leq i \leq m$, then the product $A = \prod_{i=1}^n A_i$ is an open subset of $X = \prod_{i=1}^m X_i$.*
- (b) *If F_i is closed in X_i , $1 \leq i \leq m$, then $F = \prod_{i=1}^m F_i$ is closed in the product $X = \prod_{i=1}^m X_i$.*

Proof.

(a) We prove the result for the product of two metric spaces X_1 and X_2 . Let $a = (a_1, a_2) \in A \subset X$. Since A_i is open in X_i , there exists r_i such that an open ball $B(a_i, r_i)$ in X_i is contained in A_i . Let $r = \min\{r_1, r_2\}$. We claim that $B(a, r) \subset A$. Indeed, if $x = (x_1, x_2) \in B(a, r)$, then $d(a, x) < r$ where $x = (x_1, x_2)$, and since $d_i(a_i, x_i) < d(a, x) < r \leq r_i$ we conclude that $x_i \in B(a_i, r_i)$. Hence $x_i \in A_i$, $i = 1, 2$, so that $x \in A$.

(b) The proof follows from Proposition 2.14 ■

Definition 2.33. *The **boundary** of A in X , denoted by ∂A , is the set $\overline{A} \cap \overline{X \setminus A}$.*

Hence $x \in \partial A$ if for any $r > 0$ an open ball $B(x, r)$ intersects A and $X \setminus A$ as well. Clearly, the boundary is a closed set as an intersection of closed sets.

Example 2.34. Consider \mathbb{R} with the usual metric. Then

$$\begin{aligned}\partial([0, 1]) &= \partial((0, 1)) = \{0, 1\} \\ \partial(\mathbb{Q}) &= \partial(\mathbb{R} \setminus \mathbb{Q}) = \mathbb{R}.\end{aligned}$$

We shall show the last equality. Fix $x \in \mathbb{R}$. If $x \in \mathbb{Q}$, then

$$x \neq x + \frac{1}{n} \in \mathbb{Q} \quad \text{and} \quad x \neq x + \frac{\sqrt{2}}{n} \in \mathbb{Q}^c.$$

Since

$$x = \lim_n (x + 1/n) = \lim_n (x + \sqrt{2}/n),$$

it follows that $x \in \overline{\mathbb{Q}} \cap \overline{\mathbb{Q}^c}$. So $\mathbb{Q} \subset \overline{\mathbb{Q}} \cap \overline{\mathbb{Q}^c} = \partial \mathbb{Q}$. If $x \in \mathbb{Q}^c$, then $x + 1/n \in \mathbb{Q}$ and there exists a sequence of rational numbers x_n such that

$$x = \lim (x + 1/n) = \lim x_n.$$

Hence $x \in \overline{\mathbb{Q}} \cap \overline{\mathbb{Q}^c}$ and $\partial(\mathbb{Q}) = \partial(\mathbb{Q}^c) = \overline{\mathbb{Q}} \cap \overline{\mathbb{Q}^c} = \mathbb{R}$.

Definition 2.35. A point $x \in X$ is called **isolated** if $\{x\}$ is open. A space X is called **discrete** if all of its points are isolated.

If x is an isolated point, then for some $\varepsilon > 0$, an open ball $B(x, \varepsilon) \subset \{x\}$, that is, $B(x, \varepsilon) = \{x\}$ and if $y \neq x$, then $d(x, y) \geq \varepsilon$. Conversely, if $\inf\{d(x, y) \mid y \neq x\} > 0$, then $\{x\}$ is open. Note also that $\{x\}$ is always closed. For example, consider \mathbb{N} as subspace of \mathbb{R} . Then it is discrete. Also the space $J = \{1/n \mid n \in \mathbb{N}\}$ is discrete. In a discrete space any set is open, since it is a union of one-point sets which are open. Also any set is closed being a complement of an open set. Finally, a space is discrete if and only if the only convergent sequences are those which are eventually constant (Prove this!).

Definition 2.36. A subset A of a metric space is **dense** if $\overline{A} = X$.

Example 2.37. The sets \mathbb{Q} and \mathbb{Q}^c are dense in \mathbb{R} with the usual metric.

Proposition 2.38. Let X be a metric space and $A \subset X$. Then A is dense if and only if for every non-empty open set U of X , the intersection $U \cap A \neq \emptyset$.

Definition 2.39. A subset A of X is called **nowhere dense** if $(\overline{A})^\circ = \emptyset$.

Example 2.40. The sets of all natural numbers \mathbb{N} or all integers \mathbb{Z} are nowhere dense in \mathbb{R} with the usual metric. The set of real numbers \mathbb{R} is nowhere dense in \mathbb{R}^2 with the standard metric.

Example 2.41. [Cantor set] The Cantor set is subset of $[0, 1]$ constructed as follows:

Consider the interval $C_0 = [0, 1]$. At the first step divide C_0 into three equal intervals $[0, 1/3]$, $[1/3, 2/3]$ and $[2/3, 1]$ and remove the middle open interval $(1/3, 2/3)$. Denote the remaining intervals by $C_1 = [0, 1/3] \cup [2/3, 1]$. The length of intervals which constitute C_1 is equal to $2/3$. In the second step we perform the same operations as in the first step on each of the intervals of C_1 . We remove intervals $(1/9, 2/9)$ and $(7/9, 8/9)$. Denote the four remaining intervals by C_2 . Having finished the step $(n - 1)$, we perform the n th step and obtain the set C_n consisting of 2^n intervals.

Each of the sets C_n is closed and bounded, and $C_{n+1} \subset C_n$. The Cantor set is defined as

$$C = \bigcap_{n=1}^{\infty} C_n$$

It is non-empty and since for every n , C_n is closed, C is closed. The set C does not contain any interval (show this!), and so, C has empty interior. Hence C is nowhere dense.

3 Continuity

The definition of continuity is the $\varepsilon - \delta$ definition of calculus.

Definition 3.1. Let (X, d) and (Y, ρ) be metric spaces and let $f : X \rightarrow Y$ be a function. The function f is said to be **continuous at the point** $x_0 \in X$ if the following holds: for every $\varepsilon > 0$, there exists $\delta > 0$ such that for all $x \in X$ if $d(x, x_0) < \delta$, then $\rho(f(x), f(x_0)) < \varepsilon$. The function f is said to be **continuous** if it is continuous at each point of X .

The following proposition rephrases the definition in terms of open balls.

Proposition 3.2. Let $f : X \rightarrow Y$ be a function from a metric space X to another metric space Y and let $x_0 \in X$. Then f is continuous at x_0 if and only if for every $\varepsilon > 0$ there exists $\delta > 0$ such that

$$f(B(x_0, \delta)) \subset B(f(x_0), \varepsilon).$$

Theorem 3.3. Let $f : X \rightarrow Y$ be a function from a metric space (X, d) to another metric space (Y, ρ) and let $x_0 \in X$. Then f is continuous at x_0 if and only if for every sequence $\{x_n\}$ such that $x_n \rightarrow x_0$, $f(x_n) \rightarrow f(x_0)$. And f is continuous if and only if for every convergent sequence $\{x_n\}$ in X ,

$$\lim_n f(x_n) = f(\lim_n x_n).$$

Proof. Suppose that f is continuous at x_0 and let $x_n \rightarrow x_0$. We will prove that $f(x_n) \rightarrow f(x_0)$. Let $\varepsilon > 0$ be given. By the definition of continuity at x_0 , there exists $\delta > 0$ such that for all $x \in X$,

$$\text{if } d(x, x_0) < \delta, \text{ then } \rho(f(x), f(x_0)) < \varepsilon. \quad (3)$$

Since $x_n \rightarrow x_0$, there exists an integer k such that for all $n \geq k$,

$$d(x_n, x_0) < \delta. \quad (4)$$

Combining (3) and (4), we get

$$\rho(f(x_n), f(x_0)) < \varepsilon \quad \text{for all } n \geq k. \quad (5)$$

Hence $f(x_n) \rightarrow f(x_0)$ as required. Conversely, arguing by contradiction assume that f is not continuous at x_0 . To obtain a contradiction we will construct a sequence $\{x_n\}$ such that $x_n \rightarrow x_0$ but the sequence $\{f(x_n)\}$ does not converge to $f(x_0)$. Since f is not continuous at x_0 , there is positive $\varepsilon > 0$ such that for all $\delta > 0$ there exists x satisfying $d(x, x_0) < \delta$ but $\rho(f(x), f(x_0)) \geq \varepsilon$. For each n , take $\delta = 1/n$ and then choose x_n so that $d(x_n, x_0) < 1/n$ but $\rho(f(x_n), f(x_0)) \geq \varepsilon$. Hence $x_n \rightarrow x_0$ but the sequence $\{f(x_n)\}$ does not converge to $f(x_0)$. The second part of the theorem is an immediate consequence of the first. ■

Global continuity has a simple formulation in terms of open and closed sets.

Theorem 3.4. *Let f be a function from a metric space (X, d) to (Y, ρ) . Then f is continuous if and only if for every open set $U \in Y$, $f^{-1}(U)$ is open in X .*

Proof. Suppose first that f is continuous and U is open in Y . If $x \in f^{-1}(U)$, then $f(x) \in U$. Since U is open in Y and $f(x) \in U$, there exists a positive number ε such that $B(f(x), \varepsilon) \subset U$. In view of Proposition 3.2, there exists $\delta > 0$ such that $f(B(x, \delta)) \subset B(f(x), \varepsilon)$. Hence $B(x, \delta) \subset f^{-1}(f(B(x, \delta))) \subset f^{-1}(U)$, so $f^{-1}(U)$ is open in X . Conversely, suppose that $f^{-1}(U)$ is open in X for every open set U in Y . Let $x \in X$ and let $\varepsilon > 0$ be given. Since $B(f(x), \varepsilon)$ is open in Y , the set $f^{-1}(B(f(x), \varepsilon))$ is open in X . Since $x \in f^{-1}(B(f(x), \varepsilon))$, there exists $\delta > 0$ such that $B(x, \delta) \subset f^{-1}(B(f(x), \varepsilon))$. This implies that $f(B(x, \delta)) \subset B(f(x), \varepsilon)$, and in view of Proposition 3.2, f is continuous. ■

Theorem 3.5. *Let f be a function from (X, d) to (Y, ρ) . Then f is continuous if and only if for every closed set $F \subset Y$, $f^{-1}(F)$ is closed in X .*

The proof is left as an exercise.

Theorem 3.6. *Let X , Y and Z be three metric spaces.*

- (a) *If $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are continuous, then the composition $g \circ f$ is continuous.*
- (b) *If $f : X \rightarrow Y$ is continuous, and A is a subspace of X , then the restriction of f to A , $f|_A : A \rightarrow Y$, is continuous.*

Proof. (a) Let $x_n \rightarrow x_0$. Since f is continuous at x_0 , $f(x_n) \rightarrow f(x_0)$. Since g is continuous at $f(x_0)$, $g(f(x_n)) \rightarrow g(f(x_0))$. Hence $g \circ f(x_n) \rightarrow g \circ f(x_0)$. The second statement follows from the first. Here is another proof of the

second statement. Let U be an open subset of Z . Since g is continuous, $g^{-1}(U)$ is open in Y , and since f is continuous, $f^{-1}(g^{-1}(U))$ is open in X . But $f^{-1}(g^{-1}(U)) = (g \circ f)^{-1}(U)$ and so, $(g \circ f)^{-1}(U)$ is open in X . Hence $g \circ f$ is continuous.

(b) Note that $f|_A = f \circ j$, where $j : A \rightarrow X$ is the inclusion, i.e., defined by $j(x) = x$ for $x \in A$. Since for any open set U in X , $j^{-1}(U) = U \cap A$ which is open in A , it follows that j is continuous. So (b) follows from (a). ■

Theorem 3.7. *Let (X, d) , (Y_1, ρ_1) and (Y_2, ρ_2) be metric spaces. Let f be a function from X to Y_1 and g a function from X to Y_2 . Define the function h from X to the product $Y_1 \times Y_2$ by*

$$h(x) = (f(x), g(x)), \quad \text{for } x \in X.$$

Then h is continuous at x_0 if and only if f and g are continuous at x_0 . And h is continuous if and only if both functions f and g are continuous.

The similar statement about functions from the direct product does not hold in general. Suppose that f is a function from $X \times Y$ to Z . It may happen that f is discontinuous, though the maps $x \mapsto f(x, y)$ for every $y \in Y$ and $y \mapsto f(x, y)$ for every $x \in X$ are all continuous. For example, consider a function from $\mathbb{R} \times \mathbb{R}$ to \mathbb{R} defined by

$$f(x, y) = \begin{cases} \frac{xy}{x^2 + y^2} & \text{for } (x, y) \neq (0, 0); \\ 0 & \text{for } (x, y) = (0, 0). \end{cases}$$

The function f is discontinuous at $(0, 0)$ but all the functions $x \mapsto f(x, y)$ and $y \mapsto f(x, y)$ are continuous.

Theorem 3.8 (The pasting lemma). *Let $X = A \cup B$, where A and B are closed subspaces of X . Let $f : A \rightarrow Y$ and $g : B \rightarrow Y$ be continuous. If $f(x) = g(x)$ for all $x \in A \cap B$, then the function $h : X \rightarrow Y$ defined by*

$$h(x) = \begin{cases} f(x) & \text{if } x \in A; \\ g(x) & \text{if } x \in B \end{cases}$$

is continuous.

Proof. Let C be a closed subset of Y . Then $h^{-1}(C) = f^{-1}(C) \cup g^{-1}(C)$. Since f is continuous, $f^{-1}(C)$ is closed in A . But since A is closed, $f^{-1}(C)$ is closed in X . Similarly, $g^{-1}(C)$ is closed in X . So $h^{-1}(C)$ is closed in X and the proof is finished. ■

Uniform Continuity and Uniform Convergence

Definition 3.9. A mapping f from a metric space (X, d) to a metric space (Y, ρ) is said to be **uniformly continuous**, if for every $\varepsilon > 0$, there exists $\delta > 0$ such that $\rho(f(x), f(y)) < \varepsilon$ for all $x, y \in X$ satisfying $d(x, y) < \delta$.

Obviously, a uniformly continuous function is continuous.

Example 3.10. The function $f(x) = x/(1 + x^2)$ from \mathbb{R} to \mathbb{R} is uniformly continuous. To see this observe that for any $x < y$, in view of the mean value theorem of calculus, there exists $t \in (0, 1)$ such that

$$|f(x) - f(y)| = |f'(t)| \cdot |x - y| = \left| \frac{1 - t^2}{(1 + t^2)^2} \right| \cdot |x - y| \leq |x - y|.$$

since $|f'(t)| \leq 1$. Hence for given ε , choose $\delta = \varepsilon$. Then for any x, y such that $d(x, y) = |x - y| < \delta$, we have

$$d(f(x), f(y)) = |f(x) - f(y)| \leq |x - y| = d(x, y) < \delta = \varepsilon.$$

So f is uniformly continuous.

Example 3.11. The function $f(x) = x^2$ for $x \in \mathbb{R}$ is not uniformly continuous. Indeed, for a given $\delta > 0$ we can set

$$x = 1/\delta + \delta/2 \quad \text{and} \quad y = 1/\delta,$$

then $|x - y| = \delta/2 < \delta$ but $|x^2 - y^2| > 1$. However, if we consider the same function on some bounded interval, say $[-a, a]$, then the function is uniformly continuous since if $\delta < \varepsilon/2a$ and $x, y \in [-a, a]$ with $|x - y| < \delta$, then $|x^2 - y^2| = |x - y| \cdot |x + y| < 2a|x - y| < \varepsilon$.

Let (X, d) and (Y, ρ) be metric space. Consider a sequence $\{f_n\}$ of functions $f_n : X \rightarrow Y$ and let $f : X \rightarrow Y$.

Definition 3.12. The sequence $\{f_n\}$ is said to **converge pointwise** to f if for every $x \in X$ and for every $\varepsilon > 0$, there exists an index N such that

$$\rho(f_n(x), f(x)) < \varepsilon \quad \text{for all } n \geq N.$$

The sequence $\{f_n\}$ is said to **converge uniformly** to f if for every $\varepsilon > 0$, there exists an index N such that

$$\rho(f_n(x), f(x)) < \varepsilon \quad \text{for all } n \geq N \text{ and all } x \in X.$$

Equivalently, $\{f_n\}$ converges uniformly to f on X if

$$\sup\{\rho(f_n(x), f(x)) \mid x \in X\} \rightarrow 0.$$

The notion of uniform convergence of a sequence of functions is, in general, more useful than that of pointwise convergence.

Theorem 3.13. *Let $\{f_n\}$ be a sequence of continuous functions from a metric space (X, d) to a metric space (Y, ρ) . Suppose that $\{f_n\}$ converges uniformly to f from X to Y . Then f is continuous.*

In words, the uniform limit of continuous functions is continuous.

Proof. Let $x_0 \in X$ and let $\varepsilon > 0$ be given. Since $\{f_n\}$ converges uniformly to f , there exists an index N such that for all $n \geq N$ and all $x \in X$,

$$\rho(f_n(x), f(x)) < \varepsilon/3. \quad (6)$$

Since f_N is continuous at x_0 , we can choose $\delta > 0$ so that

$$\rho(f_N(x), f_N(x_0)) < \varepsilon/3 \quad (7)$$

for all $d(x, x_0) < \delta$. Now if $d(y, x_0) < \delta$, then

$$\rho(f(y), f(x_0)) \leq \rho(f(y), f_N(y)) + \rho(f_N(y), f_N(x_0)) + \rho(f_N(x_0), f(x_0)).$$

Each term of the right-hand side is less than $\varepsilon/3$, the first and the third in view of (6) and the second in view of (7). Thus

$$\rho(f(y), f(x_0)) < \varepsilon$$

for all $d(y, x_0) < \delta$. This proves that f is continuous. ■

4 Complete Spaces

Definition 4.1. *Let (X, d) be a given metric space and let $\{x_n\}$ be a sequence of points of X . We say that $\{x_n\}$ is **Cauchy** (or satisfies the **Cauchy condition**) if for every $\varepsilon > 0$ there exists $k \in \mathbb{N}$ such that*

$$d(x_n, x_m) < \varepsilon \quad \text{for all } n, m \geq k.$$

Properties of Cauchy sequences are summarized in the following propositions.

Proposition 4.2. *If $\{x_n\}$ is a Cauchy sequence, then $\{x_n\}$ is bounded.*

Proof. Take $\varepsilon = 1$. Since $\{x_n\}$ is Cauchy, there exists an index k such that $d(x_n, x_k) < 1$ for all $n \geq k$. Let $R > 1$ be such that $d(x_i, x_k) < R$ for $1 \leq i \leq k-1$. Then $x_n \in B(x_k, R)$ for all n , so $\{x_n\}$ is bounded. ■

Proposition 4.3. *If $\{x_n\}$ is convergent, then $\{x_n\}$ is a Cauchy sequence.*

Proof. Assume that $x_n \rightarrow x$. Then for a given $\varepsilon > 0$ there exists $k \in \mathbb{N}$ such that $d(x_n, x) < \varepsilon/2$ for all $n \geq k$. Hence taking any $n, m \geq k$,

$$d(x_n, x_m) \leq d(x_n, x) + d(x, x_m) < \varepsilon/2 + \varepsilon/2 = \varepsilon.$$

So $\{x_n\}$ is Cauchy. ■

Proposition 4.4. *If $\{x_n\}$ is Cauchy and it contains a convergent subsequence, then $\{x_n\}$ converges.*

Proof. Assume that $\{x_n\}$ is Cauchy and $x_{k_n} \rightarrow x$. We will show that $x_n \rightarrow x$. Let $\varepsilon > 0$. Since $\{x_n\}$ is Cauchy, there exists k' such that $d(x_n, x_{k_n}) < \varepsilon/2$ for all $n \geq k'$. Also since $x_{k_n} \rightarrow x$, there exists k'' such that $d(x_{k_n}, x) < \varepsilon/2$ for all $n \geq k''$. Set $k = \max\{k', k''\}$. Then for $n \geq k$,

$$d(x_n, x) \leq d(x_n, x_{k_n}) + d(x_{k_n}, x) < \varepsilon/2 + \varepsilon/2 = \varepsilon$$

showing that $x_n \rightarrow x$. ■

A Cauchy sequence need not converge. For example, consider $\{1/n\}$ in the metric space $((0, 1), |\cdot|)$. Clearly, the sequence is Cauchy in $(0, 1)$ but does not converge to any point of the interval.

Definition 4.5. *A metric space (X, d) is called **complete** if every Cauchy sequence $\{x_n\}$ in X converges to some point of X . A subset A of X is called **complete** if A as a metric subspace of (X, d) is complete, that is, if every Cauchy sequence $\{x_n\}$ in A converges to a point in A .*

By the above example, not every metric space is complete; $(0, 1)$ with the usual metric is not complete.

Theorem 4.6. *The space \mathbb{R} with the usual metric is complete.*

Proof. Let $\{x_n\}$ be a Cauchy sequence in \mathbb{R} . Then it is bounded, say $|x_n| \leq M$. Set $y_n = \inf\{x_k \mid k \geq n\}$. Then $\{y_n\}$ is increasing and $y_n \leq M$ for all n . Hence $\{y_n\}$ converges, say to x (see Proposition 11.11 in Appendix). We

claim that also $x_n \rightarrow x$. To see this choose N so that $|x_n - x_m| < \varepsilon/2$ for $n, m \geq N$. In particular,

$$x_N - \varepsilon/2 < x_k < x_N + \varepsilon/2 \quad \text{for all } k \geq N.$$

Hence

$$x_N - \varepsilon/2 \leq y_n \leq x_N + \varepsilon/2 \quad \text{for all } n \geq N.$$

Let $n \rightarrow \infty$. Then

$$x_N - \varepsilon/2 \leq x \leq x_N + \varepsilon/2,$$

or equivalently, $|x_N - x| \leq \varepsilon/2$. Hence for $n \geq N$,

$$|x_n - x| \leq |x_n - x_N| + |x_N - x| < \varepsilon/2 + \varepsilon/2 = \varepsilon.$$

Thus $\{x_n\}$ converges to x . ■

A subspace of a complete metric space may not be complete. However, the following holds true.

Theorem 4.7. *If (X, d) is a complete metric space and Y is a closed subspace of X , then (Y, d) is complete.*

Proof. Let $\{x_n\}$ be a Cauchy sequence of points in Y . Then $\{x_n\}$ also satisfies the Cauchy condition in X , and since (X, d) is complete, there exists $x \in X$ such that $x_n \rightarrow x$. But Y is also closed, so $x \in Y$ showing that Y is complete. ■

Theorem 4.8. *If (X, d) is a metric space, $Y \subset X$ and (Y, d) is complete, then Y is closed.*

Proof. Let $\{x_n\}$ be a sequence of points in Y such that $x_n \rightarrow x$. We have to show that $x \in Y$. Since $\{x_n\}$ converges in X , it satisfies the Cauchy condition in X and so, it also satisfies the Cauchy condition in Y . Since (Y, d) is complete, it converges to some point in Y , say to $y \in Y$. Since any sequence can have at most one limit, $x = y$. So $x \in Y$ and Y is closed. ■

Theorem 4.9. *If (X_i, d_i) are complete metric spaces for $i = 1, \dots, m$, then the product (X, d) is a complete metric space.*

Proof. Let $x_n = (x_n^1, \dots, x_n^m)$ and $\{x_n\}$ be a Cauchy sequence in (X, d) . Then for a given $\varepsilon > 0$ there exists k such that $d(x_n, x_m) < \varepsilon$ for all $n, m \geq k$. Since

$$d_j(x_n^j, x_m^j) \leq d(x_n, x_m) < \varepsilon,$$

it follows that $\{x_n^j\}$ is Cauchy in (X_j, d_j) for $j = 1, \dots, m$. Since (X_j, d_j) is complete, for $j = 1, \dots, m$ there exists $x^j \in X_j$ such that $x_n^j \rightarrow x^j$. Then, in view of Proposition 2.14, $x_n \rightarrow x$, where $x = (x^1, \dots, x^m)$. ■

Let (X, d) and (Y, d') be metric spaces and let $C(X, Y)$ be the space of continuous and bounded functions $f : X \rightarrow Y$. If $Y = \mathbb{R}$, we abbreviate $C(X, \mathbb{R})$ by $C(X)$. Consider

$$\rho(f, g) := \sup\{d'(f(x), g(x)) \mid x \in X\}$$

for $f, g \in C(X, Y)$.

Theorem 4.10. *The space $(C(X, Y), \rho)$ is a complete metric space if (Y, d') is complete.*

Proof. The verification that ρ is a metric is left as an exercise. Suppose that Y is complete, and suppose that $\{f_n\}$ is a Cauchy sequence in $C(X, Y)$. Then for every $x \in X$,

$$d'(f_n(x), f_m(x)) \leq \rho(f_n, f_m)$$

so that $\{f_n(x)\}$ is a Cauchy sequence on Y . Hence there exists a point, denoted by $f(x) \in Y$, such that $d'(f_n(x), f(x)) \rightarrow 0$. In this way we obtain a function $f : X \rightarrow Y$ which associate with a point $x \in X$ a point which is the limit of $\{f_n(x)\}$. We must check that f is continuous and bounded, and that $\rho(f_n, f) \rightarrow 0$. Let $x \in X$, and $\varepsilon > 0$. Then there exists N such that $d'(f(x), f_N(x)) < \varepsilon/3$, and an open ball $B(x, \delta)$ such that $d'(f_N(x), f_N(y)) < \varepsilon/3$ for every $y \in B(x, \delta)$. It follows that for every $y \in B(x, \delta)$,

$$d'(f(x), f(y)) \leq d'(f(x), f_N(x)) + d'(f_N(x), f_N(y)) + d'(f_N(y), f(y)) < \varepsilon.$$

Hence f is continuous. Now given $\varepsilon > 0$, chose n_0 such that $\rho(f_n, f_m) < \varepsilon$ for all $n, m \geq n_0$. Then for every $x \in X$,

$$d'(f_n(x), f(x)) = \lim_{m \rightarrow \infty} d'(f_n(x), f_m(x)) \leq \varepsilon$$

for every $n \geq n_0$. This says that $\rho(f_n, f) \leq \varepsilon$ for $n \geq n_0$. It remains to show that f is bounded. Take $x, y \in X$ and let $N \in \mathbb{N}$ be such that

$$d'(f(x), f_N(x)) < 1/2 \quad \text{and} \quad d'(f(y), f_N(y)) < 1/2$$

Note that we can find such N since $\rho(f_n, f) \rightarrow 0$. Then

$$\begin{aligned} d'(f(x), f(y)) &\leq d'(f(x), f_N(x)) + d'(f_N(x), f_N(y)) + d'(f_N(y), f(y)) \\ &< 1 + d'(f_N(x), f_N(y)) \leq 1 + \text{diam} f_N(X). \end{aligned}$$

Since $x, y \in X$ were arbitrary, $\text{diam} f(X) \leq 1 + \text{diam} f_N(X)$. Hence f is bounded. The proof is completed. \blacksquare

Corollary 4.11. *The space $(C(X), \rho)$ is complete.*

Structure of complete metric spaces-Baire's theorem

Let (X, d) be a metric space. If U and V are open and dense, then $U \cap V$ is also open and dense. To see that $U \cap V$ is dense, we have to show that $O \cap U \cap V$ is non-empty for any open set O . Since U is dense, there is $u \in O \cap U$, and since $O \cap U$ is open, $B(u, r) \subset O \cap U$ for some $r > 0$. Since V is dense, $B(u, r) \cap V \neq \emptyset$ so that, $\emptyset \neq B(u, r) \cap V \subset O \cap U \cap V$. If U and V are assumed to be dense but not necessarily open, then the intersection $U \cap V$ does not have to be dense. For example, let U be the set of rational numbers and V the set of irrational numbers \mathbb{Q}^c . Then both sets are dense in \mathbb{R} with the usual metric, however, $U \cap V = \emptyset$. Consider, now a sequence of dense and open sets U_n . In general, the intersection $\bigcap_{n \geq 1} U_n$ may be empty. For example, consider (\mathbb{Q}, d) with the usual metric d . Let $\{q_n | n \in \mathbb{N}\}$ be enumeration of rational numbers, and let $U_n = \mathbb{Q} \setminus \{q_n\}$. Then each U_n is open since it is a complement of a closed set $\{q_n\}$, and is dense. However, $\bigcap_{n \geq 1} U_n = \bigcap_{n \geq 1} [\mathbb{Q} \setminus \{q_n\}] = \mathbb{Q} \setminus \bigcup_{n \geq 1} \{q_n\} = \emptyset$. The Baire theorem says that if (X, d) is complete, then $\bigcap_{n \geq 1} U_n$ is dense.

Theorem 4.12. *Let (X, d) be a complete metric space, and let $\{U_n\}$ be a sequence of open and dense subsets of X . Then $\bigcap_{n \geq 1} U_n$ is dense.*

Proof. It suffices to show that $B(x, r)$ contains a point belonging to $\bigcap_{n \geq 1} U_n$ for any open ball $B(x, r)$. Since U_1 is open and dense, $B(x, r) \cap U_1$ is non-empty and open. So, there exists an open ball $B(x_1, R)$ with $R < r$ such that $B(x_1, R) \subseteq B(x, r)$ and $B(x_1, R) \subseteq U_1$. Taking $r_1 < R$, we get that $\overline{B}(x_1, r_1) \subseteq B(x, r)$ and $\overline{B}(x_1, r_1) \subseteq U_1$. Similarly, since U_2 is open and dense, there exists x_2 and $r_2 < 1/2$ such that $\overline{B}(x_2, r_2) \subseteq \overline{B}(x_1, r_1) \cap U_2$. Continuing in this way we find a sequence of balls $\overline{B}(x_n, r_n)$ with $r_n < 1/n$ and $\overline{B}(x_{n+1}, r_{n+1}) \subseteq \overline{B}(x_n, r_n) \cap U_n$. We claim that $\{x_n\}$ is Cauchy. By construction, $\overline{B}_n(x_n, r_n) \subset \overline{B}_k(x_k, r_k)$ for all $n \geq k$. Given $\varepsilon > 0$ choose $k \in \mathbb{N}$ so that $1/k < \varepsilon/2$. Then, if $n, m \geq k$,

$$d(x_n, x_m) \leq d(x_n, x_k) + d(x_k, x_m) < 1/k + 1/k < \varepsilon.$$

Because (X, d) is complete, $\{x_n\}$ converges, say to y . The point y lies in all balls $\overline{B}(x_k, r_k)$ since $x_n \in \overline{B}(x_k, r_k)$ for all $n \geq k$ and $\overline{B}(x_k, r_k)$ is closed for all k , so that after taking a limit as $n \rightarrow \infty$, $y \in \overline{B}(x_k, r_k)$ for all k . In

particular, $y \in \overline{B}(x_1, r_1) \subseteq B(x, r)$ and $y \in \overline{B}(x_{n+1}, r_{n+1}) \subset U_n$ for all n . Consequently, $y \in B(x, r) \cap \bigcap_{n \geq 1} U_n$, and the proof is finished. ■

As a consequence we obtain the following theorem.

Theorem 4.13. *If (X, d) is a complete metric space and $\{F_n\}$ is a sequence of nowhere dense subsets of X , then $\bigcup F_n$ has empty interior.*

Proof. Arguing by contradiction assume that $\bigcup F_n$ has non-empty interior. So $B(x, r) \subseteq \bigcup F_n$ for some x and $r > 0$. Define $U_n = X \setminus \overline{F_n}$. Clearly, U_n is open and we claim that it is dense. Indeed, if for some open set V , we have $V \cap U_n = \emptyset$, then $V \subseteq X \setminus U_n = \overline{F_n}$ contradicting that $\overline{F_n}$ has empty interior. Consequently, in view of the above theorem, $\bigcap_{n \geq 1} U_n$ is dense. So $B(x, r) \cap \bigcap_{n \geq 1} U_n \neq \emptyset$. On the other hand, $B(x, r) \subseteq \bigcup F_n \subseteq \bigcup \overline{F_n}$ so that $\emptyset = B(x, r) \cap [X \setminus \bigcup_{n \geq 1} \overline{F_n}] = B(x, r) \cap \bigcap_{n \geq 1} [X \setminus \overline{F_n}] = B(x, r) \cap \bigcap_{n \geq 1} U_n$, contradiction. ■

Example 4.14. The metric space \mathbb{R} with the standard metric space cannot be written as a countable union of nowhere sets since it is complete. By contrast, \mathbb{Q} with the standard metric can be written as the union of one point sets $\{q_n\}$, where $\{q_n | n \in \mathbb{N}\}$ is an enumeration of \mathbb{Q} . Every one point set $\{q_n\}$ is closed in \mathbb{Q} and its interior is empty, so nowhere dense. This does not contradict Baire's theorem since \mathbb{Q} with the standard metric is not complete.

Applications

Theorem 4.15. *Let (X, d) be a complete metric space, and let $\{f_n\}$ be a sequence of continuous functions $f_n : X \rightarrow \mathbb{R}$. Assume that the sequence $\{f_n(x)\}$ is bounded for every $x \in X$. Then there exists a non-empty open set $U \subset X$ on which the sequence $\{f_n\}$ is bounded, that is, there is a constant M such that $|f_n(x)| \leq M$ for all $x \in U$ and all $n \in \mathbb{N}$.*

Proof. Since the function f_n is continuous, the set $f_n^{-1}([-m, m]) = \{x \in X \mid |f_n(x)| \leq m\}$ is closed for any pair of positive integers n and m . Thus,

$$E_m = \{x \in X \mid |f_n(x)| \leq m \text{ for all } n \in \mathbb{N}\} = \bigcap_n f_n^{-1}([-m, m])$$

is closed for every $m \in \mathbb{N}$. If x is any point in X , then $|f_n(x)| \leq k$ for some $k \in \mathbb{N}$ and all n because $\{f_n(x)\}$ is bounded. Hence $X = \bigcup_m E_m$. In view

of the Baire theorem, one of the sets E_m has non-empty interior, say E_m . Setting $U = E_m^\circ$ the conclusion follows. ■

Theorem 4.16. *There exists continuous function $f : [0, 1] \rightarrow \mathbb{R}$ which is not differentiable at every point $x \in [0, 1]$.*

Proof. Recall that f has a right-hand derivative at x , if

$$\lim_{h \rightarrow 0^+} [(f(x+h) - f(x))/h] \text{ exists.}$$

We denote this limit by $f'_+(x)$. In particular, if f is differentiable at $x \in [0, 1]$ then $f'_+(x)$ exists and is equal to $f'(x)$. Consider the complete metric space $C([0, 1], \mathbb{R})$ with a metric d given by

$$d(f, g) = \sup\{|f(x) - g(x)| | x \in [0, 1]\}.$$

Let

$$M = \{f \in C([0, 1], \mathbb{R}) \mid \text{exists } x \in [0, 1] \text{ such that } f'_+(x) \text{ exists}\}$$

and let M_m , for $m \geq 2$, be the set of all $f \in C([0, 1], \mathbb{R})$ for which exists some $x \in [0, 1 - 1/m]$ such that

$$|f(x+h) - f(x)| \leq m \cdot h \text{ for all } h \in [0, 1/m].$$

Claim 1: $M \subset \bigcup_{n \geq 2} M_n$. Let $f \in M$. Then there exists $x \in [0, 1]$ such that $f'_+(x)$ exists. We will show that $|f(x+h) - f(x)| \leq m \cdot h$ for some $m \in \mathbb{N}$ and all $0 \leq h \leq 1/m$. Since

$$\lim_{h \rightarrow 0^+} \frac{f(x+h) - f(x)}{h} = f'_+(x),$$

we have

$$\lim_{h \rightarrow 0^+} \left| \frac{f(x+h) - f(x)}{h} \right| = |f'_+(x)|. \quad (1)$$

Take an integer $k \geq 2$ such that $|f'_+(x)| \leq k$ and $x \in [0, 1 - 1/k]$. In view of (1), there exists $0 < \delta < 1/k$ such that

$$|f(x+h) - f(x)| \leq k \cdot h \text{ for all } 0 \leq h \leq \delta.$$

Since f is continuous on a closed and bounded interval, there is $C > 0$ such that $|f(x)| \leq C$ for all $x \in [0, 1]$ (this is proved in the section on

compactness). Let k' be any integer so that $2C/\delta < k'$. Then, for $\delta \leq h \leq 1$ such that $x + h \leq 1$,

$$|f(x+h) - f(x)| \leq |f(x+h)| + |f(x)| \leq 2C = \frac{2C}{\delta} \cdot \delta \leq \frac{2C}{\delta} \cdot h \leq k' \cdot h.$$

Taking $m = \max\{k, k'\}$, we have $x \in [0, 1 - 1/m]$ and $|f(x+h) - f(x)| \leq m \cdot h$ for all $h \in [0, 1/m]$, so that $f \in M$.

Claim 2: M_m is closed for all $m \geq 2$. To see this, take $f \in \overline{M_m}$. We will show that $f \in M_m$, that is, $|f(x+h) - f(x)| \leq m \cdot h$ for some $x \in [0, 1 - 1/m]$ and all $h \in [0, 1/m]$. There exists $(f_k) \subset M_m$ such that $d(f_k, f) = \sup\{|f_k(x) - f(x)| : x \in [0, 1]\} \rightarrow 0$ as $k \rightarrow \infty$. Since $f_k \in M_m$, there exists $x_k \in [0, 1 - 1/m]$ such that

$$|f_k(x_k + h) - f_k(x_k)| \leq m \cdot h \quad (2)$$

for all $h \in [0, 1/m]$. Since $\{x_k\} \subseteq [0, 1 - 1/m]$, there exists a subsequence which converges to some point $x \in [0, 1 - 1/m]$. Without loss of generality we may assume that $x_k \rightarrow x \in [0, 1 - 1/m]$. Hence, by the triangle inequality and by (2),

$$\begin{aligned} |f(x+h) - f(x)| &\leq |f(x+h) - f(x_k+h)| + |f(x_k+h) - f_k(x_k+h)| \\ &\quad + |f_k(x_k+h) - f_k(x_k)| + |f_k(x_k) - f_k(x)| + |f_k(x) - f(x)| \\ &\leq |f(x+h) - f(x_k+h)| + d(f_k, f) + m \cdot h \\ &\quad + |f_k(x_k) - f_k(x)| + d(f_k, f) \end{aligned}$$

for all $0 \leq h \leq 1/m$. Since $d(f_k, f) \rightarrow 0$, and $|f(x+h) - f(x_k+h)| \rightarrow 0$, and $|f(x) - f(x_k)| \rightarrow 0$, as $k \rightarrow \infty$, we get that

$$|f(x+h) - f(x)| \leq m \cdot h$$

for all $0 \leq h \leq 1/m$. Consequently, $f \in M_m$ and M_m is closed.

Claim 3: $M_m^\circ = \emptyset$. Let $f \in M_m$, and let $\varepsilon > 0$. Then there exists a piecewise linear function $g : [0, 1] \rightarrow \mathbb{R}$ such that $d(f, g) = \sup\{|f(x) - g(x)| : 0 \leq x \leq 1\} < \varepsilon$ and $|g'_+(x)| > m$ for all $x \in [0, 1]$. That is, $g \in B(f, \varepsilon)$ and $g \notin M_m$. (Here $B(f, \varepsilon)$ is a ball in $C([0, 1], \mathbb{R})$ with centre at f and radius ε). So $M_m^\circ = \emptyset$.

In view of the Baire's theorem, $C([0, 1], \mathbb{R}) \neq \bigcup_{m \geq 2} M_m$ since otherwise $\bigcup_{m \geq 2} M_m$ has non-empty interior. Hence there exists $f \in C([0, 1], \mathbb{R})$ so that $f \notin \bigcup_{m \geq 2} M_m$. Since $M \subseteq \bigcup_{m \geq 2} M_m$, $f \notin M$. Since M contains all functions which are differentiable at least one point in $[0, 1]$, f is not differentiable at any $x \in [0, 1]$ ■




Figure 2: The black curve is the graph of f and the grey curve is the graph of g .

Contraction mapping principle-Banach fixed point theorem

Let (X, d) be a metric space and let $f : X \rightarrow X$. A point $x \in X$ is a **fixed point** of f if $f(x) = x$. The solution of many classes of equations can be regarded as fixed points of appropriate functions. In this section we give conditions that guarantee the existence of fixed points of certain functions. A function $f : X \rightarrow X$ is called a **contraction** if there exists $\alpha \in (0, 1)$ such that

$$d(f(x), f(y)) \leq \alpha d(x, y) \quad (8)$$

for all $x, y \in X$.

Theorem 4.17 (Banach Fixed Point Theorem). *Let $f : X \rightarrow X$ be a contraction of a complete metric space. Then f has a unique fixed point p . For any $x \in X$, define $x_0 = x$ and $x_{n+1} = f(x_n)$ for $n \geq 0$. Then $x_n \rightarrow p$, and*

$$d(x, p) \leq \frac{d(x, f(x))}{1 - \alpha}. \quad (9)$$

Proof. We start with the uniqueness of the fixed point of f . Assume that $p \neq q$ and that $f(p) = p$ and $f(q) = q$. Then

$$d(p, q) = d(f(p), f(q)) \leq \alpha d(p, q)$$

so that $d(p, q) = 0$ since $\alpha \in (0, 1)$. So $p = q$, contradicting our assumption. Hence f has at most one fixed point. Fix any point $x \in X$, and let $x_0 = x$

and $x_{n+1} = f(x_n)$ for $n \geq 0$. Then for any n ,

$$d(x_{n+1}, x_n) = d(f(x_n), f(x_{n-1})) \leq \alpha d(x_n, x_{n-1})$$

and,

$$d(x_{n+1}, x_n) \leq \alpha d(x_n, x_{n-1}) \leq \alpha^2 d(x_{n-1}, x_{n-2}) \leq \cdots \leq \alpha^n d(x_1, x_0).$$

For $m > n$,

$$\begin{aligned} d(x_m, x_n) &\leq d(x_n, x_{n+1}) + d(x_{n+1}, x_{n+2}) + \cdots + d(x_{m-1}, x_m) \\ &\leq (\alpha^n + \alpha^{n+1} + \cdots + \alpha^{m-1}) d(x_1, x_0) \leq \left(\sum_{i=n}^{\infty} \alpha^i \right) d(x_1, x_0) \\ &= \alpha^n \left(\sum_{i=0}^{\infty} \alpha^i \right) d(x_1, x_0) = \frac{\alpha^n d(x_1, x_0)}{1 - \alpha}. \end{aligned}$$

Since $\alpha^n \rightarrow 0$ as $n \rightarrow \infty$ (recall $\alpha \in (0, 1)$), the sequence $\{x_n\}$ is Cauchy in X . Since (X, d) is complete, there exists $p \in X$ such that $x_n \rightarrow p$. Taking a limit $m \rightarrow \infty$ in the last inequality we find that

$$d(p, x_n) \leq \frac{\alpha^n d(x_1, x_0)}{1 - \alpha}. \quad (10)$$

Thus,

$$\begin{aligned} d(f(p), p) &\leq d(f(p), x_{n+1}) + d(x_{n+1}, p) = d(f(p), f(x_n)) + d(x_{n+1}, p) \\ &\leq d(p, x_n) + d(x_{n+1}, p) \leq \frac{\alpha^n d(x_1, x_0)}{1 - \alpha} + \frac{\alpha^{n+1} d(x_1, x_0)}{1 - \alpha} \\ &= \alpha^n \cdot \frac{(1 + \alpha) d(x_1, x_0)}{1 - \alpha} \rightarrow 0, \end{aligned}$$

and therefore $p = f(p)$. The inequality (9) follows from (10) by taking $n = 0$. ■

Here is an application of Banach fixed point theorem to the local existence of solutions of ordinary differential equations.

Theorem 4.18 (Picard's Theorem). *Let U be an open subset of \mathbb{R}^2 and let $f : U \rightarrow \mathbb{R}$ be a continuous function which satisfies the Lipschitz condition with respect to the second variable, that is,*

$$|f(x, y_1) - f(x, y_2)| \leq \alpha |y_1 - y_2|$$

for all $(x, y_1), (x, y_2) \in U$, and some $\alpha > 0$. Then for a given $(x_0, y_0) \in U$ there is $\delta > 0$ so that the differential equation

$$y'(x) = f(x, y(x))$$

has a unique solution $y : [x_0 - \delta, x_0 + \delta] \rightarrow \mathbb{R}$ such that $y(x_0) = y_0$.

Proof. Note that it is enough to show that there are $\delta > 0$ and a unique function $y : [x_0 - \delta, x_0 + \delta] \rightarrow \mathbb{R}$ such that

$$y(x) = y_0 + \int_{x_0}^x f(t, y(t)) dt.$$

Fix $(x_0, y_0) \in U$, then there exists $\delta > 0$ and $b > 0$ such that if $I = [x_0 - \delta, x_0 + \delta]$ and $J = [y_0 - b, y_0 + b]$, then $I \times J \subset U$. Since f is continuous and $I \times J$ is closed and bounded, f is bounded on $I \times J$. That is, $|f(x, y)| \leq M$ for some M and all $(x, y) \in U$. Take δ smaller so that $\alpha\delta < 1$ and $\alpha M < b$. Denote by X the set of all continuous functions $g : I \rightarrow J$. The set X with the metric $\rho(g, h) = \sup\{|g(x) - h(x)|, x \in I\}$ is a complete metric space. For $g \in X$, let

$$(Tg)(x) = y_0 + \int_{x_0}^x f(t, g(t)) dt.$$

Then $Tg : I \rightarrow \mathbb{R}$ is continuous since if $x_1, x_2 \in I$ and $x_2 > x_1$, then

$$|(Tg)(x_2) - (Tg)(x_1)| = \left| \int_{x_1}^{x_2} f(t, g(t)) dt \right| \leq \int_{x_1}^{x_2} |f(t, g(t))| dt \leq M|x_2 - x_1|.$$

For $x_0 \leq x \leq x_0 + \delta$,

$$|(Tg)(x) - y_0| = \left| \int_{x_0}^x f(t, g(t)) dt \right| \leq \int_{x_0}^x |f(t, g(t))| dt \leq M|x - x_0| < M\delta < b$$

The same inequality holds for $x_0 - \delta \leq x \leq x_0$, and so $Tg \in X$ for any $g \in X$. Since f is Lipschitz with respect to the second variable, we obtain for $g, h \in X$ and $x \in [x_0, x_0 + \delta]$,

$$\begin{aligned} |(Tg)(x) - (Th)(x)| &= \left| \int_{x_0}^x [f(t, g(t)) - f(t, h(t))] dt \right| \\ &\leq \int_{x_0}^x |f(t, g(t)) - f(t, h(t))| dt \\ &\leq \alpha|x - x_0|d(g, h) < \alpha\delta d(g, h). \end{aligned}$$

Similarly, $|(Tg)(x) - (Th)(x)| \leq \alpha|x - x_0|d(g, h) < \alpha\delta d(g, h)$ for $x \in [x_0 - \delta, x_0]$. Since $\alpha\delta < 1$, T is a contraction and in view of Banach's fixed point theorem there exists a unique continuous function $y : I \rightarrow J$ such that

$$y(x) = (Ty)(x) = y_0 + \int_{x_0}^x f(t, y(t))dt.$$

■

Completions

The space $(0, 1)$ with the usual metric is not complete but is a subspace of the complete metric space $[0, 1]$ with the usual metric. This example illustrates the general situation: every metric space X may be regarded as a subspace of a complete metric space \tilde{X} in such a way that $\overline{X} = \tilde{X}$.

We will need the following concept.

Definition 4.19. A bijective map f from (X, d) onto (Y, ρ) is called an **isometry** if

$$\rho(f(x), f(y)) = d(x, y) \quad \text{for all } x, y \in X.$$

If $f : X \rightarrow Y$ is an isometry, then $f^{-1} : Y \rightarrow X$ is also an isometry, and the spaces (X, d) and (Y, ρ) are called **isometric**. Two isometric spaces can be regarded as indistinguishable for all practical purposes that involve only distance.

Definition 4.20. A **completion** of a metric space (X, d) is a pair consisting of a complete metric space (\tilde{X}, \tilde{d}) and an isometry $\varphi : X \rightarrow \varphi(X)$ such that $\varphi(X)$ is dense in \tilde{X} .

Theorem 4.21. Let (X, d) be a metric space. Then (X, d) has a completion. The completion is unique in the following sense: If $((X_1, d_1), \varphi_1)$ and $((X_2, d_2), \varphi_2)$ are completions of (X, d) , then (X_1, d_1) and (X_2, d_2) are isometric. That is, there exists an isometry $\varphi : X_1 \rightarrow X_2$ such that $\varphi \circ \varphi_1 = \varphi_2$.

Proof.

Existence: Let $B(X)$ be the space of bounded functions defined on X equipped with the uniform norm $\sigma(f, g) = \sup_{y \in X} |f(y) - g(y)|$. Fix a point $a \in X$. With every $x \in X$ we associate a function $f_x : X \rightarrow \mathbb{R}$ defined by

$$f_x(y) = d(y, x) - d(y, a), \quad y \in X.$$

We have

$$|f_x(y)| = |d(y, x) - d(y, a)| \leq d(x, a)$$

so that f_x is bounded. Since

$$|f_{x_1}(y) - f_{x_2}(y)| \leq d(x_1, x_2) \quad \text{for all } y \in X,$$

$\sigma(f_{x_1}, f_{x_2}) = \sup_{y \in X} \{|f_{x_1}(y) - f_{x_2}(y)|\} \leq d(x_1, x_2)$. On the other hand,

$$\sigma(f_{x_1}, f_{x_2}) \geq |f_{x_1}(x_2) - f_{x_2}(x_2)| = d(x_1, x_2).$$

Hence

$$\sigma(f_{x_1}, f_{x_2}) = d(x_1, x_2),$$

and the map $\varphi : X \rightarrow C(X, \mathbb{R})$ defined by $\varphi(x) = f_x$ is an isometry onto $\varphi(X)$,

$$\sigma(\varphi(x_1), \varphi(x_2)) = d(x_1, x_2).$$

Denote by X' the closure of $\varphi(X)$ in $B(X)$ and let d' be the metric on X' induced by σ . Since $(B(X), \sigma)$ is complete and X' is closed in $B(X)$, the space (X', d') is complete.

Uniqueness:

The isometry $\varphi_1 : X \rightarrow \varphi_1(X)$ has an inverse $\varphi_1^{-1} : \varphi_1(X) \rightarrow X$. Then $\varphi_2 \circ \varphi_1^{-1}$ is an isometry from $\varphi_1(X)$ onto X_2 . Since $\varphi_1(X)$ is dense in (X_1, d_1) , $\varphi_2 \circ \varphi_1^{-1}$ extends to the map $\varphi : X_1 \rightarrow X_2$ satisfying

$$d_2(\varphi(x), \varphi(y)) = d_1(x, y), \quad x, y \in X_1.$$

Since X_1 is complete, in view of the above equation, $\varphi(X_1)$ is closed in X_2 . Since $\varphi \circ \varphi_1 = \varphi_2$, $\varphi_2(X) \subset \varphi(X_1)$. This implies that $X_2 = \overline{\varphi_2(X)} \subseteq \overline{\varphi(X_1)} = \varphi(X_1)$ since $\varphi(X_1)$ is closed in X_2 . Consequently, $\varphi(X_1) = X_2$, i.e., φ is surjective and the proof is completed. ■

5 Compact Metric Spaces

We start with the classical theorem of Bolzano-Weierstrass.

Theorem 5.1 (Bolzano-Weierstrass). *Let I be a closed and bounded interval of \mathbb{R} , and let $\{x_n\}$ be a sequence in I . Then there exists a subsequence $\{x_{n_k}\}$ which converges to a point in I .*

Proof. Without loss of generality we may assume that $I = [0, 1]$. Bisect the interval $[0, 1]$ and consider the two intervals $[0, 1/2]$ and $[1/2, 1]$. One of these subintervals must contain x_n for infinitely many n . Call this subinterval I_1 . Now bisect I_1 . Again, one of the two subintervals contains x_n for infinitely many n . Denote this subinterval I_2 the interval containing x_n for infinitely many n . Proceeding in this way we find a sequence of closed intervals I_n , each one contained in the preceding one, each one half of the length of the preceding one, and each containing x_n for infinitely many n . Choose an integer n_1 so that $x_{n_1} \in I_1$. Then choose $n_2 > n_1$ such that $x_{n_2} \in I_2$. Then choose $n_3 > n_2$ such that $x_{n_3} \in I_3$, and so on. Continuing this way we choose we find a sequence $\{x_{n_k}\}$ such that $x_{n_k} \in I_k$. If $i, j \geq k$, then $x_{n_i}, x_{n_j} \in I_k$ and so

$$|x_{n_i} - x_{n_j}| \leq 1/2^k.$$

Hence $\{x_{n_k}\}$ is Cauchy and since $[0, 1]$ is complete, $\{x_{n_k}\}$ converges to a point in $[0, 1]$ ■

Definition 5.2. A metric space (X, d) is called **compact** if every sequence in X has a convergent subsequence. A subset Y of X is compact if every sequence in Y has a subsequence converging to a point in Y .

Proposition 5.3. Let (X, d) be compact and Y a closed subset of X . Then Y is compact.

Proof. Let $\{x_n\}$ be a sequence in Y . Since X is compact, the sequence $\{x_n\}$ has a converging subsequence, say $x_{n_k} \rightarrow x$. Since Y is closed, $x \in Y$. ■

Proposition 5.4. Let X be a metric space and Y a compact subset of X . Then Y is closed and bounded.

Proof. Take any $x \in \overline{Y}$. There exists a sequence $\{x_n\}$ in Y converging to x . Since Y is compact, the sequence $\{x_n\}$ has a converging subsequence, say $x_{n_k} \rightarrow y$ with $y \in Y$. In view of the uniqueness of the limit, $y = x$. Hence Y is closed. To see that Y is bounded, we argue by contradiction and construct a sequence $\{x_n\}$ which does not have a converging subsequence. Fix any point $y \in X$. For every $n \in \mathbb{N}$, there exists a point $x_n \in Y$ so that $d(x_n, y) \geq n$ since otherwise $Y \subseteq \overline{B}(y, n)$ for some n . The sequence $\{x_n\}$ contains converging subsequence since Y is compact. Say $x_{n_k} \rightarrow x \in Y$. Let $\varepsilon = d(x, y)$, Then $d(x_{n_k}, x) \leq 1$ for all $k \geq N$. Hence by the triangle inequality,

$$d(x, y) \geq d(y, x_{n_k}) - d(x, x_{n_k}) \geq n_k - 1 \geq k - 1$$

for all $k \geq N$, contradiction. Consequently, Y is bounded. ■

Combining Proposition 5.4 with Theorem 5.1 we get

Theorem 5.5. *A subset Y of \mathbb{R} is compact if and only if Y is bounded and closed.*

The result is also valid in \mathbb{R}^n with the standard metric. A subset of \mathbb{R}^n is compact if and only if it is bounded and closed. This follows from the fact that if A_i is a compact subset of (X_i, d_i) for $1 \leq i \leq n$, then $A_1 \times A_2 \times \cdots \times A_n$ is compact in the product space $X_1 \times X_2 \times \cdots \times X_n$. In particular, using Theorem 5.1, $[-a, a]^n$ is compact in \mathbb{R}^n . So if A is bounded and closed in \mathbb{R}^n , then A is a subset of a compact set $[-a, a]^n$, and then Proposition 5.3 implies that A is compact.

Theorem 5.5 does not hold true for general metric spaces.

Example 5.6. Consider the metric space $((C([0, 1], \mathbb{R}), d)$ consisting of all continuous functions on the interval $[0, 1]$ with the supremum metric $d(f, g) = \sup\{|f(x) - g(x)| : x \in [0, 1]\}$. Let $A = \{f_1, f_2, \dots\}$, where $f_i(x) = x^i$ for $x \in [0, 1]$. The set A is bounded since $B(0, 2)$. For $k > i$, we have

$$|f_k(x) - f_i(x)| = x^i \cdot |x^{k-i} - 1|.$$

Let i be fixed. Then for x close to 1, $x^i > 1/2$ and for k large $x^{k-i} < 1/2$. Hence

$$|f_k(x) - f_i(x)| = x^i \cdot |x^{k-i} - 1| > 1/4,$$

So $d(f_i, f_k) \geq 1/4$ for k large. Let $f \in \overline{A}$. We claim that $f \in A$. Indeed, there exists a sequence $\{g_k\} \subset A$ such that $d(g_k, f) \rightarrow 0$. Hence $\{g_k\}$ is Cauchy and there is N such that $d(g_N, g_k) < 1/4$ for all $k \geq N$. Since $g_k \in A$, $g_k = f_{n_k}$. Hence $d(f_{n_N}, f_{n_k}) < 1/4$ for all $k \geq N$. From $d(f_i, f_k) \geq 1/4$ for all k large, it follows that the set $\{n_k\}$ is bounded, that is, $n_k \leq m$ for some $m \in \mathbb{N}$ and all $k \in \mathbb{N}$. Hence for all k , $g_k \in \{f_1, f_2, f_3, \dots, f_m\}$ so that the sequence $\{g_k\}$ has a constant subsequence, say $g_{n_l} = f_i$ for some $i \leq m$ and all l . Since a subsequence of a convergent sequence converges to the same limit, the sequence $\{g_k\}$ converges to f_i , that is, $f = f_i$. Hence A is closed. To see that A is not compact, consider a sequence $\{f_n\}$. If A were compact, then a subsequence of $\{f_n\}$ converges to some $f_i \in A$. But then $d(f_i, f_{n_k}) < 1/4$ for large k contradicting $d(f_i, f_k) \geq 1/4$ for large k .

Theorem 5.7. *Let (X, d) and (Y, d') be metric spaces and let $f : X \rightarrow Y$ be continuous. If a subset $K \subseteq X$ is compact, then $f(K)$ is compact in (Y, d') . In particular, if (X, d) is compact, then $f(X)$ is compact in Y .*

Proof. Let $\{y_n\}$ be any sequence in $f(K)$, and let $\{x_n\}$ be a sequence in K of points such that $f(x_n) = y_n$. Since K is compact, $\{x_n\}$ has a converging subsequence to a point in K , say $x_{n_k} \rightarrow x$ with $x \in K$. Since f is continuous, $f(x_{n_k}) \rightarrow f(x)$. That is, $y_{n_k} \rightarrow f(x)$ and since $f(x) \in f(K)$, $f(K)$ is compact. ■

As a corollary we get

Corollary 5.8. *Let $f : X \rightarrow \mathbb{R}$ be a continuous function on a compact metric space. Then f attains a maximum and a minimum value, that is, there exist a and $b \in X$ such that $f(a) = \inf\{f(x) \mid x \in X\}$ and $f(b) = \sup\{f(x) \mid x \in X\}$.*

Proof. By Theorem 5.7, $f(X)$ is compact and so, it is bounded and the $\sup\{f(x) \mid x \in X\}$ is finite. Set $C = \sup\{f(x) \mid x \in X\}$. By definition of supremum, for every $n \in \mathbb{N}$, there exists x_n such that $C - 1/n \leq f(x_n) \leq C$. The sequence $\{x_n\}$ has a converging subsequence, $x_{n_k} \rightarrow b$ because X is compact. In view of the continuity of f , $f(x_{n_k}) \rightarrow f(b)$, and since $C - 1/n \leq f(x_n) \leq C$, $f(b) = C$. Similarly, $f(a) = \inf\{f(x) \mid x \in X\}$. ■

Theorem 5.9. *Suppose $f : (X, d) \rightarrow (Y, d')$ is a continuous mapping defined on a compact metric space X . Then f is uniformly continuous.*

Proof. Suppose not. Then there is some $\varepsilon > 0$ such that for all $\delta > 0$ there exist points x, y with $d(x, y) < \delta$ but $d'(f(x), f(y)) \geq \varepsilon$. Take $\delta = 1/n$ and let x_n, y_n be points such that $d(x_n, y_n) < 1/n$ but $d'(f(x_n), f(y_n)) \geq \varepsilon$. Compactness of X implies that there is a subsequence $\{x_{n_k}\}$ converging to some point $x \in X$. Since $d(x_{n_k}, y_{n_k}) < 1/n_k \rightarrow 0$ as $k \rightarrow \infty$, the sequence $\{y_{n_k}\}$ converges to the same point x . Continuity of f implies that the sequences $\{f(x_{n_k})\}, \{f(y_{n_k})\}$ converge to $f(x)$. Then $d'(f(x_{n_k}), f(x)) < \varepsilon/2$ and $d'(f(y_{n_k}), f(x)) < \varepsilon/2$ for k large, and so,

$$d'(f(x_{n_k}), f(y_{n_k})) \leq d'(f(x_{n_k}), f(x)) + d'(f(x), f(y_{n_k})) < \varepsilon$$

for k large, contradiction that $d'(f(x_n), f(y_n)) \geq \varepsilon$ for all n . ■

Characterization of Compactness for Metric Spaces

Definition 5.10. *Let (X, d) be a metric space and let $A \subseteq X$. If $\{U_i\}_{i \in I}$ is a family of subsets of X such that $A \subset \bigcup_{i \in I} U_i$, then it is called a **cover** of A , and A is said to be **covered** by the U_i 's. If each U_i is open, then $\{U_i\}_{i \in I}$ is an **open cover**. If $J \subset I$ and still $A \subseteq \bigcup_{i \in J} U_i$, then $\{U_i\}_{i \in J}$ is a **subcover**.*

Definition 5.11. Let (X, d) be a metric space and let $A \subseteq X$. Then A has the **Heine-Borel property** if for every open cover $\{U_i\}_{i \in I}$ of A , there is a finite set $F \subseteq I$ such that $A \subseteq \bigcup_{i \in F} U_i$.

Example 5.12. Consider a set X with a discrete metric. Then every one-point set is open and the collection of all one-point sets is an open cover of X . Clearly, this cover does not have any proper subcover. Hence, a discrete metric space X has the Heine-Borel property if and only if X consists of a finite number of points.

Definition 5.13. Let (X, d) be a metric space and $A \subseteq X$. Let $\varepsilon > 0$. A finite subset S is called an ε -**net** for A if $A \subseteq \bigcup_{x \in S} B(x, \varepsilon)$. A set A is called **totally bounded** if, for every $\varepsilon > 0$, there is an ε -net for A . That is, for every $\varepsilon > 0$, there is a finite set S such that $A \subseteq \bigcup_{x \in S} B(x, \varepsilon)$.

Every totally bounded set is bounded, for if $x, y \in \bigcup_{i=1}^n B(x_i, \varepsilon)$, say $x \in B(x_1, \varepsilon), y \in B(x_2, \varepsilon)$, then

$$d(x, y) \leq d(x, x_1) + d(x_1, x_2) + d(x_2, y) \leq 2\varepsilon + \max\{d(x_i, x_j) \mid 1 \leq i, j \leq n\}.$$

The converse is in general false.

Example 5.14. Consider (\mathbb{R}, d) with $d(x, y) = \min\{|x - y|, 1\}$. Then (\mathbb{R}, d) is bounded since $d(x, y) \leq 1$ for all $x, y \in \mathbb{R}$. But (\mathbb{R}, d) is not totally bounded since it cannot be covered by a finite number of balls of radius $1/2$. Indeed, let S be any finite subset of \mathbb{R} , and let x be the largest number in S . If $y \in S$, then $d(x + 1, y) = \min\{|x + 1 - y|, 1\} = 1$ and so there is no $1/2$ -net for \mathbb{R} .

Theorem 5.15. Let A be a subset of a metric space (X, d) . Then the following conditions are equivalent:

- (a) A is compact.
- (b) A is complete and totally bounded.
- (c) A has the Heine-Borel property.

Proof. We will show that (a) implies (b), (b) implies (c), (c) implies (a).

(a) implies (b):

Let $\{x_n\}$ be a Cauchy sequence in A . We have to show that it converges to

a point in A . By compactness of A , some subsequence, $\{x_{n_k}\}$, converges to $x \in A$. Then $x_n \rightarrow x$. Indeed, let $\varepsilon > 0$. Choose n_0 such that $d(x_n, x_m) < \varepsilon/2$ for all $n, m \geq n_0$. Also choose k_0 such that $d(x_{n_k}, x) < \varepsilon/2$ for all $k \geq k_0$. Then if $k \geq k_0$, is such that $n_k \geq n_0$, then for $m \geq n_0$ we have,

$$d(x_m, x) \leq d(x_m, x_{n_k}) + d(x_{n_k}, x) < \varepsilon/2 + \varepsilon/2 = \varepsilon.$$

Hence we proved that A is complete.

Suppose that A is not totally bounded. Then there exists r so that A cannot be covered by finitely many balls of radius r . We construct a sequence $\{x_n\}$ in A which does not have a converging subsequence. Take any $x_1 \in A$. Since $B(x_1, r)$ does not cover A , there is a point in $A \setminus B(x_1, r)$. Call this point x_2 . Having chosen points x_1, \dots, x_n , we choose x_{n+1} so that it belongs to $A \setminus \bigcup_{i=1}^n B(x_i, r)$. This is possible since A is not covered by $B(x_1, r), \dots, B(x_n, r)$. Continuing in this way we get a sequence $\{x_n\}$ such that $d(x_n, x_m) \geq r$ for all n and m . Such a sequence cannot have a convergent subsequence since if $\{x_{n_k}\}$ converges, then it is Cauchy and $d(x_{n_k}, x_{n_m}) < r$ for large k and m . Hence A is not compact, contradiction.

(b) implies (c):

Let $\mathcal{U} = \{U_i\}_{i \in I}$ be a collection of open sets covering A . Arguing by contradiction we assume that \mathcal{U} does not contain a finite subcover. Total boundedness of A implies that there is a finite set of closed balls B_1, \dots, B_n of radius 1 which cover A . If each of the sets $A \cap B_i$ can be covered by a finite number of sets from \mathcal{U} , then A can also be covered by a finite subcollection of sets from \mathcal{U} . Therefore some $A \cap B_i$, denoted by B^1 , cannot be covered by a finite number of sets from \mathcal{U} . Since B^1 is a subset of A and A is totally bounded, B^1 is totally bounded. So let B_1^1, \dots, B_m^1 be a finite set of closed balls of radius $1/2$ which cover B^1 . If each $B_i^1 \cap B^1$ can be finitely covered by sets from \mathcal{U} , the same is true for B^1 . Therefore, some $B_j^1 \cap B^1$, denoted by B^2 , cannot be covered by a finite number of sets from \mathcal{U} . Continuing in this way we obtain a sequence of closed sets B^n such that $\dots \subset B^n \subset B^{n-1} \subset \dots \subset B^1$, none of which can be finitely covered and $\text{diam } B^n \leq 1/n$. From each B^n choose a point x_n . The sequence $\{x_n\}$ is Cauchy since for $n, m \geq k$, $x_n, x_m \in B_k$ and

$$d(x_n, x_m) \leq \text{diam } B^k \leq 1/k.$$

By completeness of A , the sequence $\{x_n\}$ converges, say $x_n \rightarrow x$. In fact, $x \in B^k$ for all k since $x_n \in B^k$ for all $n \geq k$ and since B^k is closed. In particular, $x \in A$. Since \mathcal{U} covers A , the point x belongs to some U_i , and

therefore, $B(x, \varepsilon) \subset U_i$ for some ε . If $y \in B^n$, then

$$d(x, y) \leq d(x, x_n) + d(x_n, y) \leq d(x, x_n) + \text{diam } B^n \leq d(x, x_n) + 1/n.$$

For large n , the right side is less than ε . So for large n , $B^n \subset B(x, \varepsilon)$. Hence $B^n \subset U_i$ which shows that B^n can be finitely covered by sets from \mathcal{U} . This contradiction shows that A has the Heine-Borel property.

(c) implies (a):

Suppose that A is not compact. Then there exists a sequence $\{x_n\}$ in A with no convergent subsequence in A . Then for every $x \in A$, there exists a ball $B(x, \varepsilon_x)$ which contains x_n for at most finitely many n . Otherwise, there exists x such that for every $r > 0$, $B(x, r)$ contains x_n for infinitely many n . Then, in particular, for every k , $B(x, 1/k)$ contains x_n for infinitely many n . Choose n_1 so that $x_{n_1} \in B(x, 1)$. Since $B(x, 1/2)$ contains x_n for infinitely many n , there is $n_2 > n_1$ such that $x_{n_2} \in B(x, 1/2)$. In this way we construct a subsequence $\{x_{n_k}\}$ such that $x_{n_k} \in B(x, 1/k)$. This implies $x_{n_k} \rightarrow x$ contradicting our assumption on $\{x_n\}$. Now the family $\{B(x, \varepsilon_x)\}_{x \in A}$ is an open cover of A from which it is impossible to choose a finite number of balls which will cover A since any finite cover by these balls contains x_n for finitely many n and since A contains x_n for all positive integers. Consequently, A is compact. ■

6 Topological Spaces

Our next aim is to push the process of abstraction a little further and define spaces without distances in which continuous functions still make sense. The motivation behind the definition is the criterion of continuity in terms of open sets. This criterion tells us that a function between metric spaces is continuous provided that the preimage of an open set is open. We make the following definition.

Definition 6.1. *Let X be a non-empty set. A **topology** on a set X is a collection \mathcal{T} of subsets of X satisfying the following properties:*

- O1** \emptyset and $X \in \mathcal{T}$;
- O2** if $\{U_i\}_{i \in I} \subset \mathcal{T}$, then $\bigcup_{i \in I} U_i \in \mathcal{T}$;
- O3** if $U_1, U_2, \dots, U_n \in \mathcal{T}$, then $\bigcap_{i=1}^n U_i \in \mathcal{T}$;

*The pair (X, \mathcal{T}) is called a **topological space**. If X is a topological space with topology \mathcal{T} , we say that a subset U of X is an **open set** in X if $U \in \mathcal{T}$.*

Here are some examples of topological spaces.

Example 6.2. Let (X, d) be a metric space. Then the family of open subsets of X with respect to the metric d is a topology on X .

Example 6.3. Let X be any non-empty set. The collection of *all* subsets of X , $\mathcal{P}(X)$, is a topology on X . This topology is called the **discrete topology**. Every subset U of X is an open set. On the other extreme, consider X and the collection $\{\emptyset, X\}$. It is also a topology on X , and is called the **indiscrete topology** or the **trivial topology**.

Example 6.4. Let $X = \mathbb{R}$ and let \mathcal{T}_u be a collection of subsets of X consisting of \emptyset , \mathbb{R} , and the unbounded open intervals $(-\infty, a)$ for all $a \in \mathbb{R}$. Then \mathcal{T}_u is a topology on \mathbb{R} . Similarly, we can define a topology \mathcal{T}_l consisting of \emptyset , \mathbb{R} and all unbounded intervals (a, ∞) , $a \in \mathbb{R}$.

Example 6.5. Let (X, \mathcal{T}) be a topological space and $Y \subseteq X$. Then $\mathcal{T}_Y = \{U \cap Y \mid U \in \mathcal{T}\}$ is a topology on Y . It is called the **subspace topology** or **relative topology** induced by \mathcal{T} .

Definition 6.6. Suppose that \mathcal{T} and \mathcal{T}' are two topologies on X . If $\mathcal{T} \subset \mathcal{T}'$ we say that \mathcal{T}' is **finer** or **larger** than \mathcal{T} . In this case we also say \mathcal{T} is **coarser** or **smaller** than \mathcal{T}' . Topologies \mathcal{T} and \mathcal{T}' are **comparable** if $\mathcal{T}' \subset \mathcal{T}$ and $\mathcal{T} \subset \mathcal{T}'$.

Along with a concept of open sets there is the companion concept of closed set. If X is a topological space, then a set $F \subset X$ is **closed** if $F^c = X \setminus F$ is open. By de Morgan's laws, the family of closed sets is closed under arbitrary intersection of closed sets and finite unions. More precisely, the class of closed sets has the following properties:

- C1** X and \emptyset are closed;
- C2** If F_i is a closed set for every $i \in I$, then $\bigcap_{i \in I} F_i$ is closed;
- C3** If F_1, \dots, F_n are closed, then $\bigcup_{i=1}^n F_i$ is closed.

Given a subset A of a topological space X , its **closure** is the intersection of all closed subsets of X containing A . The closure of A is denoted by \overline{A} . The **interior** of A , denoted by A° , is the union of all open subsets of A . If $x \in X$, then a set $A \subset X$ is called a **neighbourhood** of x , if $x \in A^\circ$.


Basis

If X is a topological space with topology \mathcal{T} , then a **basis** for \mathcal{T} is a collection $\mathcal{B} \subset \mathcal{T}$ such that every member of \mathcal{T} , i.e., every open set, is a union of elements of \mathcal{B} .

Example 6.7. The collection of all open balls forms a basis for the topology of metric space.

Theorem 6.8. *Let X be a set. Then a collection \mathcal{B} of subsets of X is a basis for a topology of X if and only if \mathcal{B} has the following two properties:*

- (1) *For every $x \in X$, there exists $B \in \mathcal{B}$ such that $x \in B$.*
- (2) *If $B_1, B_2 \in \mathcal{B}$ and $x \in B_1 \cap B_2$, then there exists $B_3 \in \mathcal{B}$ such that $x \in B_3 \subset B_1 \cap B_2$.*



Proof. Any basis satisfies (1) since the whole space X is open, and (2) since the intersection of two open sets $B_1 \cap B_2$ is open. Conversely, assume that \mathcal{B} is a collection of subsets of X with properties (1) and (2). Define \mathcal{T} to be


the collection of all subsets of X that are unions of sets in \mathcal{B} . We shall show that \mathcal{T} is topology. The condition (1) guarantees that $X \in \mathcal{T}$. Clearly, an arbitrary union of sets in \mathcal{T} belongs to \mathcal{T} in view of definition of \mathcal{T} . Assume that $U, V \in \mathcal{T}$. We have to show that $U \cap V$ is the union of sets in \mathcal{B} . Take any $x \in U \cap V$. Since U and V are unions of sets in \mathcal{B} , there exist $B_1, B_2 \in \mathcal{B}$ such that $x \in B_1 \subset U$ and $x \in B_2 \subset V$. So $x \in B_1 \cap B_2$, and, in view of (2), there exists $B_x \in \mathcal{B}$ such that $x \in B_x \subset B_1 \cap B_2$. Hence $B_x \subset U \cap V$, and consequently,

$$U \cap V = \bigcup_{x \in U \cap V} B_x.$$

This shows that $U \cap V \in \mathcal{T}$. ■

Hausdorff and normal spaces

Definition 6.9. A topological space X is called a **Hausdorff space** if for every two points $x, y \in X$ such that $x \neq y$, there exist disjoint open sets U and V satisfying $x \in U$ and $y \in V$. A space X is **normal** if for each pair A, B of disjoint closed subsets of X , there exist disjoint open sets U and V such that $A \subset U$ and $B \subset V$.



Continuity

Continuous functions in metric spaces were characterized in terms of open and closed sets (see Theorem 3.4 and Theorem 3.5). This suggests the definition of continuity in topological spaces.

Definition 6.10. Let X and Y be topological spaces and let $f : X \rightarrow Y$. The map f is **continuous at a point** x_0 if for every neighbourhood U of $f(x_0)$ in Y there exists a neighbourhood V of x_0 in X such that $f(V) \subset U$. Global continuity of f is defined in terms of open sets: f is **continuous** if $f^{-1}(U)$ is open in X for every open set U in Y .

If $f : X \rightarrow Y$ is bijective and f and f^{-1} are both continuous, f is called a **homeomorphism** and X and Y are said to be **homeomorphic**. We call a property **topological** if it is invariant under homeomorphism.

Elementary properties of continuous functions

- (1) If $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are continuous maps between topological spaces, then the composition $g \circ f : X \rightarrow Z$ is continuous.
- (2) If $f : X \rightarrow \mathbb{R}$ and $g : X \rightarrow \mathbb{R}$ are continuous, then $h : X \rightarrow \mathbb{R}^2$ given by $h(x) = (f(x), g(x))$ is continuous.
- (3) If A is a subspace of X , then the inclusion map $i : A \rightarrow X$ is continuous (this follows from the definition of the topology on the subspace A). If $f : X \rightarrow Y$ is continuous, where Y is another topological space, then the restriction map $h : A \rightarrow Y$ defined by $h(x) = f(x)$ for $x \in A$, is continuous. This follows from (1) using the fact that $h = f \circ i$.

7 Compact Topological Spaces

Theorem 4.15 gives three equivalent characterizations of compactness for metric spaces: the Bolzano-Weierstrass property, completeness together with total boundedness and the Heine-Borel property. In the case of general topological spaces the most useful is the Heine-Borel property. A subset Y of a topological space (X, \mathcal{T}) is called **compact** if for every collection $\mathcal{U} = \{U_i\}_{i \in I}$ of open sets such that $A \subseteq \bigcup_{i \in I} U_i$, there is a finite $J \subseteq I$ for which $Y \subseteq \bigcup_{i \in J} U_i$. DeMorgan's laws lead to the following characterization of compactness in terms of closed sets.

Definition 7.1. A family $\{F_i\}_{i \in I}$ of closed subsets of X is said to have the **finite intersection property** if $\bigcap_{i \in J} F_i \neq \emptyset$ for all finite $J \subseteq I$.

Theorem 7.2. A topological space X has is compact if and only if for every family $\{F_i\}_{i \in I}$ of closed subsets of X having the finite intersection property, $\bigcap_{i \in I} F_i \neq \emptyset$.

Proof. Assume that X is compact. Let $\{F_i\}_{i \in I}$ be a collection of closed sets having the finite intersection property. Arguing by contradiction assume that $\bigcap_{i \in I} F_i = \emptyset$. Denoting by $U_i = X \setminus F_i$ we have $\bigcup_{i \in I} U_i = \bigcup_{i \in I} [X \setminus F_i] = X \setminus \bigcap_{i \in I} F_i = X$. So $\{U_i\}_{i \in I}$ is an open cover of X . Hence there are U_{i_1}, \dots, U_{i_k} such that $X = U_{i_1} \cup \dots \cup U_{i_k}$. But then $\emptyset = X \setminus X = X \setminus \bigcup_{l=1}^k U_{i_l} = \bigcap_{l=1}^k F_{i_l}$, contradicting the assumption that $\{F_i\}$ has the

finite intersection property. Conversely, suppose that for every collection $\{F_i\}_{i \in I}$ having the finite intersection property we have $\bigcap_{i \in I} F_i \neq \emptyset$. Take any open cover $\{U_i\}_{i \in I}$ of X , and define $F_i = X \setminus U_i$. Then F_i 's are closed and $\bigcap_{i \in I} F_i = \bigcap_{i \in I} [X \setminus U_i] = X \setminus \bigcup_{i \in I} U_i = \emptyset$. So $\{F_i\}$ does not have the finite intersection property (otherwise $\bigcap_{i \in I} F_i \neq \emptyset$). So there is a finite set $J \subseteq I$ such that $\bigcap_{i \in J} F_i = \emptyset$. But then $X = \bigcup_{i \in J} [X \setminus F_i] = \bigcup_{i \in J} U_i$ showing that X is compact. ■

Theorem 7.3. *A closed subspace of a compact topological space is compact*

Proof. Let K be a closed subset of a topological space X , and let $\{U_{i \in J}\}$ be an open cover of X . Then the collection $\{U_{i \in J} \cup \{K^c\}\}$ is a family of open subsets of X that covers X . Since X is compact, there is a finite subfamily of this family that covers X . The corresponding subfamily of $\{U_{i \in J}\}$ covers Y . ■

Theorem 7.4. *If X is a Hausdorff space, then every compact subset of X is closed.*

Proof. Let K be a compact subset of X . Since X is Hausdorff, for every $x \in K^c$ and every $y \in K$, there are disjoint open sets U_{xy} and V_{xy} such that $x \in U_{xy}$ and $y \in V_{xy}$. Then for every $x \in K^c$, $\{V_{xy}\}_{y \in K}$ is an open cover of K . Since K is compact, there exist $y_1, \dots, y_n \in K$ such that $K \subseteq \bigcup_{i=1}^n V_{xy_i}$. Set $U = \bigcap_{i=1}^n U_{xy_i}$. Then U is open, $U \cap K = \emptyset$, and $x \in U$. Thus $x \in U \subseteq K^c$ showing that K^c is open, and consequently, that K is closed. ■

Theorem 7.5. *A compact Hausdorff space is normal*

Proof. Let A and B be disjoint closed subsets of a compact Hausdorff space. In view of Theorem 7.3, the sets A and B are compact. Proceeding like in the proof of the previous theorem, we find for every $x \in B$ disjoint open sets V_x and U_x such that $x \in V_x$ and $A \subseteq U_x$. Then the open sets $\{V_x\}_{x \in B}$ cover B . Consequently, there exist $x_1, \dots, x_n \in B$ such that $B \subseteq V_{x_1} \cup \dots \cup V_{x_n} := V$. Then $U := U_{x_1} \cap \dots \cap U_{x_n}$ is open, $U \cap V = \emptyset$, and $A \subseteq U, B \subseteq V$. ■

Theorem 7.6. *Suppose that $f : X \rightarrow Y$ is a continuous map between topological spaces X and Y . If $K \subseteq X$ is a compact set, then $f(K)$ is a compact subset of Y . In particular, if X is compact, then $f(X)$ is compact.*

Proof. Let \mathcal{U} be an open cover of $f(K)$. That is, \mathcal{U} consists of open subsets of Y such that their union contains $f(K)$. The continuity of f implies that for any set $U \in \mathcal{U}$, $f^{-1}(U)$ is an open subset of X . Moreover, the family

$\{f^{-1}(U) \mid U \in \mathcal{U}\}$ is an open cover of K . Indeed, if $x \in K$, then $f(x) \in f(K)$, and so $f(x) \in U$ for some $U \in \mathcal{U}$. This implies that $x \in f^{-1}(U)$. Since K is compact, $K \subseteq \bigcup_{i=1}^n f^{-1}(U_i)$ for some n . It follows that $f(K) \subseteq \bigcup_{i=1}^n U_i$ which proves that $f(K)$ is a compact subset of Y . This completes the proof of the theorem. ■

Theorem 7.7. *Let f be a continuous bijective function from a compact topological space X to a Hausdorff topological space Y . Then the inverse function $f^{-1} : Y \rightarrow X$ is continuous.*

Proof. Denote by $g = f^{-1} : Y \rightarrow X$. We have to show that $g^{-1}(K)$ is closed in Y for any closed set K in X . Since f is a bijection, $g^{-1}(A) = f(A)$ for any subset of Y . So $g^{-1}(K) = f(K)$. Since K is closed and X compact, K is also compact. By the previous result, $f(K)$ is compact in Y and since Y is Hausdorff, $f(K)$ is closed. So $g^{-1}(K)$ is closed in Y , as required. ■

Example 7.8. Let S^1 be the unit circle in \mathbb{R}^2 of radius 1 and centre $(0, 0)$. We consider S^1 as a subspace of \mathbb{R}^2 . Let $f : [0, 2\pi) \rightarrow S^1$ be given by $f(x) = (\cos x, \sin x)$ for $x \in [0, 2\pi)$. Show that f is a continuous bijection but the inverse map $f^{-1} : S^1 \rightarrow [0, 2\pi)$ is not continuous. Why doesn't this contradict Theorem 6.7?

8 Connected Spaces

A pair of non-empty and open sets U, V of a topological space X is called a **separation** of X if $U \cap V = \emptyset$ and $X = U \cup V$. A topological space X is called **disconnected** if there is a separation of X , and otherwise is called **connected**. A subset Y of X is said to be connected if it is connected as a subspace of X , that is, Y is not the union of two non-empty sets $U, V \in \mathcal{T}_Y$ such that $U \cap V = \emptyset$.

Example 8.1. The set X containing at least two points and considered with the discrete topology is disconnected, however, X with the indiscrete topology is connected.

Example 8.2. The subspace $\mathbb{R} \setminus \{0\}$ of \mathbb{R} is disconnected since $\mathbb{R} \setminus \{0\} = A \cup B$, where $A = \{r \in \mathbb{R} \mid r < 0\}$ and $B = \{r \in \mathbb{R} \mid r > 0\}$. If $X = \mathbb{Q}$ is considered as subspace of \mathbb{R} , then X is disconnected since $X = A \cup B$ with $A = \mathbb{Q} \cap (-\infty, r)$ and $B = \mathbb{Q} \cap (r, \infty)$, where r is irrational.

A “2-valued” function is a function from X to $\{0, 1\}$, where $\{0, 1\}$ is considered with discrete topology.

Theorem 8.3. *A space X is connected if and only if every 2-valued continuous function on X is constant. Equivalently, X is disconnected if and only if there exists a 2-valued continuous function from X onto $\{0, 1\}$*

Proof. Suppose that X is connected and $f : X \rightarrow \{0, 1\}$ is continuous. Let $A = f^{-1}(\{0\})$ and $B = f^{-1}(\{1\})$. The sets A, B are open, disjoint and $X = A \cup B$. So one of A, B has to be empty. Conversely, assume that every continuous 2-valued function is constant. Assume that $X = A \cup B$, A and B are open, and $A \cap B = \emptyset$. Define

$$f(x) = \begin{cases} 0 & \text{if } x \in A, \\ 1 & \text{if } x \in B. \end{cases}$$

Clearly, the function f is continuous. So f is constant, say $f(x) = 0$ for all $x \in X$. But then $A = X$ and $B = \emptyset$. Hence X is connected as claimed. ■

Theorem 8.4. *Let $f : X \rightarrow Y$ be a continuous function between spaces X and Y . If X is connected, then the image $f(X)$ is connected.*

Proof. Let $g : f(X) \rightarrow \{0, 1\}$ be continuous. Then the composition $g \circ f : X \rightarrow \{0, 1\}$ is continuous, hence constant since X is connected. Hence g is constant on $f(X)$ and the result follows in view of Theorem 8.3. ■

Theorem 8.5. *If A is a connected subset of a space X , then \overline{A} is also connected.*

Proof. Let $f : \overline{A} \rightarrow \{0, 1\}$ be continuous. Then $f|_A$ is continuous, and so, f is constant on A . Say $f = 0$ on A . We claim that $f = 0$ on \overline{A} . Suppose $f(x) = 1$ for some $x \in \overline{A}$. The set $\{1\}$ is open in $\{0, 1\}$ and since f is continuous $f^{-1}(\{1\})$ is an open subset of \overline{A} . Thus say $f^{-1}(\{1\}) = U \cap \overline{A}$ for some open set U in X . This means that $f = 1$ on $U \cap \overline{A}$. Since $x \in \overline{A}$, $U \cap A \neq \emptyset$, say $y \in U \cap A$. Then $f(y) = 1$ since $y \in U \cap A \subseteq U \cap \overline{A}$, but on the other hand $f(y) = 0$ since $f = 0$ on A . Therefore, \overline{A} is connected as claimed. ■

Example 8.6. The union of connected subspaces does not have to be connected. Consider \mathbb{R} with the usual topology. Then the sets $(-\infty, 0)$ and $(0, \infty)$ are connected subspaces of \mathbb{R} , but the union $(-\infty, 0) \cup (0, \infty) = \mathbb{R} \setminus \{0\}$ is disconnected.

Theorem 8.7. *If $\{A_i\}_{i \in I}$ is a family of connected subsets of X such that $\bigcap_{i \in I} A_i \neq \emptyset$, then $A = \bigcup_{i \in I} A_i$ is connected.*

Proof. Let $f : A \rightarrow \{0, 1\}$ be continuous. Then $f|_{A_i}$ is continuous for every i , so it is constant. Since $\bigcap_{i \in I} A_i \neq \emptyset$, we must have the same constant on every A_i . Hence f is constant and A is connected. ■

As an application of this theorem we have the following

Theorem 8.8. *If for any two points in a space X there exists a connected subspace of X containing these two points, then X is connected.*

Proof. Fix a point $a \in X$. For $b \in X$ denote by $C(b)$ a connected subspace of X containing a and b . Then $X = \bigcup_{b \in X} C(b)$. Since $a \in \bigcap_{b \in X} C(b)$, the result follows from the previous theorem. ■

Let $x \in X$ and let C_x be the union of all the connected subsets of X containing x . Each C_x is called a **component** (or **connected component**) of X .

Proposition 8.9. *Let C_x be the connected component of X containing x . Then*

- (a) *for each $x \in X$, C_x is connected and closed; and*
- (b) *for any two $x, y \in X$, either $C_x = C_y$ or $C_x \cap C_y = \emptyset$.*

Proof. The set C_x is connected in view of Theorem 8.7, and by Theorem 8.5, $\overline{C_x}$ is connected. Hence by the definition of C_x , $\overline{C_x} \subset C_x$, so $C_x = \overline{C_x}$ and C_x is closed. If $C_x \cap C_y \neq \emptyset$, then $C_x \cup C_y$ is connected by Theorem 8.7. So again by the definition of C_x , $C_x \cup C_y \subset C_x$. Hence $C_y \subset C_x$. Similarly, $C_x \subset C_y$, so $C_x = C_y$ as required. ■

Example 8.10. If X is equipped with the discrete topology, then every subset of X is open and closed. Hence the connected components of X are sets consisting of one point.

Next we shall determine the connected subsets of \mathbb{R} . By an **interval** $I \subset \mathbb{R}$ we mean a subset of \mathbb{R} having the following property: if $x, y \in I$ and $x \leq z \leq y$, then $z \in I$.

Theorem 8.11. *A subset of \mathbb{R} is connected if and only if it is an interval.*

Proof. Suppose that $J \subset \mathbb{R}$ is not an interval. Then there are $x, y \in J$ and $z \notin J$ with $x < z < y$. Then define $A = (-\infty, z) \cap J$ and $B = (z, \infty) \cap J$. Clearly, A, B are disjoint, non-empty, relatively open, and $A \cup B = J$. So J is not connected. Conversely, suppose that J is an interval. We will show that J is connected. Let $f : J \rightarrow \{0, 1\}$ be continuous, and suppose that f is not constant. Then there are x_1 and $y_1 \in J$ such that $f(x_1) = 0$ and $f(y_1) = 1$. For simplicity assume that $x_1 < y_1$. Let a be the midpoint of $[x_1, y_1]$. If $f(a) = 0$, then set $x_2 = x_1$ and $y_2 = a$, and otherwise, $x_2 = a$ and $y_2 = y_1$. So $x_1 \leq x_2 \leq y_2 < y_1$, $|x_2 - y_2| \leq 2^{-1}|x_1 - y_1|$, and $f(x_i) \neq f(y_i)$. Iterating this procedure we find a sequences $\{x_n\}$ and $\{y_n\}$ with the following properties: $x_1 \leq x_2 \leq \dots \leq x_n < y_n \leq \dots \leq y_1$, $|x_n - y_n| \leq 2^{-1}|x_{n-1} - y_{n-1}| \leq 2^{n-1}|x_1 - y_1|$, and $f(x_n) = 0$, $f(y_n) = 1$. Since \mathbb{R} is complete, $\{x_n\}$ converges to some z , and since $|x_n - y_n| \rightarrow 0$, $y_n \rightarrow z$. Clearly, $z \in J$. Hence $0 = \lim_n f(x_n) = f(z) = \lim_n f(y_n) = 1$, a contradiction. So f is constant, and this implies that J is connected. ■

We can apply the last theorem to analyze the structure of open subsets of \mathbb{R} . We claim that any open set $U \subset \mathbb{R}$ is a countable union of pairwise disjoint open intervals. Indeed, let $x \in U$ and let I_x be the connected component of U containing x . Thus, I_x is an interval. If $y \in I_x$, then there is $\delta > 0$ such that $(y - \delta, y + \delta) \subset U$ since U is open. Hence $I_x \cup (y - \delta, y + \delta)$ is connected and since I_x is a connected component, $(y - \delta, y + \delta) \subset I_x$. So I_x is an open interval, and U is a union of open intervals (its components). Since each must contain a different rational number, U is at most countable union of disjoint open intervals.

Here is an important application of Theorem 8.11.

Theorem 8.12 (Intermediate Value Theorem). *Let f be a continuous function defined on a connected space X . Then for any $x, y \in X$ and any $r \in \mathbb{R}$ such that $f(x) \leq r \leq f(y)$ there exists $c \in X$ such that $f(c) = r$.*

Proof. The set $f(X)$ is a connected subset of \mathbb{R} . Hence $f(X)$ is an interval, and since $f(x), f(y) \in f(X)$, it has to contain r . ■

Definition 8.13. *A space X is called **path connected** if for any two points p and $q \in X$, there exists a continuous function $f : [0, 1] \rightarrow X$ such that $f(0) = p$ and $f(1) = q$. The function f is called a **path** from $f(0)$ to $f(1)$.*

If X is path connected, then X is connected but the converse is false in general as the following example shows.

Example 8.14. Denote by $X = \{(t, \sin(\pi/t)) \mid t \in [0, 2]\} \subset \mathbb{R}^2$. Let $\varphi : \mathbb{R}^2 \rightarrow \mathbb{R}$ be the projection onto the first coordinate, that is, $\varphi(x, y) = x$. Then $\varphi : X \rightarrow (0, 2]$ is a homeomorphism and since $(0, 2]$ is connected so is X . Therefore, $\overline{X} = (\{0\} \times [-1, 1]) \cup X = J \cup X$ is connected, where we abbreviated $J = \{0\} \times [-1, 1]$. We shall show that \overline{X} is not path connected. Arguing by contradiction assume that $f : [0, 1] \rightarrow \overline{X}$ is a continuous path in \overline{X} such that $f(0) \in J$ and $f(1) \in X$. Consider $f^{-1}(J)$. It is closed in $[0, 1]$ and contains 0. Let $a = \sup\{t \in [0, 1], t \in f^{-1}(J)\}$. Since $f(1) \in X$, $a < 1$. Since f is continuous, there exists $\delta > 0$ such that $f(a + \delta) \in X$. Write $f(t) = (x(t), y(t))$. Then $x(a) = 0$ and $x(t) > 0$, $y(t) = \sin(\pi/x(t))$ for $t \in (0, a + \delta]$. For every large n find r_n such that $0 < r_n < x(a + 1/n)$ and $\sin(\pi/r_n) = (-1)^n$. Since the function x is continuous by the Intermediate Value Theorem there is $t_n \in (a, a + \delta]$ such that $x(t_n) = r_n$ and $y(t_n) = (-1)^n$. So $t_n \rightarrow a$ but $y(t_n)$ does not converge contradicting the fact that f is continuous. Hence \overline{X} is not path connected.

9 Product Spaces

We define a topology on a finite product of topological spaces. Consider a finite collection X_1, \dots, X_n of topological spaces. The **product topology** on the product $X = X_1 \times \dots \times X_n$ is the topology for which a basis of open sets is given by “rectangles”

$$\{U_1 \times \dots \times U_n \mid U_j \text{ is open in } X_j \text{ for } 1 \leq j \leq n\}. \quad (11)$$

Observe that the intersection of two such sets is again a set of this form. Indeed,

$$(U_1 \times \dots \times U_n) \cap (V_1 \times \dots \times V_n) = (U_1 \cap V_1) \times \dots \times (U_n \cap V_n)$$


Consequently, the family (11) forms a basis. Let $\pi_j : X \rightarrow X_j$ be the projection of X onto the j th factor, defined by

$$\pi_j(x_1, \dots, x_n) = x_j, \quad (x_1, \dots, x_n) \in X.$$

For an open set $U_j \subset X_j$, we have

$$\pi_j^{-1}(U_j) = X_1 \times \dots \times X_{j-1} \times U_j \times X_{j+1} \times \dots \times X_n$$

which is a basic open set. Hence each projection π_j is continuous.



Theorem 9.1. *Let X be the product of the topological spaces X_1, \dots, X_n , and let π_j be the projection of X onto X_j . The product topology for X is the smallest topology for which each of the projections π_j is continuous.*

Proof. Let \mathcal{T} be another topology on X such that the projections π_j are \mathcal{T} -continuous. Take open sets $U_j \subset X_j$, $1 \leq j \leq n$. Then each $\pi_j^{-1}(U_j)$ belongs to \mathcal{T} since π_j is \mathcal{T} -continuous. Since

$$\pi_1^{-1}(U_1) \cap \dots \cap \pi_n^{-1}(U_n) = U_1 \times \dots \times U_n$$

the basic set $U_1 \times \dots \times U_n$ belongs to \mathcal{T} and \mathcal{T} includes the product topology. ■

Call a function f from one topological space to another **open** if it maps open sets onto open sets.

Theorem 9.2. *Let X be the product of the topological spaces X_1, \dots, X_n . Then each projection π_j of X onto X_j is open.*

Proof. Let $U = U_1 \times \dots \times U_n$ be a basic open set in X . Then $\pi_j(U) = U_j$, and since the maps preserve unions, the image of any open set is open. ■

Theorem 9.3. *Let Y be a topological space and let f be a continuous map from Y to the product $X = X_1 \times \dots \times X_n$. Then f is continuous if and only if $\pi_j \circ f$ is continuous for all $1 \leq j \leq n$.*

Proof. If f is continuous, the $\pi_j \circ f$ is continuous as a composition of continuous maps. Conversely, suppose that $\pi_j \circ f$ is continuous for all $1 \leq j \leq n$. Take a basic open set $U = U_1 \times \dots \times U_n$ in X . Then

$$f^{-1}(U) = (\pi_1 \circ f)^{-1}(U_1) \cap \dots \cap (\pi_n \circ f)^{-1}(U_n)$$

is a finite intersection of open sets and hence is open. Since the inverses of functions preserve unions, the inverse image of any open set is open, and consequently, f is continuous. ■

We next study which properties of topological spaces are valid for the product $X = X_1 \times \cdots \times X_m$ whenever they hold for X_1, X, \dots, X_m .

Theorem 9.4. *Let X be the product of Hausdorff spaces X_1, \dots, X_n . Then X is Hausdorff.*

Proof. Take two different points $x = (x_1, \dots, x_n)$, $y = (y_1, \dots, y_n)$, and choose an index i so that $x_i \neq y_i$. Since X_i is Hausdorff, there exist open sets U_i and V_i in X_i such that $U_i \cap V_i = \emptyset$. Then $\pi_i^{-1}(U_i)$ and $\pi_i^{-1}(V_i)$ are open and disjoint sets containing x and y , respectively. Consequently, X is Hausdorff as required. ■

Theorem 9.5. *Let X be the product of path-connected spaces X_1, \dots, X_n . Then X is path-connected.*

Proof. Take two points $x = (x_1, \dots, x_n)$, $y = (y_1, \dots, y_n)$ in X . Since each X_j is path-connected, for each $1 \leq j \leq n$ there exists a path $\gamma_j : [0, 1] \rightarrow X_j$ from x_j to y_j . Define $\gamma : [0, 1] \rightarrow X$ by setting

$$\gamma(t) = (\gamma_1(t), \dots, \gamma_n(t)), \quad t \in [0, 1].$$

Then γ is a path connecting x with y . So X is path-connected. ■

To study connectedness of the product of connected spaces we will need the following fact. Fix points $x_2 \in X_2, \dots, x_n \in X_n$ and define a map $h : X_1 \rightarrow X$ by setting $h(x_1) = (x_1, \dots, x_n)$. Then h is a homeomorphism of X_1 onto the “slice” $X_1 \times \{x_2\} \times \cdots \times \{x_n\}$ of X . Indeed, if $U = U_1 \times \cdots \times U_n$ is a basic open set in X , then $h^{-1}(U) = U_1$ is open so that h is continuous. Since the inverse of h is equal to $\pi_1|_{X_1 \times \{x_2\} \times \cdots \times \{x_n\}}$, h^{-1} is continuous and h is a homeomorphism. Similarly, for each j fixed and fixed points $x_i \in X_i$, $i \neq j$, the map $X_j \rightarrow \{x_1\} \times \cdots \times \{x_{j-1}\} \times X_j \times \{x_{j+1}\} \times \cdots \times \{x_n\}$ is a homeomorphism.

Theorem 9.6. *Let X be the product of connected spaces X_1, \dots, X_n . Then X is connected.*

Proof. We prove the theorem for the product of two connected spaces X_1 and X_2 . We apply Theorem 8.8. Take two points $a = (a_1, a_2), b = (b_1, b_2) \in X$ and consider sets $C_1 = \{(x, b_2) \in X \mid x \in X_1\}$ and $C_2 = \{(a_1, y) \in X \mid y \in X_2\}$. By the above remark, the sets C_1, C_2 are connected. Then, in view of Theorem 8.7, $C = C_1 \cup C_2$ is connected since $C_1 \cap C_2 = \{(a_1, b_2)\}$. Applying Theorem 8.8, the space X is connected since $a, b \in C$. ■

To study compactness of the product of compact spaces we need the following lemma.

Lemma 9.7. *Let Y be a topological space and let \mathcal{B} be a basis for the topology of Y . If every open cover of Y by sets in \mathcal{B} has a finite subcover, then Y is compact.*

Proof. Let $\{U_i\}_{i \in I}$ be an open cover of Y . For each $y \in Y$, choose $V_y \in \mathcal{B}$ and an index j so that $y \in V_y \subset U_j$. The family $\{V_y\}_{y \in Y}$ forms an open cover of Y by sets belonging to \mathcal{B} . In view of the assumption, there exists a finite number of the V_y 's that cover Y . Since each of these V_y 's is contained in at least one of the U_j 's, we obtain a finite number of U_j 's that cover Y . Hence Y is compact. ■

Theorem 9.8 (Tichonoff's Theorem for the finite product). *Let X be the product of compact spaces X_1, \dots, X_n . Then X is compact.*

Proof. We consider only the product of two compact spaces X_1 and X_2 . Let \mathcal{R} be a cover of $X_1 \times X_2$ by basic open sets of the form $U \times V$, U open in X_1 and V open in X_2 . In view of Lemma 9.7, it is enough to show that \mathcal{R} has a finite subcover. Fix $z \in X_2$. The slice $X_1 \times \{z\}$ is compact. Hence there are finitely many sets $U_1 \times V_1, \dots, U_n \times V_n$ in \mathcal{R} covering the slice $X_1 \times \{z\}$. We may assume that $z \in V_j$ for all $1 \leq j \leq n$. The set $V(z) = V_1 \cap \dots \cap V_n$ is an open set containing z , and the set $\pi_2^{-1}(V(z))$ is covered by sets $U_j \times V_j$, $1 \leq j \leq n$. The collection $\{V(z)\}_{z \in X_2}$ is an open cover of X_2 , and since X_2 is compact, $X_2 = V(z_1) \cup \dots \cup V(z_l)$ for some finite number of points $z_j \in X_2$. Then $X = \pi_2^{-1}(V(z_1)) \cup \dots \cup \pi_2^{-1}(V(z_l))$. Each $\pi_2^{-1}(V(z_j))$ is covered by finitely many sets in \mathcal{R} . Consequently, X can be covered by finitely many sets in \mathcal{R} , and, in view of Lemma 9.7, X is compact. ■

Compactness in function spaces: Ascoli-Arzelà theorem

Next we study compact subsets of the space of continuous functions. Let X be a compact topological space and (M, σ) a complete metric space. By $C(X, M)$ we denote the set of all continuous functions from X to M . We consider $C(X, M)$ with the metric

$$d(f, g) = \sup\{\sigma(f(x), g(x)) \mid x \in X\}$$

Definition 9.9. Let X be a topological space and (M, σ) a compact metric space, and let \mathcal{F} be a family of functions from X to M . The family \mathcal{F} is called **equicontinuous at** $x \in X$ if for every $\varepsilon > 0$ there exists a neighbourhood U_ε of x such that

$$\sigma(f(y), f(x)) < \varepsilon \text{ for all } y \in U_\varepsilon \text{ and all } f \in \mathcal{F}.$$

The family \mathcal{F} is called **equicontinuous** if it is equicontinuous at each $x \in X$.

Example 9.10. Consider two metric spaces (X, ρ) and (M, σ) . Given $M > 0$ let \mathcal{F} be a set of all functions $f : X \rightarrow M$ such that

$$\sigma(f(x), f(y)) \leq M\rho(x, y)$$

for all $x, y \in X$. Then \mathcal{F} is an equicontinuous family of functions. For if $\varepsilon > 0$, take $U_\varepsilon = B(x, \varepsilon/M)$. Then if $y \in U_\varepsilon$ and $f \in \mathcal{F}$, we have

$$\sigma(f(x), f(y)) \leq M\rho(x, y) < M \cdot \varepsilon/M = \varepsilon.$$

Theorem 9.11 (Ascoli-Arzelà Theorem). Let X be a compact space and let (M, σ) be a complete metric space. Let $\mathcal{F} \subset C(X, M)$. Then the closure $\overline{\mathcal{F}}$ is compact in $C(X, M)$ if and only if two of the following conditions hold:

- (1) \mathcal{F} is equicontinuous.
- (2) for each $x \in X$. the set $\mathcal{F}(x) = \{f(x) \mid f \in \mathcal{F}\}$ has a compact closure in M .

Proof. Since $C(X, M)$ is a complete metric space, $\overline{\mathcal{F}}$ is compact if and only if \mathcal{F} is totally bounded. Assume first that the conditions (1) and (2) are satisfied. In view of the above remark we have to show that \mathcal{F} is totally bounded. Given $\varepsilon > 0$, for each $x \in X$ there exists an open neighbourhood $V(x)$ such that if $y \in V(x)$, then $\sigma(f(x), f(y)) < \varepsilon$ for all $f \in \mathcal{F}$. Since $\{V(x)\}_{x \in X}$ is an open cover of X and X is compact by assumption, there exist a finite number of points x_1, \dots, x_n such that $V(x_1), \dots, V(x_n)$ cover X . The sets $\mathcal{F}(x_j)$ are totally bounded in M , hence so is the union $\mathcal{S} = \mathcal{F}(x_1) \cup \dots \cup \mathcal{F}(x_n)$. Let $\{a_1, \dots, a_m\}$ be an ε -net for \mathcal{S} . For every map $\varphi : \{1, \dots, n\} \rightarrow \{1, \dots, m\}$ denote by

$$B_\varphi = \{f \in \mathcal{F} \mid \sigma(f(x_j), a_{\varphi(j)}) < \varepsilon \text{ for all } j = 1, \dots, n\}.$$

Observe that there is only a finite number of sets B_φ and every $f \in \mathcal{F}$ belongs to one of such sets. Moreover, if $f, g \in \mathcal{F}$, then

$$\begin{aligned} \sigma(f(y), g(y)) &\leq \sigma(f(y), f(x_k)) + \sigma(f(x_k), a_{\varphi(k)}) \\ &\quad + \sigma(a_{\varphi(k)}, g(x_k)) + \sigma(g(x_k), g(y)) \\ &< \varepsilon + \varepsilon + \varepsilon + \varepsilon = 4\varepsilon \end{aligned}$$

for all $y \in V(x_k)$. So if $f, g \in B_\varphi$, then $d(f, g) < 4\varepsilon$. Consequently, the diameter of B_φ is $< 4\varepsilon$, and since there are finitely many such B_φ and they cover \mathcal{F} , the set \mathcal{F} is totally bounded.

Conversely, assume that \mathcal{F} is totally bounded. Note that the mapping $\Psi : \mathcal{F} \rightarrow M$ given by $\Psi(f) = f(x)$ is distance decreasing, i.e.,

$$\sigma(\Psi(f), \Psi(g)) = \sigma(f(x), g(x)) \leq d(f, g).$$

It follows that for every $x \in X$, the set $\mathcal{F}(x) \subset M$ is totally bounded and (2) holds. To see that (1) holds, let $\varepsilon > 0$ and let f_1, \dots, f_n be an ε -net of \mathcal{F} . Given $x \in X$ we find open neighbourhood $V(x)$ of x such that $\sigma(f_j(x), f_j(y)) < \varepsilon$ for all $y \in V(x)$ and all $j = 1, \dots, n$. Then if $f \in \mathcal{F}$ choose an index j so that $d(f, f_j) < \varepsilon$. It follows that if $y \in V(x)$, then

$$\begin{aligned} \sigma(f(x), f(y)) &\leq \sigma(f(x), f_j(x)) + \sigma(f_j(x), f_j(y)) + \sigma(f_j(y), f(y)) \\ &< \varepsilon + \varepsilon + \varepsilon = 3\varepsilon. \end{aligned}$$

Therefore, the family \mathcal{F} is equicontinuous at x , and since x was an arbitrary point of X , \mathcal{F} is equicontinuous as required. \blacksquare

Corollary 9.12. *Let X be a compact topological space and Y a compact metric space. Let \mathcal{F} be an equicontinuous a of $C(X, Y)$. Then every sequence in \mathcal{F} has a uniformly convergent subsequence.*

Definition 9.13. A family \mathcal{F} of maps $f : X \rightarrow Y$, where Y is a metric space is called **pointwise bounded** if $\{f(x) \mid f \in \mathcal{F}\}$ is bounded in Y for every $x \in X$.

Lemma 9.14. Assume that X is a compact metric space and let \mathcal{F} be an equicontinuous and pointwise bounded family in $C(X)$. Then there is a constant M such that $f(X) \subset [-M, M]$ for all $f \in \mathcal{F}$.

Proof. For each $x \in X$, there is M_x such that $|f(x)| \leq M_x$ for all $f \in \mathcal{F}$. Since \mathcal{F} is equicontinuous, for each x there is an open set U_x such that $|f(x) - f(y)| \leq 1$ for all $f \in \mathcal{F}$ and $y \in U_x$. Then

$$|f(y)| \leq |f(y) - f(x)| + |f(x)| \leq 1 + M_x = K_x$$

for all $y \in U_x$. The sets U_x form an open covering of X and since X is compact, there exists a finite subcovering U_{x_1}, \dots, U_{x_n} . Set now $M = \max\{K_{x_1}, \dots, K_{x_n}\}$. Then $|f(x)| \leq M$ for all $x \in X$. ■

Corollary 9.15 (Arzelà - Ascoli Theorem, classical version). Let X be a compact topological space. Assume that \mathcal{F} is a pointwise bounded and equicontinuous subset of $C(X)$. Then every sequence in \mathcal{F} has a uniformly convergent subsequence.

Proof. In view of the above exercise the set \mathcal{F} is uniformly bounded, that is, $|f(x)| \leq M$ for all $f \in \mathcal{F}$ and $x \in X$. Set $Y = [-M, M]$. Then Y is compact in \mathbb{R} , and \mathcal{F} is a subset of $C(X, Y)$. So the corollary follows from Corollary 9.12. ■

10 Uryshon's and Thietze's Theorems

We show the existence of continuous functions on normal topological spaces. We start with the following characterisation of normal spaces.

Lemma 10.1. A topological space X is normal if and only if for every closed subset $A \subset X$ and every open subset $B \subset X$ containing A , there exists an open set U such that $A \subset U \subset \overline{U} \subset B$.

Proof. Assume first that X is normal and A and B are as above. Then the sets A and $X \setminus B$ are closed and disjoint. So, in view of normality of X , there exist open disjoint sets U and V such that $A \subset U$ and $X \setminus B \subset V$. Then $\overline{U} \subset X \setminus V \subset B$, so that U has the required properties.

Conversely, let A and B be closed disjoint subsets of X . Then $V = X \setminus B$

is open and $A \subset V$. By assumption there exists an open set U such that $A \subset U \subset \overline{U} \subset V$. Then U and $X \setminus \overline{U}$ are disjoint open sets satisfying $A \subset U$ and $B \subset X \setminus \overline{U}$. So X is normal and the proof is completed. ■

Theorem 10.2 (Urysohn's Lemma). *Let A and B be closed subspace of a normal space X . Then we can find a continuous function $f : X \rightarrow [0, 1]$ such that $f(a) = 0$ for all $a \in A$ and $f(b) = 1$ for all $b \in B$.*

Proof. For the proof recall that a dyadic rational number is a number which can be written in the form $p = \frac{m}{2^n}$, with n, m being integers. Set $V = X \setminus B$, an open set which contains A . By Lemma 10.1, there exists an open set $U_{1/2}$ such that

$$A \subset U_{1/2} \subset \overline{U}_{1/2} \subset V.$$

Applying Lemma 10.1 again to the open set $U_{1/2}$ containing A and to the open set V containing $\overline{U}_{1/2}$, we obtain open sets $U_{1/4}$ and $U_{3/4}$ such that

$$A \subset U_{1/4} \subset \overline{U}_{1/4} \subset U_{1/2} \subset \overline{U}_{1/2} \subset U_{3/4} \subset \overline{U}_{3/4} \subset V.$$

Continuing in this way, we associate to each such number $p \in D$ an open subset $U_p \subset X$ having the following properties

$$\overline{U}_p \subset U_q, \quad 0 < p < q < 1, \quad (12)$$

$$A \subset U_q, \quad 0 < p < 1, \quad (13)$$

$$U_p \subset V, \quad 0 < p < 1. \quad (14)$$

Next we shall construct the function f which is continuous and such that the sets ∂U_p are level sets of f on which f assumes the value p . Define $f(x) = 0$ if $x \in U_p$ for all $p > 0$ and $f(x) = \sup\{p \mid x \notin U_p\}$ otherwise. Clearly, $0 \leq f \leq 1$, $f(x) = 0$ for all $x \in A$ and $f(x) = 1$ for all $x \in B$. It remains to show that f is continuous. Take $x \in X$. We only consider the case that $0 < f(x) < 1$. (The remaining cases $f(x) = 0$ and $f(x) = 1$ are left as an exercise). Let $\varepsilon > 0$ and choose dyadic rationals p and q such that $0 < p, q < 1$ and

$$f(x) - \varepsilon < p < f(x) < q < f(x) + \varepsilon.$$

Then $x \notin U_r$ for dyadic rationals r between p and $f(x)$ so that, in view of (12), $x \notin \overline{U}_p$. On the other hand $x \in U_q$. So $W = U_q \setminus \overline{U}_p$ is an open neighbourhood of x . Then $p \leq f(y) \leq q$ for any $y \in W$ which shows that $|f(x) - f(y)| < \varepsilon$ for all $y \in W$. Hence f is continuous and the proof is completed. ■

Theorem 10.3 (Thietze's extension theorem). *Let A be a closed subset of a normal space X and let f be a bounded continuous real valued function on A . Then there exists a bounded continuous function $h : X \rightarrow \mathbb{R}$ such that $f = h$ on A .*

Proof. Set $a_0 = \sup\{|f(a)| \mid a \in A\}$. (Note that $a < \infty$ since f is bounded). Define sets

$$B_0 = \{a \in A \mid f(a) \leq -a_0/3\} \quad C_0 = \{a \in A \mid f(a) \geq a_0/3\}.$$

Since f is continuous on A and A is closed, the sets B and C are closed and disjoint subsets of X . Taking a linear combination of constant function and the function from Uryshon's lemma, we find a continuous function $g_0 : X \rightarrow \mathbb{R}$ such that $-a_0/3 \leq g_0 \leq a_0/3$, $g_0 = -a_0/3$ on B_0 and $g_0 = a_0/3$ on C_0 . In particular,

$$\begin{aligned} |g_0| &\leq a_0/3 \\ |f - g_0| &\leq 2a_0/3. \end{aligned}$$

Iterating this process we construct the sequence of functions $\{g_n\}$ satisfying

$$|g_n| \leq 2^n a_0 / 3^{n+1} \tag{15}$$

$$|f - g_0 - g_1 - \cdots - g_n| \leq 2^{n+1} a_0 / 3^{n+1} \quad \text{on } A. \tag{16}$$

Indeed, suppose that the functions g_0, \dots, g_{n-1} have been constructed. To construct g_{n+1} , set

$$a_{n-1} = \sup\{|f(a) - g_0(a) - g_1(a) - \cdots - g_{n-1}(a)| \mid a \in A\},$$

and repeat the above argument with a_{n-1} replacing a_0 and $f - g_0 - g_1 - \cdots - g_{n-1}$ replacing f . This gives the function g_n such that

$$\begin{aligned} |g_n| &\leq a_{n-1}/3 \\ |f - g_0 - g_1 - \cdots - g_n| &\leq 2a_{n-1}/3 \quad \text{on } A. \end{aligned}$$

Since $a_{n-1} \leq 2^n a_0 / 3^n$, the function g_n satisfies (15)-(16). Set

$$h_n = g_0 + \cdots + g_n, \quad n \geq 1.$$

If $n > m$, then

$$\begin{aligned} |h_n - h_m| &= |g_{m+1} + \cdots + g_n| \leq \left(\left(\frac{2}{3} \right)^{m+1} + \cdots + \left(\frac{2}{3} \right)^n \right) \cdot \frac{a_0}{3} \\ &\leq \left(\frac{2}{3} \right)^{m+1} \cdot a_0. \end{aligned}$$

Consequently, $\{h_n\}$ is Cauchy in $C(X, \mathbb{R})$. Hence there exists a continuous function $h : X \rightarrow \mathbb{R}$ such that $h_n \rightarrow h$. In addition,

$$|h| = |\lim h_n| = \lim |h_n| \leq \lim \sum_{k=0}^n |g_k| \leq \frac{a_0}{3} \sum_{k=1}^{\infty} \left(\frac{2}{3}\right)^k = a_0,$$

so that h is bounded. Finally, in view of (16), $|f - h| = \lim |f - h_n| \leq \lim 2^n a_0 / 3^{n+1} \rightarrow 0$ on A , so that $f = h$ on A . The proof is completed. ■

Both theorems are valid in metric spaces as the following theorem shows.

Theorem 10.4. *Every metric space is normal.*

Proof. Let A and B be disjoint closed subsets of X . Define

$$f(x) = \inf\{d(x, a) \mid a \in A\}$$

for $x \in X$. Observe that $f(x) = 0$ if and only if $x \in A$ since A is closed. The function f is continuous. Similarly, let $g(x) = \inf\{d(x, b) \mid b \in B\}$. Then g is continuous and $g(x) = 0$ if and only if $x \in B$. Since A and B are disjoint, $f(x) + g(x) > 0$ for all $x \in X$. Set

$$h(x) = \frac{f(x)}{f(x) + g(x)}, \quad x \in X$$

Then h is continuous, $h(x) = 0$ if and only if $x \in A$ and $h(x) = 1$ if and only if $x \in B$. Take now $U = \{x \mid h(x) > 3/4\}$ and $V = \{x \mid h(x) < 1/4\}$. Clearly, $U \cap V = \emptyset$, U, V are open, and $A \subset U$, $B \subset V$, so that X is normal. ■

11 Appendix

Sets

A **set** is considered to be a collection of objects. The objects of a set A are called **elements** (or **members**) of A . If x is an element of a set A we write $x \in A$, and if x is not an element of A we write $x \notin A$. Two sets A and B are called **equal**, $A = B$, if A and B have the same elements. A set A is a **subset** of a set B , written $A \subset B$, if every element of A is also an element of B . The **empty set** \emptyset has no elements; it has the property that it is a subset of any set, that is, $\emptyset \subset A$ for any set A . Given two sets A and B we define:

(a) the **union** $A \cup B$ of A and B as the set

$$A \cup B = \{x \mid x \in A \text{ or } x \in B\};$$

(b) the **intersection** $A \cap B$ of A and B as the set

$$A \cap B = \{x \mid x \in A \text{ and } x \in B\};$$

(c) the set **difference** $A \setminus B$ of A and B as the set

$$A \setminus B = \{x \mid x \in A \text{ and } x \notin B\}.$$

Sets A and B are called **disjoint** if $A \cap B = \emptyset$. The concept of union and intersection of two sets extends to unions and intersections of arbitrary families of sets. By a **family of sets** we mean a nonempty set \mathcal{F} whose elements are sets themselves. If \mathcal{F} is a family of sets, then

$$\bigcup_{A \in \mathcal{F}} A = \{x \mid x \in A \text{ for some } A \in \mathcal{F}\}$$
$$\bigcap_{A \in \mathcal{F}} A = \{x \mid x \in A \text{ for all } A \in \mathcal{F}\}.$$

When it is understood that all sets under considerations are subsets of a fixed set X , then the **complement** A^c of a set $A \subset X$ is defined by

$$A^c = X \setminus A = \{x \in X \mid x \notin A\}.$$

In this situation we have **deMorgan's laws**:

$$\left(\bigcup_{i \in I} A_i \right)^c = \bigcap_{i \in I} A_i^c, \quad \left(\bigcap_{i \in I} A_i \right)^c = \bigcup_{i \in I} A_i^c.$$

The set of all subsets of a given set X is called the **power set** and is denoted by $\mathcal{P}(X)$.

If X and Y are sets, their **cartesian product** $X \times Y$ is the set consisting of ordered pairs (x, y) with $x \in X$ and $y \in Y$.

Given two sets X and Y , a **relation** from X to Y is subset R of $X \times Y$. We say that R is a relation on X if $X \times X$, that is, $R \subset X \times X$. Quite often we write xRy instead of $(x, y) \in R$.

The most important example of a relation is a **function**. A relation f from X to Y is called a function if for each $x \in X$ there exists exactly one $y \in Y$

such that xfy . If xfy , we write $y = f(x)$; y is called the **value** of f at x . We also will write $f : X \rightarrow Y$ to mean that f is a function from X to Y . Here X is called the **domain** of f , and the set $\{f(x) \mid x \in X\}$ is called the **range** of f . If $f : X \rightarrow Y$ is a function, $A \subset X$ and $B \subset Y$, then the **image** of A and the **preimage** of B under f are sets defined by

$$f(A) = \{f(x) \mid x \in A\}, \quad f^{-1}(B) = \{x \mid f(x) \in B\}.$$

We say that f is **injective**, or **one-one**, if $f(x) = f(y)$ only when $x = y$, and we say that f is **surjective**, or **onto**, if $f(X) = Y$, that is, if the image of f is the whole of Y . A function which is both injective and surjective is called **bijective**. Sometimes we will use words a “map” or a “mapping” instead of a function. Unions and Intersections behave nicely under inverse image:

$$\begin{aligned} f^{-1}\left(\bigcup_{i \in I} A_i\right) &= \bigcup_{i \in I} f^{-1}A_i. \\ f^{-1}\left(\bigcap_{i \in I} A_i\right) &= \bigcap_{i \in I} f^{-1}A_i. \\ f^{-1}(A^c) &= (f^{-1}(A))^c. \end{aligned}$$

Given two functions $f : X \rightarrow Y$ and $g : Y \rightarrow Z$, we define the **composition** $g \circ f$ of f and g as the function $g \circ f : X \rightarrow Z$ defined by the equation $g \circ f(x) = g(f(x))$. If $f : X \rightarrow Y$ is one-one, then f has the **inverse** f^{-1} . The inverse f^{-1} is defined on the range $f(X)$ and takes values in X ; it is given by the formula $f^{-1}(y) = x$ if and only if $f(x) = y$.

Countable and Uncountable Sets

A set A is called **finite** if for some $n \in \mathbb{N}$, there is a bijection f from $\{1, \dots, n\}$ to A . The number n is uniquely determined and is called the **cardinality of** A . We denote this fact by $\sharp A = n$ or $\text{card}(A) = n$. If A is not finite, then it is called **infinite**. If A is infinite, then there is an injective function f from the set of natural numbers \mathbb{N} into A . If there exists a bijection between \mathbb{N} and A , then we say that X is **countably infinite** (or just countable). So A is countably infinite if and only if its elements can be listed in an infinite sequence $X = \{x_1, x_2, \dots\}$. If there is no bijection between \mathbb{N} and A , then A is called **uncountable**.

Example 11.1. The set \mathbb{Z} of all integers is countably infinite. To see this

consider the function $f : \mathbb{N} \rightarrow \mathbb{Z}$ defined by

$$f(n) = \begin{cases} n/2 & \text{if } n \text{ is even;} \\ -(n-1)/2 & \text{if } n \text{ is odd.} \end{cases}$$

Check that the function f is a bijection from \mathbb{N} to \mathbb{Z} so that \mathbb{Z} is countably infinite.

Example 11.2. Consider the interval $I = [0, 1]$. Then I is uncountable. Seeking a contradiction, suppose that I is countable. Hence all elements of I can be listed as an infinite sequence $\{x_1, x_2, \dots\}$:

$$\begin{aligned} x_1 &= 0.a_1^1 a_2^1 a_3^1 \cdots \\ x_2 &= 0.a_1^2 a_2^2 a_3^2 \cdots \\ x_3 &= 0.a_1^3 a_2^3 a_3^3 \cdots \\ &\vdots \quad \quad \vdots \end{aligned}$$

Define

$$b_n = \begin{cases} 1 & \text{if } a_n^n \neq 1 \\ 2 & \text{if } a_n^n = 1 \end{cases}$$

and $x = 0.b_1 b_2 b_3 \cdots$. Then $x \in [0, 1]$ but it is not a member of $\{x_n\}$, contradiction.

Proposition 11.3. *Let A be a non-empty set. Then the following are equivalent:*

- (a) A is countable.
- (b) There exists a surjection $f : \mathbb{N} \rightarrow A$.
- (c) There exists an injection $g : A \rightarrow \mathbb{N}$.

Proof. Assume that A is countable. If A is countably infinite, then there exists a bijection $f : \mathbb{N} \rightarrow A$. If A is finite, then there is bijection $h : \{1, \dots, n\} \rightarrow A$ for some n . Define $f : \mathbb{N} \rightarrow A$ by

$$f(i) = \begin{cases} h(i) & \text{if } 1 \leq i \leq n, \\ h(n) & \text{if } i > n. \end{cases}$$

Check that f is a surjection. So the implication $(a) \implies (b)$ is proved. To prove the implication $(b) \implies (c)$. Let $f : \mathbb{N} \rightarrow A$ be a surjection. Define $g : A \rightarrow \mathbb{N}$ by the equation $g(a) = \text{smallest number in } f^{-1}(a)$. Since f is a surjection, $f^{-1}(a)$ is non-empty for any $a \in A$, so that g is well-defined. Next check that if $a \neq a'$, then $f^{-1}(a)$ and $f^{-1}(a')$ are disjoint, so they have different smallest elements. The injectivity of g follows. Now the implication $(c) \implies (a)$. Assume that $g : A \rightarrow \mathbb{N}$ is injective. We want to show that A is countable. Note that g from A to $g(A)$ is a bijection. So it suffices to show that any subset B of \mathbb{N} is countable. This is obvious when B is finite. Hence assume that B is an infinite subset of \mathbb{N} . We define a bijection $h : \mathbb{N} \rightarrow B$. Let $h(1)$ be the smallest element of B . Since B is infinite, it is non-empty and so $h(1)$ is well-defined. Having already defined $h(n-1)$, let $h(n)$ be the smallest element of the set $\{k \in B \mid k > h(n-1)\}$. Again this set is non-empty, so $h(n)$ is well-defined. Now check that the function h is a bijection from \mathbb{N} to B . ■

Corollary 11.4. *The set $\mathbb{N} \times \mathbb{N}$ is countable.*

Proof. In view of the previous proposition, it is enough to construct an injective function $f : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$. For example, let $f(n, m) = 2^n 3^m$. Suppose that $2^n 3^m = 2^k 3^l$. If $n < k$, then $3^m = 2^{k-n} 3^l$. The left side of this equality is an odd number whereas the right is an even number. So $n = k$, and $3^m = 3^l$. But then also $m = l$. Hence f is injective as required. ■

Proposition 11.5. *If A and B are countable, then $A \times B$ is countable.*

Proof. Since A and B are countable, there exist surjective functions $f : \mathbb{N} \rightarrow A$ and $g : \mathbb{N} \rightarrow B$. Define $h : \mathbb{N} \times \mathbb{N} \rightarrow A \times B$ by $h(n, m) = (f(n), g(m))$. The function h is surjective and $\mathbb{N} \times \mathbb{N}$ is countable, so $A \times B$ is countable. ■

Corollary 11.6. *The set \mathbb{Q} of all rational numbers is countable.*

Proposition 11.7. *If I is a countable set and A_i is countable for every $i \in I$, then $\bigcup_{i \in I} A_i$ is countable.*

Proof. For each $i \in I$, there exists a surjection $f_i : \mathbb{N} \rightarrow A_i$. Moreover, since I is countable, there exists a surjection $g : \mathbb{N} \rightarrow I$. Now define $h : \mathbb{N} \times \mathbb{N} \rightarrow \bigcup_{i \in I} A_i$ by $h(n, m) = f_{g(n)}(m)$. Check that h is surjective so that $\bigcup_{i \in I} A_i$ is countable. ■

Real numbers, Sequences

The set of all real numbers, \mathbb{R} , has the following properties:

- (a) the arithmetic properties,
- (b) the ordering properties, and
- (c) the completeness property.

The arithmetic properties start with the fact that any two real numbers a, b can be added to produce a real number $a + b$, the sum of a and b . The rules for addition are $a + b = b + a$, $(a + b) + c = a + (b + c)$. There is a real number 0, called zero, such that $a + 0 = 0 + a = a$ for all real numbers a . Each real number a has a negative $-a$ such that $a + (-a) = 0$. Besides addition, we have multiplication; two real numbers a, b can be multiplied to produce the product of a and b , $a \cdot b$. The rules for multiplication are $ab = ba$ and $(ab)c = a(bc)$. There is a real number 1, called one, such that $a1 = 1a = a$, and for each $a \neq 0$, there is a reciprocal $1/a$ such that $a(1/a) = 1$.

The ordering properties start with the fact that there is a subset \mathbb{R}^+ of \mathbb{R} , the set of positive real numbers. The set \mathbb{R}^+ is characterized by the property: if $a, b \in \mathbb{R}^+$, then $a + b$ and $ab \in \mathbb{R}^+$. The fact that $a \in \mathbb{R}^+$ is denoted by $0 < a$. The set of negative real numbers $\mathbb{R}^- = -\mathbb{R}^+$ is the set of negatives of elements in \mathbb{R}^+ . For every $a \in \mathbb{R}$, we have $a \in \mathbb{R}^+$ or $a = 0$ or $a \in \mathbb{R}^-$. The notation $a < b$ (or $b > a$) means that $b - a \in \mathbb{R}^+$. We also write $a \leq b$ to mean $a < b$ or $a = b$. The order properties of real numbers are as follows:

- (a) $a < b$ and $b < c$, then $a < c$.
- (b) $a < b$ and $c > 0$, then $ac < bc$.
- (c) $a < b$ and $c \in \mathbb{R}$, then $a + c < b + c$.
- (d) $a < b$ and $a, b > 0$, then $1/b < 1/a$.

If $A \subset \mathbb{R}$, a number M is called an **upper bound** for A if $a \leq M$ for all $a \in A$. Similarly, m is a **lower bound** for A if $m \leq a$ for all $a \in A$. A subset A of \mathbb{R} is called **bounded above** if it has an upper bound, and is called **bounded below** if it has a lower bound. If A has an upper and lower bound, then it is called **bounded**. A given subset of \mathbb{R} may have several upper bounds. If A has an upper bound M such that $M \leq b$ for any upper bound b of A , then we call M a **least upper bound** of A or **supremum** of A , and denote it by $M = \sup A$. Similarly, a real number m is called **greatest lower bound** of A or **infimum** of A if m is a lower bound of A and $b \leq m$

for all lower bounds b of A . If m is the greatest lower bound of A , we write $m = \inf A$.

The **completeness property** of \mathbb{R} asserts that every non-empty subset $A \subset \mathbb{R}$ that is bounded above has a least upper bound, and that every non-empty subset $S \subset \mathbb{R}$ which is bounded below has a greatest lower bound. Useful characterisations of a least upper bound and a greatest lower bound are contained in the following propositions:

Proposition 11.8. *Let $A \subset \mathbb{R}$ be bounded above. Then $a = \sup A$ if and only if $x \leq a$ for any $x \in A$, and for any $\varepsilon > 0$ there exists $x \in A$ such that $a < x + \varepsilon$.*

Proof. Assume first that $a = \sup A$. Clearly, $x \leq a$ for any $x \in A$. Take $\varepsilon > 0$. If for all $x \in A$, $x + \varepsilon \leq a$, then $x \leq a - \varepsilon$ for all x . Hence $a - \varepsilon$ is an upper bound of A contradicting the definition of a as the least upper bound of A . Conversely, from $x \leq a$ for any $x \in A$ follows that a is an upper bound of A . Assume that there is an upper bound b such that $b < a$. Then we get a contradiction with the fact that for any $\varepsilon > 0$ there exists $x \in A$ such that $a < x + \varepsilon$. Let $\varepsilon := (a - b)/2$ and $x \in A$. Then $x + \varepsilon \leq b + \varepsilon = (a + b)/2 < a$. ■

There is also a similar characterisation of $\inf A$ provided that A is bounded from below.

Proposition 11.9. *Let $A \subset \mathbb{R}$ be bounded from below. Then $a = \inf A$ if and only if $a \leq x$ for any $x \in A$, and for any $\varepsilon > 0$ there exists $x \in A$ such that $x - \varepsilon < a$.*

The proof of the proposition follows from the previous one by observing two facts: if A is bounded from below then the set $-A = \{x \mid -x \in A\}$ is bounded from above and that $\sup(-A) = -\inf A$.

It is useful to introduce the **extended real number system**, $\overline{\mathbb{R}} = \mathbb{R} \cup \{-\infty, \infty\}$ by adjoining symbols ∞ and $-\infty$ subject to the ordering rule $-\infty < a < \infty$ for all $a \in \mathbb{R}$. If A is not bounded above, then we write $\sup A = \infty$, and if A is not bounded below we write $\inf A = -\infty$. For example, we have $\inf \mathbb{R} = -\infty$ and $\sup \mathbb{R} = \infty$. We also have $\sup \emptyset = -\infty$ and $\inf \emptyset = \infty$, and for all non-empty sets A , $\inf A \leq \sup A$. With this terminology, the completeness property asserts that every subset of \mathbb{R} has a least upper bound and a greatest lower bound. The arithmetic operations

on \mathbb{R} can be partially extended to $\overline{\mathbb{R}}$. In particular we have:

$$\begin{aligned}\pm\infty + r &= r + \pm\infty = \pm\infty \quad \text{for } r \in \mathbb{R} \\ (\infty) + (\infty) &= \infty, \quad \text{and} \quad (-\infty) + (-\infty) = -\infty.\end{aligned}$$

Subtraction is defined in a similar way with the exception that

$$(\infty) + (-\infty) \quad \text{and} \quad (-\infty) + (\infty)$$

are not defined. We also define multiplication by

$$r(\pm\infty) = (\pm\infty)r = \begin{cases} \pm\infty, & \text{if } r > 0, \\ \mp\infty, & \text{if } r < 0, \end{cases}$$

and

$$(\pm\infty)(\pm\infty) = \infty, \quad (\pm\infty)(\mp\infty) = -\infty.$$

The multiplication $0 \cdot (\pm\infty)$ is not defined.

If a is an upper bound of A and $a \in A$, then a is a **maximum** of A , and we write $a = \max A$. Similarly, if $a \in A$ is a lower bound of A , then a is a **minimum** of A and this fact is denoted by $a = \min A$. If A and $B \subset \mathbb{R}$, then $A + B = \{a + b \mid a \in A, b \in B\}$, $A + a = \{x + a \mid x \in A\}$, and $aA = \{ax \mid x \in A\}$. Here are some properties of supremum and infimum:

- (a) **monotonicity property:** $A \subset B$, then $\sup A \leq \sup B$ and $\inf B \leq \inf A$.
- (b) **reflection property:** $\sup(-A) = -\inf A$ and $\inf(-A) = -\sup A$.
- (c) **translation property:** $\sup(A + a) = \sup A + a$ and $\inf(A + a) = \inf A + a$.
- (d) **dilation property:** $\sup(aA) = a \sup A$ and $\inf(aA) = a \inf A$ provided that $a > 0$.
- (e) **addition property:** $\sup(A + B) = \sup A + \sup B$ and $\inf(A + B) = \inf A + \inf B$.

A **sequence** of real numbers is a function $f : \mathbb{N} \rightarrow \mathbb{R}$. We often write the sequence as $\{f(n)\}$ or $\{f_n\}$. A sequence $\{a_n\}$ of real numbers is said to **converge** to a real number a if for every $\varepsilon > 0$ there is an integer n_0 such that if $n \geq n_0$, then $|a_n - a| < \varepsilon$. In this situation we call a the **limit** of $\{a_n\}$; a convergent sequence has a unique limit. We also write $a_n \rightarrow a$ or $\lim_{n \rightarrow \infty} a_n = a$. A sequence $\{a_n\}$ which does not converge to any limit in

\mathbb{R} is said to **diverge**. We say that $a_n \rightarrow \infty$, if for every $M > 0$, there is n_0 such that $a_n > M$ for all $n \geq n_0$. Similarly, $a_n \rightarrow -\infty$, if for every $M < 0$ there exists n_0 such that $a_n < M$ for all $n \geq n_0$. A sequence $\{a_n\}$ is **bounded** if $|a_n| < M$ for some number M and all $n \in \mathbb{N}$. A convergent sequence is always bounded. Here are some elementary properties of limits of sequences:

Proposition 11.10. *Let $\{a_n\}$ and $\{b_n\}$ be convergent sequences to a and b , respectively. Let $c \in \mathbb{R}$. Then we have:*

- (a) $\{ca_n\}$ converges to ca .
- (b) the sequence $\{a_n + b_n\}$ converges to $a + b$
- (c) the sequence $\{a_n \cdot b_n\}$ converges to $a \cdot b$
- (d) if $b_n \neq 0$ for all n and $b \neq 0$, then the sequence $\{a_n/b_n\}$ converges to a/b

A sequence $\{a_n\}$ is called **monotone increasing** if $a_n \leq a_{n+1}$ for all $n \in \mathbb{N}$. It is **monotone decreasing** if $a_n \geq a_{n+1}$ for all $n \in \mathbb{N}$.

Proposition 11.11. *If $\{a_n\}$ is a monotone increasing sequence that is bounded above, $a_n \leq M$ for all n , then $\{a_n\}$ is convergent. If $\{a_n\}$ is monotone increasing and it is unbounded from above, then $a_n \rightarrow \infty$. If $\{a_n\}$ is monotone decreasing and it is bounded below, $M \leq a_n$ for all n , then $\{a_n\}$ is convergent, and if $\{a_n\}$ is unbounded from below, then $a_n \rightarrow -\infty$.*

Proof. If $\{a_n\}$ is unbounded from above, then for every M there is k such that $a_k > M$. Since the sequence is increasing, $a_n \geq a_k \geq M$ for all $n \geq k$. Thus $a_n \rightarrow \infty$. Next assume that $\{a_n\}$ is bounded above. Then $a := \sup\{a_n \mid n \in \mathbb{N}\} < \infty$. Let $\varepsilon > 0$. By the definition of supremum, $a_n \leq a$ for all n and there is an integer n_0 such that $a < a_{n_0} + \varepsilon$. Since $\{a_n\}$ is monotone increasing, $a_n \leq a < a_n + \varepsilon$ for all $n \geq n_0$, that is, $|a_n - a| < \varepsilon$ for all $n \geq n_0$. Thus the sequence converges to a . The proof for monotonically decreasing sequences is similar and is left as an exercise. ■

Let $\{a_n\}$ be a sequence. If $0 < n_1 < n_2 < \dots$ are positive integers, then $\{a_{n_k}\}$ is called a **subsequence** of $\{a_n\}$.

Proposition 11.12. *If $\{a_n\}$ is a convergent sequence with the limit a , then every subsequence of $\{a_n\}$ converges to a . Conversely, if a sequence $\{a_n\}$ has the property that each of its subsequences is convergent, then $\{a_n\}$ itself converges.*

Proof. Let $\{a_{n_k}\}$ be a subsequence of $\{a_n\}$. For a given $\varepsilon > 0$ choose n_0 such that $|a_n - a| < \varepsilon$ for all $n > n_0$. Note that if $k > n_0$, then $n_k > n_0$ and so $|a_{n_k} - a| < \varepsilon$ for all $k > n_0$. Therefore, $\{a_{n_k}\}$ converges to a . The converse follows from the fact that the sequence $\{a_n\}$ is a subsequence of itself. ■

Let $\{a_n\}$ be a bounded sequence. For each $n \in \mathbb{N}$, let $b_n = \sup_{m \geq n} a_m = \sup\{a_n, a_{n+1}, \dots\}$. Then $\{b_n\}$ is monotone decreasing, and it is bounded since $\{a_n\}$ is bounded. In view of Proposition 11.11, $\{b_n\}$ converges. The limit is called **upper limit** of $\{a_n\}$. Similarly, let $c_n = \inf_{m \geq n} a_m = \inf\{a_n, a_{n+1}, \dots\}$. Then $\{c_n\}$ is monotone increasing, and it is bounded since $\{a_n\}$ is bounded. The limit of $\{c_n\}$ is called **lower limit** of $\{a_n\}$. If $\{a_n\}$ is not bounded above, then its upper limit is equal to ∞ , and if $\{a_n\}$ is not bounded below, then its lower limit is equal to $-\infty$. Summarizing

$$\begin{aligned}\limsup a_n &= \overline{\lim} a_n = \inf_{n \geq m} \sup_{k \geq n} a_k = \lim_{n \rightarrow \infty} \sup_{k \geq n} a_k \\ \liminf a_n &= \underline{\lim} a_n = \sup_{n \geq m} \inf_{k \geq n} a_k = \lim_{n \rightarrow \infty} \inf_{k \geq n} a_k.\end{aligned}$$

A useful characterisation of the upper limit is the following proposition.

Proposition 11.13. *Let $\{a_n\}$ be a sequence in \mathbb{R} . Then the following are equivalent:*

- (a) $\overline{\lim} a_n = a$;
- (b) for every $b > a$, $a_n < b$ for all but finitely many n and for every $c < a$, $a_n > c$ for infinitely many n .

Proof. Assume $\overline{\lim} a_n = a$. Then for any $b > a$, there exists m such that $\sup_{n \geq m} a_n < b$. In particular, $a_n < b$ for all $n \geq m$. Since the sequence $\{\sup_{n \geq m} a_n\}$ is decreasing and convergent to a , it follows that $a \leq \sup_{n \geq m} a_n$ for all m . Hence if $c < a$, then for every m there exists $n \geq m$ such that $c < a_n$. This shows the implication (a) \implies (b). Conversely, assume that (b) holds. Then for every $b > a$, there exists m such that $a_n < b$ for all $n \geq m$. Hence $\sup_{n \geq m} a_n \leq b$. This implies that $\limsup a_n \leq b$ for every $b > a$ so that $\limsup a_n \leq a$. If for every $c < a$ and for every m there exists $n \geq m$ such that $a_n > c$, then for every m , $\sup_{n \geq m} a_n \geq c$. This gives $\limsup a_n > c$ and since this holds for every $c < a$, we have $\limsup a_n \geq a$. Thus $\limsup a_n = a$ and the implication (b) \implies (a) is proved. ■

As an exercise formulate and prove the corresponding statement for the lower limit. The basic properties of the upper and the lower limits are listed in the following proposition:

Proposition 11.14. *If $\{a_n\}$ and $\{b_n\}$ are sequences of real numbers, then:*

- (a) $\limsup(-a_n) = -\liminf a_n$ and $\liminf(-a_n) = -\limsup a_n$.
- (c) $\limsup(ca_n) = c\limsup a_n$ and $\liminf(ca_n) = c\liminf a_n$ for any $c > 0$.
- (d) $\limsup(a_n + b_n) \leq \limsup a_n + \limsup b_n$ and $\liminf a_n + \liminf b_n \leq \liminf(a_n + b_n)$.
- (e) $\liminf a_n \leq \limsup a_n$, with equality if and only if $\{a_n\}$ converges. In this case $\limsup a_n = \lim a_n$.
- (f) If $\{a_{n_k}\}$ is a subsequence of $\{a_n\}$, then $\liminf a_n \leq \liminf a_{n_k} \leq \limsup a_{n_k} \leq \limsup a_n$.

The proof is left as an exercise.

Theorem 11.15 (Bolzano-Weierstrass Theorem).

Let $\{a_n\}$ be a bounded sequence in \mathbb{R} . Then there is a subsequence that converges.

Proof. Set $a = \limsup a_n$. We will construct inductively a subsequence $\{a_{n_k}\}$ of $\{a_n\}$ which converges to a . In view of Proposition 11.13, there exists n_1 such that $a_{n_1} > a - 1$. Having obtained $n_1 < n_2 < \dots < n_k$ such that $a_{n_j} > a - 1/j$ for $1 \leq j \leq k$, we find, again by applying Proposition 11.13, $n_{k+1} > n_k$ such that $a_{n_{k+1}} > a - 1/(k+1)$. Hence $a \leq \liminf a_{n_k} \leq \limsup a_{n_k} \leq \limsup a_n = a$. So $\lim a_{n_k} = a$ and the proof is finished. ■

12 Problem Sheets

12.1 Problem Sheet 1

1. Check if the following functions are metrics on X .

- (a) $d(x, y) = |x^2 - y^2|$ for $x, y \in X = \mathbb{R}$
- (b) $d(x, y) = |x^2 - y^2|$ for $x, y \in X = (-\infty, 0]$
- (c) $d(x, y) = |\arctan x - \arctan y|$ for $x, y \in X = \mathbb{R}$

2. Let $X = \mathbb{R}^2$ and let d be the usual metric. Denote by $\mathbf{0} = (0, 0)$ and define


$$d_{\mathbf{0}}(x, y) = \begin{cases} 0 & \text{if } x = y; \\ d(x, \mathbf{0}) + d(\mathbf{0}, y) & \text{if } x \neq y. \end{cases}$$

Verify that $d_{\mathbf{0}}$ is a metric on X . (The metric $d_{\mathbf{0}}$ is called *the post office metric*).

3. Let $X = \mathbb{R}^2$. For $x = (x_1, x_2)$ and $y = (y_1, y_2)$ define

$$d(x, y) = \begin{cases} 1/2 & \text{if } x_1 = y_1 \text{ and } x_2 \neq y_2 \text{ or if } x_1 \neq y_1 \text{ and } x_2 = y_2; \\ 1 & \text{if } x_1 \neq y_1 \text{ and } x_2 \neq y_2; \\ 0 & \text{otherwise.} \end{cases}$$

Verify that d is a metric and that the rectangles in the figure have different “area” if d is used to measure the length of sides.



4.

(a) Show that if $0 \leq a \leq b$, then $\frac{a}{1+a} \leq \frac{b}{1+b}$.

(b) If $a, b, c \geq 0$ and $a \leq b + c$, show that $\frac{a}{1+a} \leq \frac{b}{1+b} + \frac{c}{1+c}$.

(c) Use (b) to show that if d is a metric on X , then

$$\tilde{d}(x, y) = \frac{d(x, y)}{1 + d(x, y)} \quad \text{for } x, y \in X$$

is a metric on X .

5. Let (X_i, d_i) be a metric space for $1 \leq i \leq n$ and let $X = \prod_{i=1}^n X_i$. Define

$$d_2(x, y) = \left[\sum_{i=1}^n d_i(x_i, y_i)^2 \right]^{1/2},$$

$$d_\infty(x, y) = \max\{d_i(x_i, y_i) \mid 1 \leq i \leq n\},$$

where $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n) \in X$. Verify that d_2 and d_∞ are metrics on X .

6. Fix a positive integer n . Denote by \mathcal{P}_n the set of all polynomials $p(x) = a_k x^k + a_{k-1} x^{k-1} + \dots + a_1 x + a_0$ with real coefficients a_i and $k \leq n$. For $p(x) = a_k x^k + a_{k-1} x^{k-1} + \dots + a_1 x + a_0 \in \mathcal{P}_n$ set

$$\|p\| = \max\{|a_0|, |a_1|, \dots, |a_k|\}.$$

Verify that $\|\cdot\|$ is a norm on \mathcal{P}_n .

7. Sketch the open ball $B(0, 1)$ in the metric space (\mathbb{R}^3, d_i) , where d_i is defined by

$$\begin{aligned} d_1(x, y) &= |x_1 - y_1| + |x_2 - y_2| + |x_3 - y_3| \\ d_2(x, y) &= \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2 + (x_3 - y_3)^2} \\ d_\infty(x, y) &= \max\{|x_1 - y_1|, |x_2 - y_2|, |x_3 - y_3|\}. \end{aligned}$$

for $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3) \in \mathbb{R}^3$.

8. Show that $\text{diam } B(x_0, r) \leq 2r$. Give an example showing that the strict inequality is possible.

12.2 Problem Sheet 2

1. Calculate $d(A, B)$ for the following pairs of subsets in \mathbb{R}^2 equipped with the standard metric:

- (a) $A = \{(x, 0) \mid x \in \mathbb{R}\}$ and $B = \{(x, 1) \mid x \in \mathbb{R}\}$.
- (b) A is the set of points on the x -axis whose x -coordinate satisfies $2n < x < 2n + 1$ for some $n \in \mathbb{Z}$ and B is the set of points on the line $y = 1$ for which $2m - 1 < x < 2m$ for some $m \in \mathbb{Z}$.
- (c) $A = B(x_0, r_0)$ and $B = B(x_1, r_1)$ where $x_0, x_1 \in \mathbb{R}^2$.

2. Let d and d' be two metrics on X such that

$$\alpha d(x, y) \leq d'(x, y) \leq \beta d(x, y)$$

for all $x, y \in X$ and positive constants α and β . Show that d and d' are equivalent. Give an example of X and two equivalent metrics in X for which

the above inequality does not hold. Use the above fact to show that

$$\begin{aligned}d_1(x, y) &= \sum_{i=1}^n |x_i - y_i| \\d_2(x, y) &= \left[\sum_{i=1}^n |x_i - y_i|^2 \right]^{1/2} \\d_\infty(x, y) &= \max\{|x_i - y_i| \mid i = 1, \dots, n\}.\end{aligned}$$

are equivalent on $X = \mathbb{R}^n$.

3 Consider the set $X = [-1, 1]$ as a subspace of \mathbb{R} a metric subspace of \mathbb{R} with the standard metric. Which of the following sets are open in X ? Which are open in \mathbb{R} ? Which are closed in X and which are closed in \mathbb{R} ?

- (a) $A = \{x \in X \mid 1/2 < |x| < 2\}$
- (b) $B = \{x \in X \mid 1/2 < |x| \leq 2\}$
- (c) $C = \{x \in \mathbb{R} \mid 1/2 \leq |x| < 1\}$
- (d) $D = \{x \in \mathbb{R} \mid 1/2 \leq |x| \leq 1\}$
- (e) $E = \{x \in \mathbb{R} \mid 0 < |x| \leq 1 \text{ and } 1/x \notin \mathbb{Z}\}$

4. Sketch (where possible) the following sets, and decide whether it is an open subset, or a closed subset, or neither of \mathbb{R}^2 with the standard metric:

- (a) $A = \{(x, y) \mid -1 < x \leq 1 \text{ and } -1 < y < 1\}$
- (b) $B = \{(x, y) \mid xy = 0\}$
- (c) $C = \{(x, y) \mid x \in \mathbb{Q}, y \in \mathbb{R}\}$
- (d) $D = \{(x, y) \mid -1 < x < 1 \text{ and } y = 0\}$
- (e) $E = \bigcup_{n=1}^{\infty} \{(x, y) \mid x = 1/n \text{ and } |y| \leq n\}$

5. Find the interior, the closure and the boundary of each of the following subsets of \mathbb{R}^2 with the standard metric:

- (a) $A = \{(x, y) \mid x > 0 \text{ and } y \neq 0\}$
- (b) $B = \{(x, y) \mid x \in \mathbb{N}, y \in \mathbb{R}\}$
- (c) $C = A \cup B$
- (d) $D = \{(x, y) \mid x \text{ is rational}\}$
- (e) $F = \{(x, y) \mid x \neq 0 \text{ and } y \leq 1/x\}$

6. Let A be a subset of a metric space X . Is the interior of A equal to the interior of the closure of A . Is the closure of the interior of A equal to the closure of A itself?

7 Consider a collection A_i of subsets of a metric space X . Show that

$$\begin{array}{ll} \bigcup_{i \in I} A_i^\circ \subset \left(\bigcup_{i \in I} A_i \right)^\circ & \overline{\bigcap_{i \in I} A_i} \subset \bigcap_{i \in I} \overline{A_i} \\ \left(\bigcap_{i \in I} A_i \right)^\circ \subset \bigcap_{i \in I} A_i^\circ & \bigcup_{i \in I} \overline{A_i} \subset \overline{\bigcup_{i \in I} A_i} \end{array}$$

8. Let U be open in X and let A be closed in X . Show that $U \setminus A$ is open in X and $A \setminus U$ is closed in X .

9. Let X and Y be metric spaces and A, B are non-empty subsets of X and Y , respectively.

- (a) Prove that if $A \times B$ is open in $X \times Y$, then A and B are open in X and Y , respectively.
- (b) Prove that if $A \times B$ is closed in $X \times Y$, then A and B are closed in X and Y , respectively.

12.3 Problem Sheet 3

- 1.** Show that A° and ∂A are disjoint, and $\overline{A} = A^\circ \cup \partial A$. Conclude A is open if and only if $\partial A = \overline{A} \setminus A$.
- 2.** Show that A is closed if and only if $\partial A \subset A$.
- 3.** Let $A, B \subset X$. Show that $\partial(A \cup B) \subset \partial A \cup \partial B$ and give an example in \mathbb{R} in which these sets are different. Show that if $\overline{A} \cap \overline{B} = \emptyset$, then $\partial(A \cup B) = \partial A \cup \partial B$.

4. Let X and Y be metric spaces and A, B are dense subsets of X and Y , respectively. Show that $A \times B$ is dense in $X \times Y$.

5. Let (X, d) be a metric space and let a be a fixed point of X . Show that

$$|d(x, a) - d(y, a)| \leq d(x, y) \quad (17)$$

for all $x, y \in X$.

6 Use (17) to show the following result. Let A be a dense subset of (X, d) . Show that a sequence $\{x_n\}$ of points on X converges to x if and only if

$$d(x_n, a) \rightarrow d(x, a)$$

for all $a \in A$.

7. Use (17) to show that the function $f : X \rightarrow \mathbb{R}$ defined by $f(x) = d(x, a)$, $x \in X$, is continuous.

8. Define $f : \mathbb{R} \rightarrow \mathbb{R}$ and $g : \mathbb{R} \rightarrow \mathbb{R}$ by

$$f(x) = \begin{cases} -1 & \text{if } x \geq 0 \\ 1 & \text{if } x < 0 \end{cases} \quad \text{and} \quad g(x) = \begin{cases} x & \text{if } x \geq 0 \\ 1 & \text{if } x < 0. \end{cases}$$

Show that f and g are not continuous at $x = 0$.

9. Let (X, d_X) and (Y, d_Y) be metric spaces. Show that $f : X \rightarrow Y$ is continuous if and only if

$$f(\overline{A}) \subset \overline{f(A)}$$

for all subsets A of X .

10. Let (X, d_X) and (Y, d_Y) be metric spaces. Show that $f : X \rightarrow Y$ is continuous if and only if

$$\overline{f^{-1}(B)} \subset f^{-1}(\overline{B}).$$

for all subsets B of Y .

12.4 Problem Sheet 4

1. Let Y be a subset of a metric space (X, d) . Show that

$$\overline{Y} = \{x \in X | d(x, Y) = 0\}.$$

Show that the function

$$f(x) = d(x, Y)$$

is continuous on X . Conclude that

- (a) $B(Y, \varepsilon) = \{x \in X \mid d(x, Y) < \varepsilon\}$ is open in X , and
- (b) $B[Y, \varepsilon] = \{x \in X \mid d(x, Y) \leq \varepsilon\}$ is closed in X .

2. Let A and B be disjoint non-empty closed subsets of a metric space X . Define

$$f(x) = \frac{d(x, A)}{d(x, A) + d(x, B)} \quad \text{for } x \in X.$$

Show that f is a continuous function on X whose image is in $[0, 1]$, and that $f(x) = 0$ if and only if $x \in A$ and $f(x) = 1$ if and only if $x \in B$. Next define sets $U = f^{-1}([0, 1/2))$ and $V = f^{-1}((1/2, 1])$. Show that U and V are open and disjoint, and that $A \subseteq U$, $B \subseteq V$.

3. Which of the following functions are uniformly continuous?

- (a) $f(x) = \sin x$ on $[0, \infty)$
- (b) $g(x) = \frac{1}{1-x}$ on $(0, 1)$
- (c) $h(x) = \sqrt{x}$ on $[0, \infty)$
- (d) $k(x) = \sin(1/x)$, on $(0, 1)$

4. Which of the following sequences converge uniformly on $[0, 1]$.

- (a) $f_n(x) = \frac{x}{1+nx}$
- (b) $g_n(x) = \frac{x^n}{1+x^n}$

5. Let (X, d) and (Y, ρ) be metric spaces and let $f : X \rightarrow Y$. Show that f is uniformly continuous if and only if for any two sequences $\{x_n\}$ and $\{y_n\}$ such that $d(x_n, y_n) \rightarrow 0$ it follows that $\rho(f(x_n), f(y_n)) \rightarrow 0$.

6. Suppose that $\{x_n\}$ is a sequence in a metric space (X, d) such that $d(x_n, x_{n+1}) \leq 2^{-n}$ for all $n \in \mathbb{N}$. Prove that $\{x_n\}$ is a Cauchy sequence.

7. Suppose that $\{x_n\}$ and $\{y_n\}$ are Cauchy sequences in a metric space (X, d) . Prove that the sequence of real numbers $\{d(x_n, y_n)\}$ converges.

8. Decide if the following metric spaces are complete:

- (a) $((0, \infty), d)$, where $d(x, y) = |x^2 - y^2|$ for $x, y \in (0, \infty)$.
- (b) $((-\pi/2, \pi/2), d)$, where $d(x, y) = |\tan x - \tan y|$ for $x, y \in (-\pi/2, \pi/2)$.

12.5 Problem Sheet 5

- Let $X = (0, 1]$ be equipped with the usual metric $d(x, y) = |x - y|$. Show that (X, d) is not complete. Let $\tilde{d}(x, y) = \left| \frac{1}{x} - \frac{1}{y} \right|$ for $x, y \in X$. Show that \tilde{d} is a metric on X that is equivalent to d , and that (X, \tilde{d}) is complete.
- Consider the space X consisting of all continuous functions $f : [a, b] \rightarrow \mathbb{R}$. For $f, g \in X$, define

$$d(f, g) = \int_a^b |f(x) - g(x)| \, dx.$$

Show that d is a metric on X . Is (X, d) complete?

- Cantor's Intersection Theorem** Let (X, d) be a complete metric space and let $\{F_n\}$ be a sequence of non-empty closed subsets of X such that $F_{n+1} \subseteq F_n$ for all n and $\text{diam}(F_n) \rightarrow 0$ as $n \rightarrow \infty$. Prove that $\bigcap_{n \in \mathbb{N}} F_n$ consists of exactly one point.

Show that, if any of the conditions,

- (i) (X, d) is complete, (ii) F_n is closed, (iii) $\text{diam}(F_n) \rightarrow 0$

is omitted, then $\bigcap_{n \in \mathbb{N}} F_n$ may be empty.

- Suppose that (X, d) and (Y, \tilde{d}) are metric spaces and that $f : X \rightarrow Y$ is a bijection such that both f and f^{-1} are uniformly continuous. Show that (X, d) is complete if and only if (Y, \tilde{d}) is complete.
- Let $\{f_n\}$ be a sequence of continuous functions $f_n : \mathbb{R} \rightarrow \mathbb{R}$ with the property that $\{f_n(x)\}$ is unbounded for all $x \in \mathbb{Q}$. Using Baire's theorem show that there is at least one $x \in \mathbb{Q}^c$ such that $\{f_n(x)\}$ is unbounded.
- Let (X, d) and (Y, \tilde{d}) be metric spaces such that (X, d) is complete. Let $\{f_n\}$ be a sequence of continuous functions from X to Y such that $\{f_n(x)\}$ converges for every $x \in X$. Using Baire's theorem show that for every $\varepsilon > 0$ there exist $k \in \mathbb{N}$ and a non-empty open subset U of X such that $\tilde{d}(f_n(x), f_m(x)) < \varepsilon$ for all $x \in U$ and all $n, m \geq k$.

12.6 Problem Sheet 6

1.

- (a) Let $f(x) = x^2$ for $x \in (0, a]$, and let $X = (0, a]$ with the usual metric. Find for what values of a is f a contraction. Show that $f : X \rightarrow X$ does not have a fixed point.
- (b) Let $f(x) = x + \frac{1}{x}$ for $x \geq 1$, and let $X = [1, \infty)$ with the usual metric d . Show that $f : X \rightarrow X$, that $d(f(x), f(y)) < d(x, y)$ for all $x \neq y$ but f does not have a fixed point.

Reconcile (a) and (b) with Banach fixed point theorem.

2. Consider $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by $f(x) = Ax$, where

$$A = \begin{bmatrix} 0.7 & 0.8 \\ 0.2 & -0.05 \end{bmatrix}.$$

Is f a contraction if \mathbb{R}^2 is equipped with the metric d_1, d_2, d_∞ ?

3. Consider the system of nonlinear equations

$$\begin{aligned} x_1 &= b_1 + \sin(a_{11}x_1) + \sin(a_{12}x_2) + \cdots + \sin(a_{1n}x_n), \\ x_2 &= b_2 + \sin(a_{21}x_1) + \sin(a_{22}x_2) + \cdots + \sin(a_{2n}x_n), \\ &\vdots \\ x_n &= b_n + \sin(a_{n1}x_1) + \sin(a_{n2}x_2) + \cdots + \sin(a_{nn}x_n), \end{aligned}$$

where a_{ik} , $1 \leq i, k \leq n$, and b_k , $1 \leq k \leq n$, are given real numbers. Show that the system has a unique solution $x = (x_1, \dots, x_n)$ if $\sum_{1 \leq i, k \leq n} a_{ik}^2 < 1$.

4. For $f \in C([0, 1], \mathbb{R})$, define

$$(Tf)(x) = e^x + \frac{1}{2} \cdot \frac{f(x)}{1 + f(x)^2}, \quad x \in [0, 1].$$

Show that $(Tf) \in C([0, 1], \mathbb{R})$ and that $T : C([0, 1], \mathbb{R}) \rightarrow C([0, 1], \mathbb{R})$ is a contraction. Use this fact to show that there exists exactly one function $f \in C([0, 1], \mathbb{R})$ such that

$$f(x)^3 - e^x f(x)^2 + \frac{1}{2} f(x) = e^x$$

for all $x \in [0, 1]$.

The next problems provide a different construction of the completion of (X, d) .

An **equivalence relation** on a set X is a relation \sim having the following three properties:

- (a) (Reflexivity) $x \sim x$ for every $x \in X$
- (b) (Symmetry) If $x \sim y$, then $y \sim x$
- (c) (Transitivity) If $x \sim y$ and $y \sim z$, then $x \sim z$.

The **equivalence class** determined by x , denoted by $[x]$, is defined by $[x] = \{y \in X \mid y \sim x\}$. We have $[x] = [y]$ if and only if $x \sim y$.

5. Let (X, d) be a metric space and let X^* be the set of Cauchy sequences $\mathbf{x} = \{x_n\}$ in (X, d) . Define a relation \sim in X^* by declaring $\mathbf{x} = \{x_n\} \sim \mathbf{y} = \{y_n\}$ to mean $d(x_n, y_n) \rightarrow 0$.

- (a) Show that \sim is an equivalence relation.
Denote by $[\mathbf{x}]$ the equivalence class of $\mathbf{x} \in X^*$, and let \tilde{X} denote the set of these equivalence classes.
- (b) Show that if $\mathbf{x} = \{x_n\}$ and $\mathbf{y} = \{y_n\} \in X^*$, then $\lim_{n \rightarrow \infty} d(x_n, y_n)$ exists. Show that if $\mathbf{x}' = \{x'_n\} \in [\mathbf{x}]$ and $\mathbf{y}' = \{y'_n\} \in [\mathbf{y}]$, then

$$\lim_{n \rightarrow \infty} d(x_n, y_n) = \lim_{n \rightarrow \infty} d(x'_n, y'_n).$$

For $[\mathbf{x}], [\mathbf{y}] \in \tilde{X}$, define

$$D([\mathbf{x}], [\mathbf{y}]) = \lim_{n \rightarrow \infty} d(x_n, y_n).$$

Note that the definition of D is unambiguous in view of the above equality.

- (c) Show that (\tilde{X}, D) is a complete metric space.
Hint: Let $[\mathbf{x}^n]$ be Cauchy in (\tilde{X}, D) . Then $\mathbf{x}^n = \{x_1^n, x_2^n, x_3^n, \dots\}$ is Cauchy in (X, d) . So for every $n \in \mathbb{N}$, there exists $k_n \in \mathbb{N}$ such that

$$d(x_m^n, x_{k_n}^n) < 1/n \quad \text{for all } m \geq k_n.$$

Set $\mathbf{x} = \{x_{k_1}^1, x_{k_2}^2, x_{k_3}^3, \dots\}$. Then show that \mathbf{x} is Cauchy in (X, d) and $D([\mathbf{x}^n], [\mathbf{x}]) \rightarrow 0$.

- (d) If $x \in X$, let $\varphi(x)$ be the equivalence class of the constant sequence $\mathbf{x} = (x, x, x, \dots)$. That is, $\varphi(x) = [\mathbf{x}] = [\{x, x, x, \dots\}]$. Show that $\varphi : X \rightarrow \varphi(X)$ is an isometry.
- (e) Show that $\varphi(X)$ is dense in (\tilde{X}, D) .
Hint: Let $[\mathbf{x}] \in \tilde{X}$ with $\mathbf{x} = \{x_1, x_2, x_3, \dots\}$. Denote by \mathbf{x}^n the constant sequence $\{x_n, x_n, x_n, \dots\}$ and show that $D([\mathbf{x}^n], [\mathbf{x}]) \rightarrow 0$.

12.7 Problem Sheet 7

1. Let (X, d) be a complete metric space and $f : X \rightarrow X$ be a function such that

$$d(f(x), f(y)) \leq \alpha d(x, y)$$

for all $x, y \in \overline{B}(x_0, r_0)$, where $0 < \alpha < 1$ and $d(x_0, f(x_0))/(1 - \alpha) = r_0$. Show that f has a unique fixed point $p \in \overline{B}(x_0, r_0)$.

2. (a) Let $I = [x_0 - a, x_0 + a]$ and let $r > 0$. If $f, g : I \rightarrow \mathbb{R}$ are continuous, define

$$d(f, g) = \sup\{e^{-r(x-x_0)}|f(x) - g(x)| \mid x \in I\}.$$

Show that d defines a metric on $C(I, \mathbb{R})$ which is equivalent to the supremum metric $\rho(f, g) = \sup\{|f(x) - g(x)| \mid x \in I\}$.

- (b) Let $K = I \times J$ where $I = [x_0 - a, x_0 + a]$, $J = [y_0 - b, y_0 + b]$, and let $f : K \rightarrow \mathbb{R}$ be a continuous function satisfying $|f(t, y_1) - f(t, y_2)| \leq \alpha|y_1 - y_2|$ for all $(t, y_1), (t, y_2) \in K$. Let $C = \sup\{|f(t, y)| \mid (t, y) \in K\}$, and let $\delta = \min\{a, b/C\}$. For a continuous function $y : [x_0 - \delta, x_0 + \delta] \rightarrow J$, set

$$Ty(x) = y_0 + \int_{x_0}^x f(t, y(t))dt, \quad x \in [x_0 - \delta, x_0 + \delta].$$

Show that $T : C([x_0 - \delta, x_0 + \delta], J) \rightarrow C([x_0 - \delta, x_0 + \delta], J)$ is a contraction with respect to the metric $d(y_1, y_2) = \sup\{e^{-2\alpha(x-x_0)}|f(x) - g(x)| \mid x \in I\}$.

Remark: The above device simplifies the last step in the prove of Picard's theorem given in lectures. Recall that in the last step we had to take $\delta > 0$ such that $C\delta < 1$ in order to guarantee that T is a contraction with respect to ρ . Using d we don't have to adjust δ .

3. Let $I = [x_0 - a, x_0 + a]$ and let $U = I \times \mathbb{R}$. Let $f : U \rightarrow \mathbb{R}$ be continuous.
 (a) Let $y_0, y_1 \in \mathbb{R}$. Show that a continuous function $y : I \rightarrow \mathbb{R}$ is a solution of

$$\begin{aligned} y''(x) &= f(x, y(x)), & x \in I \\ y(x_0) &= y_0 \\ y'(x_0) &= y_1 \end{aligned}$$

if and only if

$$y(x) = y_0 + (x - x_0)y_1 + \int_{x_0}^x (x - s)f(s, y(s))ds, \quad x \in I.$$

(b) Assume, in addition, that f satisfies the Lipschitz condition with respect to the second variable, $|f(x, y_1) - f(x, y_2)| \leq \alpha|y_1 - y_2|$ for all $(x, y_1), (x, y_2) \in U$, and some $\alpha > 0$. Prove that for given $y_0, y_1 \in \mathbb{R}$ there exists $\delta > 0$ such that the equation $y''(x) = f(x, y(x))$ has a unique solution $y : [x_0 - \delta, x_0 + \delta] \rightarrow \mathbb{R}$ satisfying $y(x_0) = y_0$ and $y'(x_0) = y_1$.

4. Let (X, d) be a metric space with the property that if Y is any non-empty closed subset of X and $f : Y \rightarrow Y$ is any contraction, then f has a fixed point. Show that (X, d) is complete.

Hint: Arguing by contradiction assume that there exists a Cauchy sequence which does not converge in X . We may assume that $x_n \neq x_m$ for all $n \neq m$. For $x \in X$, let $F(x) = \inf\{d(x, x_n) | n \in \mathbb{N}\}$. Show that $F(x) > 0$ for all x . Choose $\alpha \in (0, 1)$ and define sequence of integers $\{n_k\}$ as follows. Set $n_1 = 1$, let n_2 be an integer satisfying $n_2 > n_1$ and $d(x_i, x_j) \leq \alpha F(x_{n_1})$ for all integers $i, j \geq n_2$. If n_1, n_2, \dots, n_{k-1} are chosen, then $n_k > n_{k-1}$ is an integer such that $d(x_i, x_j) \leq \alpha F(x_{n_{k-1}})$ for all integers $i, j \geq n_k$. Now let $Y = \{x_{n_1}, x_{n_2}, x_{n_3}, \dots\}$ and let $f : Y \rightarrow Y$ be given by $f(x_{n_k}) = x_{n_{k+1}}$ for all $k \geq 1$. Show that Y is closed, f is a contraction but does not have a fixed point.

5. Let A be a dense subset of a metric space (X, d) , and let (Y, ρ) be complete. Consider a uniformly continuous function $f : A \rightarrow Y$. Show that there exists a unique uniformly continuous function $F : X \rightarrow Y$ such that $F(x) = f(x)$ for all $x \in A$.

12.8 Problem Sheet 8

1.

Show that if (X, d_X) and (Y, d_Y) are compact metric spaces, then the product metric space $(X \times Y, d)$ is compact. (Here d is the product metric).

2.

Show that if A_1, \dots, A_k are compact subsets of a metric space (X, d) , then $\bigcup_{i=1}^k A_i$ is compact.

3.

Which of the following subsets of \mathbb{R} and \mathbb{R}^2 are compact? (\mathbb{R} and \mathbb{R}^2 are

considered with the usual metrics).

- (a) $A = \mathbb{Q} \cap [0, 1]$
- (b) $B = \{(x, y) \in \mathbb{R}^2 | x^2 + y^2 = 1\}$
- (c) $C = \{(x, y) \in \mathbb{R}^2 | x^2 + y^2 < 1\}$
- (d) $D = \{x | |x| + |y| \leq 1\}$
- (e) $E = \{x | x \geq 1 \text{ and } 0 \leq y \leq 1/x\}$

4.

Consider (\mathbb{Q}, d) where d is the usual metric. Give an example of a set in this metric space that is closed and bounded but is not compact.

5.

Let A be a non-empty compact subset of a metric space (X, d) .

- (a) Let $x \in X$. Show that $d(x, A) = d(x, a)$ for some $a \in A$.
- (b) Let $U \subseteq X$ be open and $A \subseteq U$.

Show that there exists $\varepsilon > 0$ such that $S = \{x \in X | d(x, A) < \varepsilon\} \subseteq U$. Does this hold if A is only closed but not compact?

6.

Show that if A is a totally bounded subset of a metric space (X, d) , then for every $\varepsilon > 0$ there exists a finite subset $\{a_1, \dots, a_n\}$ of A such that $A \subseteq \bigcup_{i=1}^n B(a_i, \varepsilon)$.

7.

Show that a metric space (X, d) is totally bounded if and only if every sequence $\{x_n\} \subseteq X$ contains a Cauchy subsequence.

8.

Let X be a compact metric space and let \mathcal{U} be an open cover of X . Show that there exists a number $r > 0$ with the property: For every $x \in X$, there exists $U \in \mathcal{U}$ such that $B(x, r) \subseteq U$. The number r is called a **Lebesgue's number** of the cover \mathcal{U} .

12.9 Problem Sheet 9

1.

Show that (X, \mathcal{T}) is a topological space.

- (a) Let X be infinite set. Let

$$\mathcal{T} = \{A \subseteq X | A = \emptyset \text{ or } A = X \text{ or } X \setminus A \text{ is finite}\}.$$

This is called **co-finite topology** or **finite complement topology**.

(b) Let X be uncountable set. Define \mathcal{T} by

$$\mathcal{T} = \{A \subseteq X \mid A = \emptyset \text{ or } A = X \text{ or } X \setminus A \text{ is countable}\}.$$

This is called **co-countable topology** or **countable complement topology**.

(c) Let $X = \mathbb{R}$ and let

$$\mathcal{T} = \{A \subseteq \mathbb{R} \mid A = \emptyset \text{ or } A = \mathbb{R} \text{ or } A = (a, \infty) \text{ with } a \in \mathbb{R}\}.$$

2.

Let $\mathcal{B} = \{[a, b) \mid a, b \in \mathbb{R}\}$. Show that \mathcal{B} is a basis for a topology on \mathbb{R} . This topology, denoted \mathcal{T}_l , is called the **lower-limit topology** on \mathbb{R} . Show that the lower-limit topology is larger than the usual topology on \mathbb{R} . Find the closures of $[a, b)$, (a, b) , $(a, b]$ and $[a, b]$ in $(\mathbb{R}, \mathcal{T}_l)$.

3.

Let $\mathcal{T} = \{A \subseteq \mathbb{R} \mid 0 \notin A \text{ or } A = \mathbb{R}\}$. Show that \mathcal{T} is a topology on \mathbb{R} . What are the closed sets in $(\mathbb{R}, \mathcal{T})$? What is $\overline{\{1\}}$? Is this topology Hausdorff?

4.

Let A, B be subsets of a topological space (X, \mathcal{T}) . Show that $\overline{A \cap B} \subseteq \overline{A} \cap \overline{B}$ and $\overline{A \cup B} = \overline{A} \cup \overline{B}$.

5.

Let $A \subset X$ where (X, \mathcal{T}) is a topological space. Show that $\overline{X \setminus A} = X \setminus A^\circ$ and $(X \setminus A)^\circ = X \setminus \overline{A}$.

6.

Prove the following statements about continuous functions and discrete and indiscrete topological spaces.

- (a) If X is discrete, then every function $f : X \rightarrow Y$, where Y is any topological space, is continuous.
- (b) If X is not discrete, then there exists a topological space Y and a function $f : X \rightarrow Y$ that is not continuous.
- (c) If Y is an indiscrete topological space, then every function $f : X \rightarrow Y$, where X is any topological space, is continuous.
- (d) If Y is not indiscrete, then there exists a topological space X and a function $f : X \rightarrow Y$ that is not continuous.

7

Let X be infinite set and let \mathcal{T} be a co-finite topology on X . Show that any continuous function $f : X \rightarrow \mathbb{R}$ is constant. (\mathbb{R} is equipped with the usual metric topology).

8.

Let X and Y be topological spaces and let \mathcal{B} be a base of open sets for Y . Show that a function $f : X \rightarrow Y$ is continuous if and only if $f^{-1}(U)$ is open in X for every $U \in \mathcal{B}$.

9.

(a) Show that (a, b) is homeomorphic to (c, d) , (c, ∞) and \mathbb{R} . (All spaces are equipped with the usual topology).

(b) Show that $\mathbb{R}^2 \setminus \{(0, 0)\}$ is homeomorphic to $\mathbb{R}^2 \setminus \overline{B}((0, 0), 1)$.

10.

A **topological property** is a property that, if possessed by a topological space X , is also possessed by any topological space homeomorphic to X .

(a) Show that if $f : X \rightarrow Y$ is a homeomorphism, then $f(U)$ is open in Y for any open set $U \subseteq X$.

(b) Show that Hausdorff is a topological property.

(c) Is completeness a topological property of metric spaces?

12.10 Problem Sheet 10

1.

Let (X, \mathcal{T}) be a compact topological space and let A, B are closed subsets of (X, \mathcal{T}) . Show that $A \cup B$ is compact.

2.

Let $X = (0, 1)$ and let

$$\mathcal{T} = \{A \subseteq \mathbb{R} \mid A = \emptyset \text{ or } A = (0, 1) \text{ or } A = (0, 1 - 1/n) \text{ for } n \geq 2\}.$$

Show that every open set A different than X is compact. Is X compact?

3.

Let \mathcal{T} be a co-countable topology on \mathbb{R} , that is,

$$\mathcal{T} = \{A \subseteq \mathbb{R} \mid A = \emptyset \text{ or } \mathbb{R} \setminus A \text{ is countable}\}.$$

Is $[0, 1]$ compact in $(\mathbb{R}, \mathcal{T})$? What are the compact sets in $(\mathbb{R}, \mathcal{T})$?

4.

Consider $(\mathbb{R}, \mathcal{M})$, where \mathcal{M} is the usual metric topology in \mathbb{R} . Let

$$\mathcal{T} = \{A \subset \mathbb{R} \mid A = \emptyset \text{ or } \mathbb{R} \setminus A \text{ is compact in } (\mathbb{R}, \mathcal{M})\}.$$

(a) Show that \mathcal{T} is a topology on \mathbb{R} .

(b) Show that $(\mathbb{R}, \mathcal{T})$ is compact but not Hausdorff.

5.

Let A be closed and let B be compact in (X, \mathcal{T}) . Show that $A \cap B$ is compact.

6.

Let (X, \mathcal{T}) be compact and let $f : X \rightarrow \mathbb{R}$ be a continuous function. Show that f is bounded, that is, there is $M > 0$ such that $|f(x)| \leq M$ for all $x \in X$. Show that f attains its maximum and its minimum value.

7.

Let (X, \mathcal{T}) be compact and (Y, \mathcal{S}) Hausdorff. Show that if $f : X \rightarrow Y$ is a continuous bijection, then f is a homeomorphism.

8.

Let (X, \mathcal{T}) be a compact Hausdorff space and let \mathcal{T}' be another topology on X . Show that:

(a) if $\mathcal{T} \subseteq \mathcal{T}'$ but $\mathcal{T} \neq \mathcal{T}'$, then (X, \mathcal{T}') is Hausdorff but not compact.

(b) if $\mathcal{T}' \subseteq \mathcal{T}$ but $\mathcal{T} \neq \mathcal{T}'$, then (X, \mathcal{T}') is compact but not Hausdorff.

Hint: Use Problem 7.

9.

Let X be infinite set with the co-finite topology \mathcal{T} . Show that (X, \mathcal{T}) is connected.

10.

Is the topological space $(\mathbb{R}, \mathcal{T})$ from Problem 4 connected?

12.11 Problem Sheet 11

1.

Show that if A is a connected subspace of a topological space (X, \mathcal{T}) and if $A \subset B \subset \overline{A}$, then B is connected.

2.

If A and B are connected subsets of a topological space (X, \mathcal{T}) such that $\overline{A} \cap B \neq \emptyset$, then $A \cup B$ is connected.

3.

Let $\{A_n\}$ be a sequence of connected subsets of a topological space (X, \mathcal{T}) such that $A_n \cap A_{n+1} \neq \emptyset$ for all $n \in \mathbb{N}$. Prove that $\bigcup_{n \in \mathbb{N}} A_n$ is connected.

4.

Let (X, d) be a metric space. Call a function $f : X \rightarrow \mathbb{R}$ locally constant if for every x there exists $r > 0$ such that $f|_{B(x, r)} : B(x, r) \rightarrow \mathbb{R}$ is constant.

Show that if (X, d) is connected, then every locally constant function is constant.

5.

A metric space (X, d_X) is called a **chain connected** if for every pair x, y of points in X and every $\varepsilon > 0$, there are finitely many points $x = x_0, x_1, x_2, \dots, x_n = y$ such that $d_X(x_{i+1}, x_i) < \varepsilon$ for $i = 0, 1, \dots, n-1$. Prove that a compact, chain connected metric space is connected.

6.

A point $p \in X$ is called a *cut point* if $X \setminus \{p\}$ is disconnected. Show that the property of having a cut point is a topological property.

7.

Show that no two of the intervals (a, b) , $(a, b]$, and $[a, b]$ are homeomorphic.

8.

Show that \mathbb{R} and \mathbb{R}^2 are not homeomorphic (\mathbb{R} and \mathbb{R}^2 are equipped with the usual topologies) .

9.

Let A be countable set. Show that $\mathbb{R}^2 \setminus A$ is path connected.

10.

Show that if A is an open connected subset of \mathbb{R}^n , then A is path connected.

References

[Mu] J. R. Munkres, *Topology* , Prentice Hall, 2000.

[M] B. Mendelson, *Introduction to topology*, Dover Publications.

[C] F. Croom, *Introduction to topology* , .