

# MULTIVARIABLE AND VECTOR ANALYSIS

W W L CHEN

© W W L Chen, 1997, 2003.

This chapter is available free to all individuals, on the understanding that it is not to be used for financial gains, and may be downloaded and/or photocopied, with or without permission from the author.

However, this document may not be kept on any information storage and retrieval system without permission from the author, unless such system is not accessible to any individuals other than its owners.

## Chapter 6

### CHANGE OF VARIABLES

#### 6.1. Introduction

We consider double integrals of the type

$$\iint_S f(x, y) \, dx \, dy,$$

where  $f : S \rightarrow \mathbb{R}$  is continuous in the region  $S \subseteq \mathbb{R}^2$ . Suppose that we wish to make a substitution  $x = x(u, v)$  and  $y = y(u, v)$ . Then as in the theory of real valued functions of one real variable, we expect to have

$$(1) \quad \iint_S f(x, y) \, dx \, dy = \iint_{S^*} f(x(u, v), y(u, v)) J(u, v) \, du \, dv,$$

where  $J(u, v)$  is a term arising from the substitution, and where  $S^*$  is a region in the  $uv$ -plane corresponding to the region  $S$  in the  $xy$ -plane.

Before we go any further, let us consider the following simple example in the theory of real valued functions of one real variable.

EXAMPLE 6.1.1. To evaluate the integral

$$\int_0^1 \sqrt{1-x^2} \, dx,$$

we may use the substitution  $x = \cos \theta$ . Then  $dx = -\sin \theta \, d\theta$ , and  $\theta = \pi/2$  when  $x = 0$  and  $\theta = 0$  when  $x = 1$ . Hence

$$\int_0^1 \sqrt{1-x^2} \, dx = \int_{\pi/2}^0 \sqrt{1-\cos^2 \theta} (-\sin \theta) \, d\theta = \int_0^{\pi/2} \sin^2 \theta \, d\theta = \dots = \frac{\pi}{4}.$$

Clearly there is great potential for making a mistake by omitting a minus sign somewhere in the argument. We may instead proceed in the following way. Consider the function  $T : [0, \pi/2] \rightarrow [0, 1]$ , given by  $T(\theta) = \cos \theta$  for every  $\theta \in [0, \pi/2]$ . Clearly this function is one-to-one and onto. We may well suspect that for any continuous function  $f : [0, 1] \rightarrow \mathbb{R}$ , we have

$$\int_{[0,1]} f(x) \, dx = \int_{[0,\pi/2]} f(T(\theta)) \left| \frac{dT}{d\theta} \right| d\theta.$$

Indeed, this is true when  $f(x) = \sqrt{1-x^2}$  for every  $x \in [0, 1]$ , and is in fact true for every continuous function  $f : [0, 1] \rightarrow \mathbb{R}$ .

## 6.2. Planar Transformations

In this section, we shall study functions of the type  $T : S^* \rightarrow S$ , where  $S^*$  and  $S$  are sets in  $\mathbb{R}^2$ .

Suppose that  $S^* \subseteq \mathbb{R}^2$  is given. Consider a function  $T : S^* \rightarrow \mathbb{R}^2$ . Then for any  $(u, v) \in S^*$ , the point  $T(u, v)$  is called the image of  $(u, v)$  under  $T$ , and the set

$$T(S^*) = \{T(u, v) : (u, v) \in S^*\}$$

is called the range of the function  $T$ .

EXAMPLE 6.2.1. Suppose that  $S^* = [0, 1] \times [0, 2\pi]$ . The function  $T : S^* \rightarrow \mathbb{R}^2 : (r, \theta) \mapsto (r \cos \theta, r \sin \theta)$  has range


$$T(S^*) = \{(r \cos \theta, r \sin \theta) : (r, \theta) \in [0, 1] \times [0, 2\pi]\},$$

the closed disc in  $\mathbb{R}^2$  with radius 1 and centre  $(0, 0)$ .

EXAMPLE 6.2.2. Suppose that  $S^* = [0, 1] \times [0, 1]$ . Consider the function  $T : S^* \rightarrow \mathbb{R}^2$ , given for every  $(u, v) \in S^*$  by  $T(u, v) = (x, y)$ , where

$$\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} \quad \text{and} \quad x + iy = (1 + i)(u + iv).$$

Clearly  $T(S^*)$  is a square with vertices  $(0, 0)$ ,  $(0, 2)$  and  $(\pm 1, 1)$ .



DEFINITION. A function  $T : S^* \rightarrow \mathbb{R}^2$  is said to be one-to-one if for every  $(u', v'), (u'', v'') \in S^*$ , we have  $(u', v') = (u'', v'')$  whenever  $T(u', v') = T(u'', v'')$ . In other words, no two distinct  $(u', v'), (u'', v'') \in S^*$  can have the same image.

EXAMPLE 6.2.3. In Example 6.2.1, the function  $T : S^* \rightarrow \mathbb{R}^2$  is not one-to-one. It is easy to check that  $T(0, 0) = T(0, \pi) = 0$ .

EXAMPLE 6.2.4. In Example 6.2.2, it is easily seen that the function  $T : S^* \rightarrow \mathbb{R}^2$  is a rotation about the point  $(0, 0)$  through an angle  $\pi/4$ , together with a magnification by  $\sqrt{2}$  about the point  $(0, 0)$ . It is clear that if  $T(u', v') = (x', y')$ ,  $T(u'', v'') = (x'', y'')$  and  $(x', y') = (x'', y'')$ , then since the matrix  $\begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}$  is invertible, we must have  $(u', v') = (u'', v'')$ . Hence  $T : S^* \rightarrow \mathbb{R}^2$  is one-to-one.

EXAMPLE 6.2.5. Let us vary Example 6.2.1, and consider the rectangle  $S^* = (0, 1] \times [0, 2\pi)$ . Then the function  $T : S^* \rightarrow \mathbb{R}^2 : (r, \theta) \mapsto (r \cos \theta, r \sin \theta)$  is one-to-one, with range

$$T(S^*) = \{(r \cos \theta, r \sin \theta) : (r, \theta) \in (0, 1] \times [0, 2\pi)\},$$

the closed disc in  $\mathbb{R}^2$  with radius 1 and centre  $(0, 0)$ , punctured at the point  $(0, 0)$ .

In practice, our problem is the following: Given a region  $S \subseteq \mathbb{R}^2$ , we need to find a region  $S^* \subseteq \mathbb{R}^2$  and a one-to-one function  $T : S^* \rightarrow \mathbb{R}^2$  such that  $T(S^*) = S$ . It is therefore convenient to consider functions of the type  $T : S^* \rightarrow S$  instead of  $T : S^* \rightarrow \mathbb{R}^2$ . Note that this change does not affect our definition of one-to-one functions given earlier, and we can rephrase it as follows.

DEFINITION. A function  $T : S^* \rightarrow S$  is said to be one-to-one if for every  $(u', v'), (u'', v'') \in S^*$ , we have  $(u', v') = (u'', v'')$  whenever  $T(u', v') = T(u'', v'')$ .

To characterize the property  $T(S^*) = S$ , we have the following definition.

DEFINITION. A function  $T : S^* \rightarrow S$  is said to be onto if for every  $(x, y) \in S$ , there exists  $(u, v) \in S^*$  such that  $T(u, v) = (x, y)$ .

EXAMPLE 6.2.6. Suppose that  $S^* = [0, 1] \times [0, 2\pi]$  and  $S = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 \leq 1\}$ . Then the function  $T : S^* \rightarrow \mathbb{R}^2 : (r, \theta) \mapsto (r \cos \theta, r \sin \theta)$  is onto.

EXAMPLE 6.2.7. In Example 6.2.2, if we change the codomain of the function  $T$  to the closed rectangle  $S$  with vertices  $(0, 0)$ ,  $(0, 2)$  and  $(\pm 1, 1)$ , then  $T : S^* \rightarrow S$  is one-to-one and onto.

EXAMPLE 6.2.8. Suppose that  $S^* = (0, 1] \times [0, 2\pi)$  and  $S = \{(x, y) \in \mathbb{R}^2 : 0 < x^2 + y^2 \leq 1\}$ . Then the function  $T : S^* \rightarrow \mathbb{R}^2 : (r, \theta) \mapsto (r \cos \theta, r \sin \theta)$  is one-to-one and onto.

### 6.3. The Jacobian

In this section, we shall turn our attention to the term  $J(u, v)$  in the expression (1). We shall begin by studying the special case when  $f(x, y) = 1$  for every  $(x, y) \in S$ , so that the expression (1) is of the form

$$(2) \quad \iint_S dx dy = \iint_{S^*} J(u, v) du dv.$$

Note that the left hand side of (2) represents the area of  $S$ .

EXAMPLE 6.3.1. We study rectangular coordinates  $(x, y)$  and polar coordinates  $(r, \theta)$ , related by

$$x = r \cos \theta \quad \text{and} \quad y = r \sin \theta.$$

Consider the unit disc

$$S = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 \leq 1\}.$$

Let

$$S^* = \{(r, \theta) \in \mathbb{R}^2 : 0 \leq r \leq 1 \text{ and } 0 \leq \theta \leq 2\pi\}.$$

Then the function

$$T : S^* \rightarrow \mathbb{R}^2 : (r, \theta) \mapsto (r \cos \theta, r \sin \theta)$$

is one-to-one and onto, provided that we remove the lines  $r = 0$  and  $\theta = 2\pi$  from  $S^*$  and the point  $(x, y) = (0, 0)$  from  $S$ . Clearly

$$\iint_S dx dy = \pi.$$

On the other hand,

$$\frac{\partial(x, y)}{\partial(r, \theta)} = \det \begin{pmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{pmatrix} = \det \begin{pmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{pmatrix} = r.$$

By Fubini's theorem, we have

$$\iint_{S^*} \left| \frac{\partial(x, y)}{\partial(r, \theta)} \right| dr d\theta = \iint_{S^*} r dr d\theta = \int_0^{2\pi} \left( \int_0^1 r dr \right) d\theta = \pi.$$

Here we detect a small problem. The function  $T : S^* \rightarrow S$  may not be one-to-one and onto, unless we remove some subsets of  $S^*$  and  $S$ . Let us first of all consider such exceptional subsets.

**DEFINITION.** Suppose that  $S \subseteq \mathbb{R}^2$ . Then we say that a subset  $E \subseteq S$  is exceptional if  $E$  is contained in a finite union of curves of type 1 or 2 in  $S$ .

**DEFINITION.** Suppose that  $S^*$  and  $S$  are sets in  $\mathbb{R}^2$ . Then we say that a function  $T : S^* \rightarrow S$  is essentially one-to-one and onto if there exist exceptional subsets  $E^* \subseteq S^*$  and  $E \subseteq S$  such that the function  $T : S^* \setminus E^* \rightarrow S \setminus E$  is one-to-one and onto.

**EXAMPLE 6.3.2.** In Example 6.3.1, we can take

$$E^* = \{(r, \theta) \in S^* : r = 0 \text{ or } \theta = 2\pi\} \quad \text{and} \quad E = \{(0, 0)\}.$$

Then

$$S^* \setminus E^* = \{(r, \theta) \in \mathbb{R}^2 : 0 < r \leq 1 \text{ and } 0 \leq \theta < 2\pi\} \quad \text{and} \quad S \setminus E = \{(x, y) \in \mathbb{R}^2 : 0 < x^2 + y^2 \leq 1\}.$$

We have shown in Example 6.2.8 that the function

$$T : S^* \setminus E^* \rightarrow S \setminus E : (r, \theta) \mapsto (r \cos \theta, r \sin \theta)$$

is one-to-one and onto.

**THEOREM 6A.** Suppose that  $S^* \subseteq \mathbb{R}^2$  and  $S \subseteq \mathbb{R}^2$  are elementary regions, and that the function  $T : S^* \rightarrow S$ , where  $T(u, v) = (x(u, v), y(u, v))$  for every  $(u, v) \in S^*$ , has continuous partial derivatives. Suppose further that  $T : S^* \rightarrow S$  is essentially one-to-one and onto. Then for any Riemann integrable function  $f : S \rightarrow \mathbb{R}$ , we have


$$\iint_S f(x, y) dx dy = \iint_{S^*} f(x(u, v), y(u, v)) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv,$$

where the Jacobian determinant of  $T$  is given by

$$\frac{\partial(x, y)}{\partial(u, v)} = \det \begin{pmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{pmatrix}.$$

REMARK. An alternative approach is to restrict  $S^*$  to open sets and take  $S = T(S^*)$ . However, the problem with this approach is that since  $S$  is usually given, it may not always be possible to find a suitable open set  $S^*$  such that  $T(S^*) = S$  precisely, and we face the same difficulty, albeit in a slightly different disguise.


We shall not prove Theorem 6A. Instead we shall only briefly indicate the ideas. Consider a small rectangle



in  $S^*$  with bottom left vertex  $(u_0, v_0)$  and top right vertex  $(u, v) = (u_0 + \Delta u, v_0 + \Delta v)$ . We know from our discussion of differentiability in Chapter 2 that a good approximation to  $T(u, v)$  is given by the linear mapping

$$(4) \quad T(u_0, v_0) + (\mathbf{D}T)(u_0, v_0) \begin{pmatrix} u - u_0 \\ v - v_0 \end{pmatrix} = T(u_0, v_0) + (\mathbf{D}T)(u_0, v_0) \begin{pmatrix} \Delta u \\ \Delta v \end{pmatrix},$$

where  $(\mathbf{D}T)(u_0, v_0)$  denotes the total derivative of  $T$  at  $(u_0, v_0)$ . The image of the rectangle (3) under the mapping (4) is a parallelogram



with one vertex at  $T(u_0, v_0)$  and adjacent sides, corresponding to  $\Delta u$  and  $\Delta v$ , given by the vectors

$$(\mathbf{D}T)(u_0, v_0) \begin{pmatrix} \Delta u \\ 0 \end{pmatrix} = \begin{pmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{pmatrix} \begin{pmatrix} \Delta u \\ 0 \end{pmatrix} = \Delta u \begin{pmatrix} \frac{\partial x}{\partial u} \\ \frac{\partial y}{\partial u} \end{pmatrix}$$

and

$$(\mathbf{D}T)(u_0, v_0) \begin{pmatrix} 0 \\ \Delta v \end{pmatrix} = \begin{pmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{pmatrix} \begin{pmatrix} 0 \\ \Delta v \end{pmatrix} = \Delta v \begin{pmatrix} \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial v} \end{pmatrix}$$

respectively (here the partial derivatives are supposed to be evaluated at  $(u_0, v_0)$ ). Clearly the area of this parallelogram is

$$\left| \det \begin{pmatrix} \Delta u \frac{\partial x}{\partial u} & \Delta v \frac{\partial x}{\partial v} \\ \Delta u \frac{\partial y}{\partial u} & \Delta v \frac{\partial y}{\partial v} \end{pmatrix} \right| = \left| \det \begin{pmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{pmatrix} \right| \Delta u \Delta v = \left| \frac{\partial(x, y)}{\partial(u, v)} \right| \Delta u \Delta v.$$

EXAMPLE 6.3.3. Suppose that  $0 \leq \theta_1 < \theta_2 \leq 2\pi$ . Consider the region

$$S = \{(r \cos \theta, r \sin \theta) : \theta_1 \leq \theta \leq \theta_2 \text{ and } g_1(\theta) \leq r \leq g_2(\theta)\},$$

where  $g_1(\theta) \leq g_2(\theta)$  for every  $\theta \in [0, 2\pi]$ . Writing  $(x, y) = (r \cos \theta, r \sin \theta)$ , we can calculate the area of  $S$  as

$$A(S) = \iint_S dx dy = \iint_{S^*} \left| \frac{\partial(x, y)}{\partial(r, \theta)} \right| dr d\theta,$$

where

$$S^* = \{(r, \theta) \in \mathbb{R}^2 : \theta_1 \leq \theta \leq \theta_2 \text{ and } g_1(\theta) \leq r \leq g_2(\theta)\}.$$

Then

$$A(S) = \int_{\theta_1}^{\theta_2} \left( \int_{g_1(\theta)}^{g_2(\theta)} \left| \frac{\partial(x, y)}{\partial(r, \theta)} \right| dr \right) d\theta = \int_{\theta_1}^{\theta_2} \left( \int_{g_1(\theta)}^{g_2(\theta)} r dr \right) d\theta = \int_{\theta_1}^{\theta_2} \frac{g_2^2(\theta) - g_1^2(\theta)}{2} d\theta.$$

EXAMPLE 6.3.4. Consider the region

$$S = \{(x, y) \in \mathbb{R}^2 : x, y \geq 0 \text{ and } 1 \leq x^2 + y^2 \leq 4\}$$

in the first quadrant between the two circles of radii 1 and 2 centred at  $(0, 0)$ . We showed in Example 5.4.4 that

$$\iint_S (x^2 + y^2) dx dy = \frac{15\pi}{8}.$$

We now use the substitution  $x = r \cos \theta$  and  $y = r \sin \theta$  instead, and consider the rectangle

$$S^* = \{(r, \theta) \in \mathbb{R}^2 : 1 \leq r \leq 2 \text{ and } 0 \leq \theta \leq \pi/2\}.$$

Note that the function

$$T : S^* \rightarrow S : (r, \theta) \mapsto (r \cos \theta, r \sin \theta)$$


is one-to-one and onto. By Theorem 6A, we have

$$\iint_S (x^2 + y^2) \, dx \, dy = \iint_{S^*} (r^2 \cos^2 \theta + r^2 \sin^2 \theta) \left| \frac{\partial(x, y)}{\partial(r, \theta)} \right| \, dr \, d\theta = \iint_{S^*} r^3 \, dr \, d\theta = \dots = \frac{15\pi}{8}.$$

EXAMPLE 6.3.5. Consider the parallelogram  $S$  with vertices  $(\pi, 0)$ ,  $(\pi, \pi)$  and  $(0, \pm\pi/2)$ . We showed in Example 5.4.2 that

$$\iint_S \sin x \cos y \, dx \, dy = \frac{8}{3}.$$

We now use the substitution  $x = 2u$  and  $y = u + v$  instead, and consider the rectangle  $S^*$  with vertices  $(0, \pm\pi/2)$  and  $(\pi/2, \pm\pi/2)$ .



Note that the function

$$T : S^* \rightarrow S : (u, v) \mapsto (2u, u + v)$$

is one-to-one and onto. By Theorem 6A, we have

$$\begin{aligned} \iint_S \sin x \cos y \, dx \, dy &= \iint_{S^*} \sin 2u \cos(u + v) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| \, du \, dv = 2 \iint_{S^*} \sin 2u (\cos u \cos v - \sin u \sin v) \, du \, dv \\ &= 2 \left( \int_0^{\pi/2} \sin 2u \cos u \, du \right) \left( \int_{-\pi/2}^{\pi/2} \cos v \, dv \right) - 2 \left( \int_0^{\pi/2} \sin 2u \sin u \, du \right) \left( \int_{-\pi/2}^{\pi/2} \sin v \, dv \right) \\ &= 2 \left( \int_0^{\pi/2} \sin 2u \cos u \, du \right) \left( \int_{-\pi/2}^{\pi/2} \cos v \, dv \right) = \dots = \frac{8}{3}. \end{aligned}$$

EXAMPLE 6.3.6. Consider the repeated integral

$$\int_0^1 \left( \int_0^1 (x^2 + y^2)^{3/2} \, dx \right) \, dy = \iint_R (x^2 + y^2)^{3/2} \, dx \, dy,$$

where  $R = [0, 1] \times [0, 1]$ . By splitting  $R$  into two triangles along the line  $y = x$ , it is easy to see that

$$\begin{aligned} \iint_R (x^2 + y^2)^{3/2} dx dy &= \int_0^1 \left( \int_0^x (x^2 + y^2)^{3/2} dy \right) dx + \int_0^1 \left( \int_0^y (x^2 + y^2)^{3/2} dx \right) dy \\ &= 2 \iint_S (x^2 + y^2)^{3/2} dx dy, \end{aligned}$$

where  $S$  is the triangular region with vertices  $(0, 0)$ ,  $(1, 0)$  and  $(1, 1)$ . We use the substitution  $x = r \cos \theta$  and  $y = r \sin \theta$ , and consider the region

$$S^* = \{(r, \theta) \in \mathbb{R}^2 : 0 \leq \theta \leq \pi/4 \text{ and } 0 \leq r \leq \sec \theta\}.$$

Note that the function

$$T : S^* \rightarrow S : (r, \theta) \mapsto (r \cos \theta, r \sin \theta)$$

is one-to-one and onto, provided that we remove the line  $r = 0$  from  $S^*$  and the point  $(x, y) = (0, 0)$  from  $S$ . By Theorem 6A, we have

$$\begin{aligned} \iint_S (x^2 + y^2)^{3/2} dx dy &= \iint_{S^*} (r^2 \cos^2 \theta + r^2 \sin^2 \theta)^{3/2} \left| \frac{\partial(x, y)}{\partial(r, \theta)} \right| dr d\theta = \int_0^{\pi/4} \left( \int_0^{\sec \theta} r^4 dr \right) d\theta \\ &= \frac{1}{5} \int_0^{\pi/4} \sec^5 \theta d\theta = \left[ \frac{1}{20} \tan \theta + \sec^3 \theta + \frac{3}{40} \tan \theta \sec \theta + \frac{3}{40} \log |\tan \theta + \sec \theta| \right]_0^{\pi/4} \\ &= \frac{7\sqrt{2}}{40} + \frac{3}{40} \log(1 + \sqrt{2}). \end{aligned}$$

#### 6.4. Triple Integrals

Our discussion so far can be extended to triple integrals with no extra complication.

**DEFINITION.** Suppose that  $S \subseteq \mathbb{R}^3$ . Then we say that a subset  $E \subseteq S$  is exceptional if  $E$  is contained in a finite union of surfaces in  $S$  representing one of the three variables as a continuous function of the other two variables.

**DEFINITION.** Suppose that  $S^*$  and  $S$  are sets in  $\mathbb{R}^3$ . Then we say that a function  $T : S^* \rightarrow S$  is essentially one-to-one and onto if there exist exceptional subsets  $E^* \subseteq S^*$  and  $E \subseteq S$  such that the function  $T : S^* \setminus E^* \rightarrow S \setminus E$  is one-to-one and onto.

**THEOREM 6B.** Suppose that  $S^* \subseteq \mathbb{R}^3$  and  $S \subseteq \mathbb{R}^3$  are elementary regions, and that the function  $T : S^* \rightarrow S$ , where  $T(u, v, w) = (x(u, v, w), y(u, v, w), z(u, v, w))$  for every  $(u, v, w) \in S^*$ , has continuous partial derivatives. Suppose further that  $T : S^* \rightarrow S$  is essentially one-to-one and onto. Then for any Riemann integrable function  $f : S \rightarrow \mathbb{R}$ , we have

$$\iiint_S f(x, y, z) dx dy dz = \iiint_{S^*} f(x(u, v, w), y(u, v, w), z(u, v, w)) \left| \frac{\partial(x, y, z)}{\partial(u, v, w)} \right| du dv dw,$$

where the Jacobian determinant of  $T$  is given by

$$\frac{\partial(x, y, z)}{\partial(u, v, w)} = \det \begin{pmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} & \frac{\partial x}{\partial w} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} & \frac{\partial y}{\partial w} \\ \frac{\partial z}{\partial u} & \frac{\partial z}{\partial v} & \frac{\partial z}{\partial w} \end{pmatrix}.$$



REMARKS. (1) The image under  $T$  of a rectangular box in  $S^*$  with side lengths  $\Delta u$ ,  $\Delta v$  and  $\Delta w$  can be approximated by a parallelepiped in  $S$  with volume

$$\left| \det \begin{pmatrix} \Delta u \frac{\partial x}{\partial u} & \Delta v \frac{\partial x}{\partial v} & \Delta w \frac{\partial x}{\partial w} \\ \Delta u \frac{\partial y}{\partial u} & \Delta v \frac{\partial y}{\partial v} & \Delta w \frac{\partial y}{\partial w} \\ \Delta u \frac{\partial z}{\partial u} & \Delta v \frac{\partial z}{\partial v} & \Delta w \frac{\partial z}{\partial w} \end{pmatrix} \right| = \left| \det \begin{pmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} & \frac{\partial x}{\partial w} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} & \frac{\partial y}{\partial w} \\ \frac{\partial z}{\partial u} & \frac{\partial z}{\partial v} & \frac{\partial z}{\partial w} \end{pmatrix} \right| \Delta u \Delta v \Delta w = \left| \frac{\partial(x, y, z)}{\partial(u, v, w)} \right| \Delta u \Delta v \Delta w.$$

(2) Cylindrical coordinates  $(r, \theta, z)$  are related to rectangular coordinates  $(x, y, z)$  in  $\mathbb{R}^3$  by

$$x = r \cos \theta, \quad y = r \sin \theta, \quad z = z.$$

Hence

$$\frac{\partial(x, y, z)}{\partial(r, \theta, z)} = \det \begin{pmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} & \frac{\partial x}{\partial z} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} & \frac{\partial y}{\partial z} \\ \frac{\partial z}{\partial r} & \frac{\partial z}{\partial \theta} & \frac{\partial z}{\partial z} \end{pmatrix} = \det \begin{pmatrix} \cos \theta & -r \sin \theta & 0 \\ \sin \theta & r \cos \theta & 0 \\ 0 & 0 & 1 \end{pmatrix} = r,$$


so that

$$\iiint_S f(x, y, z) \, dx \, dy \, dz = \iiint_{S^*} f(r \cos \theta, r \sin \theta, z) r \, dr \, d\theta \, dz.$$

Here, we essentially leave the variable  $z$  alone, and replace the variables  $x$  and  $y$  by the variables  $r$  and  $\theta$  in the same way as with rectangular and polar coordinates in  $\mathbb{R}^2$ .

(3) Spherical coordinates  $(r, \phi, \theta)$  are related to rectangular coordinates  $(x, y, z)$  in  $\mathbb{R}^3$  by

$$x = r \sin \phi \cos \theta, \quad y = r \sin \phi \sin \theta, \quad z = r \cos \phi.$$



We have

$$\begin{aligned}
 \frac{\partial(x, y, z)}{\partial(r, \phi, \theta)} &= \det \begin{pmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \phi} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \phi} & \frac{\partial y}{\partial \theta} \\ \frac{\partial z}{\partial r} & \frac{\partial z}{\partial \phi} & \frac{\partial z}{\partial \theta} \end{pmatrix} = \det \begin{pmatrix} \sin \phi \cos \theta & r \cos \phi \cos \theta & -r \sin \phi \sin \theta \\ \sin \phi \sin \theta & r \cos \phi \sin \theta & r \sin \phi \cos \theta \\ \cos \phi & -r \sin \phi & 0 \end{pmatrix} \\
 &= (\cos \phi) \det \begin{pmatrix} r \cos \phi \cos \theta & -r \sin \phi \sin \theta \\ r \cos \phi \sin \theta & r \sin \phi \cos \theta \end{pmatrix} + (r \sin \phi) \det \begin{pmatrix} \sin \phi \cos \theta & -r \sin \phi \sin \theta \\ \sin \phi \sin \theta & r \sin \phi \cos \theta \end{pmatrix} \\
 &= (\cos \phi)(r^2 \cos \phi \sin \phi \cos^2 \theta + r^2 \cos \phi \sin \phi \sin^2 \theta) + (r \sin \phi)(r \sin^2 \phi \cos^2 \theta + r \sin^2 \phi \sin^2 \theta) \\
 &= r^2 \sin \phi \cos^2 \phi + r^2 \sin \phi \sin^2 \phi = r^2 \sin \phi,
 \end{aligned}$$

so that

$$\iiint_S f(x, y, z) \, dx dy dz = \iiint_{S^*} f(r \sin \phi \cos \theta, r \sin \phi \sin \theta, r \cos \phi) r^2 \sin \phi \, dr d\phi d\theta.$$

EXAMPLE 6.4.1. Consider the integral

$$\iiint_S (x^2 + y^2 + z^2)^{1/2} \, dx dy dz,$$

where

$$S = \{(x, y, z) \in \mathbb{R}^3 : y, z \geq 0 \text{ and } 4 \leq x^2 + y^2 + z^2 \leq 9\}.$$

We use spherical coordinates and the substitution

$$x = r \sin \phi \cos \theta, \quad y = r \sin \phi \sin \theta, \quad z = r \cos \phi,$$

and consider the set

$$S^* = \left\{ (r, \phi, \theta) \in \mathbb{R}^3 : r \in [2, 3], \phi \in \left[0, \frac{\pi}{2}\right] \text{ and } \theta \in [0, \pi] \right\}.$$

Note that the function

$$T : S^* \rightarrow S : (r, \phi, \theta) \mapsto (r \sin \phi \cos \theta, r \sin \phi \sin \theta, r \cos \phi)$$

is one-to-one and onto. By Theorem 6B and noting Remark (3) above, we have

$$\begin{aligned}
 \iiint_S (x^2 + y^2 + z^2)^{1/2} \, dx dy dz &= \iiint_{S^*} (r^2 \sin^2 \phi \cos^2 \theta + r^2 \sin^2 \phi \sin^2 \theta + r^2 \cos^2 \phi)^{1/2} r^2 \sin \phi \, dr d\phi d\theta \\
 &= \iiint_{S^*} r^3 \sin \phi \, dr d\phi d\theta = \left( \int_2^3 r^3 \, dr \right) \left( \int_0^{\pi/2} \sin \phi \, d\phi \right) \left( \int_0^\pi d\theta \right) = \frac{65\pi}{4}.
 \end{aligned}$$

EXAMPLE 6.4.2. Consider the integral

$$\iiint_S (x^2 + y^2 + z^2)^{5/2} dx dy dz,$$

where

$$S = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 + z^2 \leq 1\}.$$

We use spherical coordinates and the substitution

$$x = r \sin \phi \cos \theta, \quad y = r \sin \phi \sin \theta, \quad z = r \cos \phi,$$

and consider the set

$$S^* = \{(r, \phi, \theta) \in \mathbb{R}^3 : r \in [0, 1], \phi \in [0, \pi] \text{ and } \theta \in [0, 2\pi]\}.$$

Note that the function

$$T : S^* \rightarrow S : (r, \phi, \theta) \mapsto (r \sin \phi \cos \theta, r \sin \phi \sin \theta, r \cos \phi)$$

is one-to-one and onto, provided that we remove the planes  $r = 0$  and  $\theta = 2\pi$  from  $S^*$  and the point  $(0, 0, 0)$  from  $S$ . By Theorem 6B and noting Remark (3) above, we have

$$\iiint_S (x^2 + y^2 + z^2)^{5/2} dx dy dz = \iiint_{S^*} r^7 \sin \phi dr d\phi d\theta = \left( \int_0^1 r^7 dr \right) \left( \int_0^\pi \sin \phi d\phi \right) \left( \int_0^{2\pi} d\theta \right) = \frac{\pi}{2}.$$

EXAMPLE 6.4.3. Consider the integral

$$\iiint_S (x^2 + y^2)^{1/2} z dx dy dz,$$

where

$$S = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 \leq 1 \text{ and } 0 \leq z \leq 1\}.$$

We use cylindrical coordinates and the substitution

$$x = r \cos \theta, \quad y = r \sin \theta, \quad z = z,$$

and consider the set

$$S^* = \{(r, \theta, z) \in \mathbb{R}^3 : r \in [0, 1], \theta \in [0, 2\pi] \text{ and } z \in [0, 1]\}.$$

Note that the function

$$T : S^* \rightarrow S : (r, \theta, z) \mapsto (r \cos \theta, r \sin \theta, z)$$

is one-to-one and onto, provided that we remove the planes  $r = 0$  and  $\theta = 2\pi$  from  $S^*$  and the point  $(0, 0, 0)$  from  $S$ . By Theorem 6B and noting Remark (2) above, we have

$$\begin{aligned} \iiint_S (x^2 + y^2)^{1/2} z dx dy dz &= \iiint_{S^*} (r^2 \cos^2 \theta + r^2 \sin^2 \theta)^{1/2} z r dr d\theta dz \\ &= \left( \int_0^1 r^2 dr \right) \left( \int_0^{2\pi} d\theta \right) \left( \int_0^1 z dz \right) = \frac{\pi}{3}. \end{aligned}$$

## PROBLEMS FOR CHAPTER 6

1. By changing to polar coordinates, evaluate the integral  $\int_0^1 \left( \int_0^{\sqrt{1-x^2}} e^{-x^2-y^2} dy \right) dx$ .
2. By writing  $2x = r - s$  and  $2y = r + s$ , evaluate the integral  $\int_0^{1/2} \left( \int_x^{1-x} \left( \frac{x-y}{x+y} \right)^2 dy \right) dx$ .
3. By using the transformation  $x + y = u$  and  $y = uv$ , show that  $\int_0^1 \left( \int_0^{1-x} e^{y/(x+y)} dy \right) dx = \frac{e-1}{2}$ .  
[REMARK: Take great care when calculating the Jacobian.]

4. Suppose that  $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  and  $g : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  are differentiable. Suppose further that  $(u, v) = f(s, t)$  and  $(x, y) = g(u, v)$ , so that  $(x, y) = (g \circ f)(s, t)$ .  
a) Use the Chain rule  $(\mathbf{D}(g \circ f))(s, t) = (\mathbf{D}g)(u, v)(\mathbf{D}f)(s, t)$  to show that

$$\frac{\partial(x, y)}{\partial(s, t)} = \frac{\partial(x, y)}{\partial(u, v)} \frac{\partial(u, v)}{\partial(s, t)}.$$

- b) Show that  $\frac{\partial(x, y)}{\partial(u, v)} \frac{\partial(u, v)}{\partial(x, y)} = 1$  provided that  $\frac{\partial(x, y)}{\partial(u, v)} \neq 0$ .
5. Let  $S$  be the region in the first quadrant of  $\mathbb{R}^2$  bounded by  $x^2 - y^2 = 1$ ,  $x^2 - y^2 = 9$ ,  $xy = 2$  and  $xy = 4$ . Consider also the substitution  $u = x^2 - y^2$  and  $v = 2xy$ .  
a) Draw a picture of the region  $S$ .  
b) Use the result in Question 4(b) to find  $\partial(x, y)/\partial(u, v)$ .  
c) Show that  $2x^2 = (u^2 + v^2)^{1/2} + u$  and  $2y^2 = (u^2 + v^2)^{1/2} - u$ .  
d) Let  $S^* = \{(u, v) \in \mathbb{R}^2 : u \in [1, 9] \text{ and } v \in [4, 8]\}$ . Show that  $T : S^* \rightarrow S : (u, v) \mapsto (x, y)$  is one-to-one and onto.  
e) By changing variables, show that  $\iint_S (x^2 + y^2) dx dy = 8$ .

6. Evaluate each of the following triple integrals by using a suitable substitution:

- a)  $\iiint_S (x^2 + y^2 + z^2) dx dy dz$ , where  $S = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 \leq 4 \text{ and } 0 \leq z \leq 2\}$
- b)  $\iiint_S (x^2 + y^2 + z^2)^{1/2} e^{-(x^2+y^2+z^2)} dx dy dz$ , where  $S = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 + z^2 \leq 4\}$
- c)  $\iiint_S \frac{dx dy dz}{(x^2 + y^2 + z^2)^{3/2}}$ , where  $S = \{(x, y, z) \in \mathbb{R}^3 : a^2 \leq x^2 + y^2 + z^2 \leq b^2\}$  with  $b > a > 0$

7. Let  $S$  be the region in  $\mathbb{R}^3$  bounded by the paraboloid  $z = x^2 + y^2$  and the cone  $z = \sqrt{x^2 + y^2}$ .  
a) Find the set of points  $(x, y, z) \in \mathbb{R}^3$  where the paraboloid intersects the cone.  
b) Draw the cross section of the region  $S$  on the plane  $y = 0$ .  
c) Show that the volume of  $S$  is equal to

$$\iint_T (\sqrt{x^2 + y^2} - (x^2 + y^2)) dx dy,$$

where  $T = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 \leq 1\}$ .

- d) By changing to polar coordinates, show that the volume of  $S$  is equal to  $\pi/6$ .

8. By considering a suitable set of coordinates in  $\mathbb{R}^3$  and making the necessary change of variables, show that

$$\iiint_S \sqrt{x^2 + y^2 + z^2} \, dx \, dy \, dz = \frac{27\pi(2\sqrt{2} - 1)}{2},$$

where  $S$  is the region in  $\mathbb{R}^3$  bounded by the plane  $z = 3$  and the cone  $z = \sqrt{x^2 + y^2}$ .