

# Bajas Segregadas en el HS (Chile9

**René D. Garreaud**


Departament of Geophysics

Universidad de Chile

Santiago, Chile


# Outline

- Structure and evolution
- Climatological distribution
- Why COLs are so frequent/persistent off western South America
- The March 2015 Atacama rainstorm




## Cut off Lows...

- Horizontal scale of a few hundred km.
- Lifecycle of several days
- Erratic displacement, hard to predict.
- Can cause deep convection and intense precipitation (case study 2)
- Can also bring strong winds, heavy snowfall, and unusual cold conditions to high-elevation regions (e.g., Vuille and Ammann 1997)
- Increase stratosphere–troposphere exchange (STE) of trace gases






**2021-03-25**  
**20:40:17 UTC**










GFS 27 Mayo 2021 - 21  
UTC  
(Pronostico de 12 horas)


Contornos negros: PNM cada 10 mb  
Contornos celeste: Z500  
Colores: Agua Precipitable (rojo: 50 mm)




(a) Z500 (GFS) + GOES13 (Vis), 18 UTC, 09-03-2019




(b) Z500 + SLP (GFS), 18 UTC, 09-03-2019



(c) Temperatura máxima, 09-03-2019




(d) GOES16, Natural Color, 16 UTC, 09-03-2019






+ GOES13 (Vis), 18 UTC, 09-03-2019




(b) Z500 + SLP (GFS), 18 UTC, 09-03-2019






## BS: Un factor clave en las precipitaciones del norte de Chile



**Fig. 2.8** Distribution of annual precipitation due to cut-off lows: (a) spatial distribution and (c) latitudinal distribution from CR2MET daily rainfall product for the period 1979 to 2014 (Boisier et al. 2018), using the cut-off low database developed by Barahona (2016); (b) spatial distribution and (d) latitudinal distribution of the percentage of annual precipitation due to cut-off lows

La pregunta del millón...se segrega o no? Pasa o no pasa?


500-hPa geo. height (black, dam), temp. (red, C), ascent (blue,  $5 \times 10^{-3}$  hPa/s), cyc. rel. vort. ( $\times 10^{-5}$  s $^{-1}$ ),  
Initialized: 0600 UTC 27 May 2020 | Forecast hour: 126 | Valid: 1200 UTC 1 Jun 2020  
wind (barbs, kt)





# La pregunta del millón...se segrega o no? Pasa o no pasa?


500-hPa geo. height (black, dam), temp. (red, C), ascent (blue,  $5 \times 10^{-3}$  hPa/s), cyc. rel. vort. ( $\times 10^{-5}$  s $^{-1}$ ),  
Initialized: 0600 UTC 27 May 2020 | Forecast hour: 150 | Valid: 1200 UTC 2 Jun 2020  
wind (barbs, kt)






# La pregunta del millón...se segrega o no? Pasa o no pasa?

500-hPa geo. height (black, dam), temp. (red, C), ascent (blue,  $5 \times 10^{-3}$  hPa/s), cyc. rel. vort. ( $\times 10^{-5}$  s $^{-1}$ ),  
Initialized: 0600 UTC 27 May 2020 | Forecast hour: 174 | Valid: 1200 UTC 3 Jun 2020  
wind (barbs, kt)




# La pregunta del millón...se segrega o no? Pasa o no pasa?

500-hPa geo. height (black, dam), temp. (red, C), ascent (blue,  $5 \times 10^{-3}$  hPa/s), cyc. rel. vort. ( $\times 10^{-5}$  s $^{-1}$ ),  
Initialized: 0600 UTC 27 May 2020 | Forecast hour: 192 | Valid: 0600 UTC 4 Jun 2020  
wind (barbs, kt)









# COL structure and evolution

To illustrate the **structural evolution of a typical COL** in the SH we integrated MM5 (25 km resolution) for a 6 day period forced by NCEP-NCAR Reanalysis (2.5x2.5 lat-lon) in their lateral boundaries. Full topography and standard parameterizations.






Geopotential height  
and wind speed at  
300 hPa




Day of March 2005  
18:00 UTC



# COL structure and evolution

Upward - Downward







300 hPa geopotential height  
and vertical velocity





Day of March 2005  
18:00 UTC




Z(300 hPa) and  
mid level clouds


No cloud until  
reaching the  
continent!



00:00:00  
1999104  
1 of 97  
Wednesday



Vis5D



Vis5D


# COL structure and evolution

300 hPa geopotential height. Wind vectors and potential vorticity at 340 K


Weak CVA

Strong AVA


a. 1800 UTC 7 March 2005




b. 1800 UTC 8 March 2005




c. 1800 UTC 9 March 2005




d. 1800 UTC 10 March 2005










Vis5D

# Long-term mean distribution of COLs

Dataset: NCEP-NCAR Reanalysis (6 hourly, 2.5x2.5 lat-lon grids). 1979-2000, 2015  
Search and track(\*) closed lows at 300 hPa equatorward of the main westerly jet.  
Lows must satisfy criteria of intensity, duration (>1 day) and cold-core at upper levels.




(\*) Tracking algorithm following Murray and Simmonds, 1991



# Long-term mean distribution of COLs

Most COLS in three subtropical regions: Australia, South America and South Africa



Austral Summer  
(SA, SAF min)

Austral Winter  
(SA max)

Dir 1  
2

BS1  
X


X  
X  
BS2  
X






# Annual mean COL distribution

Some changes depending on level of identification and dataset



# Seasonal distribution of surface cyclones


In contrast to COL distribution, surface cyclone density maximizes in a circumpolar band at  $60^\circ$  with less asymmetry






# Annual mean COL distribution in the NH

DJF (NH Winter)






JJA (NH Summer)



500 hPa closed lows to the south of the Jet (from Bell and Bosart 1989)


# Others aspects of COLs in the SH




# Others aspects of COLs in the SH

For each COL we calculate the precipitation (TRMM) in a  $10^\circ \times 10^\circ$  lat-lon box around the COL center.

Area (size) and Intensity (colors) of mean precipitation for each COL



Percentage annual of precipitation accounted by COLs







# COL distribution in the SH

High frequency of COL off the west coast of South America (but in summer) due to:

Dry conditions over the SE Pacific → little diabatic heating → COLs tend to be last longer  
Dynamical forcing enhance their genesis (jet exit region, frequent blocking farther south, Andes cordillera)



Blocking anticyclone  
distribution





# Numerical experiments using WRF



12 UTC 10 March 2005

06 UTC 12 March 2005



*CTR*



*Red Topo  
(0.2\*H)*





*Dry  
(No SH)*



# Numerical experiments using WRF

400-hPa air temperature and cloud mixing ratio integrated between 600 and 300 hPa at 1800 UTC 11 Mar 2005 in CTR.

Green arrows are 36 hr back trajectories arriving at the 500 hPa level






**The March 2015 Atacama Storm.** Three days of intense rainfall triggered landslides and widespread flooding. More than 80 casualties and major damage to public and private infrastructure. Most acute impact during the event but many problems (e.g. public health) in subsequent months.




# Precipitation during the Event




# Distribución observada de la Precipitación diaria en estaciones DGA zona de Copiapo (1970-2013)

Precipitación media anual  $\approx 15$  mm






# Precipitation during the Event




It was a **warm storm**. Freezing level during the storm remained above 4000 m ASL...much of the Andean slope receive rain instead of snow

March 24, 2015

Lightning detected by WWLL  
One every 10 strokes




# Synoptic environment I: Z500 and SLP






# Large scale context

OLR & H250 19-22 Mar 2015




COL developed off northern Chile in connection with a large-scale ridge over the South Pacific that in turn growth from energy propagating from the tropics by the PSA mode....

# Synoptic environment I: PW and 850 winds




# What caused such a intense storm over Atacama?

Dynamic features




Thermodynamic features






Plausible suspect: marked, sudden SST warming off South America (EN 2015)  
Destabilize the atmosphere and provide extra moisture

Niño 1-2  
SST




SST anomaly  
23 March 2015




# Numerical experiment using RegCM (forced by ERA)

In a sensitivity run the SST was kept equal to the field at March 10 (prior to the warming)  
thus causing a sfc BC cooler than the control run




## CTR PW (contours) and SENS-CTR PW (colors)






# RegCM simulated precipitation


CTR




Sens



Sens-CTR



Sens/CTR %



# Conclusions

- \*COLs in the SH have similar structure and characteristics than their NH counterparts

- \*COLs in the SH tends to cluster in three subtropical areas: Australia, South America and Africa, away from baroclinically active regions





- \*The Andes cordillera has little influence on the COL formation and intensification. Rather, the cyclone segregation appears mostly driven by the large-scale, upper-level circulation.

- \*The Andes delays the COL demise by blocking the inflow of warm, moist air from the interior of the continent that would otherwise initiate deep convection.

- \*Given enough moisture, COLs can cause heavy precipitation even in the driest place on earth (Atacama dessert)

# Precipitación acumulada en 6 horas


## Pronóstico valido a las 18 hrs, 24 Mar 2015






# Annual mean COL distribution in the NH

DJF (NH Winter)



JJA (NH Summer)



Light blue shading: 500 hPa closed lows to the south of the Jet (from Bell and Bosart 1989)  
Warm colors: Surface cyclones density (from Raible et al. 2008)