

Psychosocial Factors in Medication Adherence and Diabetes Self-Management: Implications for Research and Practice

Jeffrey S. Gonzalez

Yeshiva University and Albert Einstein College of Medicine

Molly L. Tanenbaum

Stanford University

Persis V. Commissariat

Joslin Diabetes Center, Boston, Massachusetts

Diabetes is a chronic illness that places a significant self-management burden on affected individuals and families. Given the importance of health behaviors—such as medication adherence, diet, physical activity, blood glucose self-monitoring—in achieving optimal glycemic control in diabetes, interventions designed and delivered by psychologists hold promise in assisting children, adolescents, and adults with diabetes in improving their health status and lowering their risk of serious complications. This article first provides an overview of diabetes self-management and associated challenges and burdens. Socioeconomic status factors that may influence diabetes management and outcomes are briefly highlighted. We then review the evidence base for select psychosocial factors that may be implicated in diabetes self-management. Modifiable targets of psychological intervention are presented across 3 overarching domains: (a) knowledge, beliefs, and related cognitive constructs; (b) emotional distress and well-being; and (c) behavioral skills and coping. Important methodological issues facing future research are discussed, along with opportunities for psychologists in improving the care and treatment outcomes of individuals and families living with diabetes. In conclusion, we advocate for continued research emphasis on improving psychosocial aspects of living with diabetes, with greater attention to the situational context in which the self-regulatory processes underlying self-management occur. Psychologists have important roles to play in reducing emotional distress, improving patient knowledge, and providing training in behavioral skills to promote successful self-management and to support patient-centered diabetes care.

Keywords: diabetes, adherence, self-management, glycemic control, psychosocial factors

As the prevalence of Type 1 (T1D) and Type 2 diabetes (T2D) continues to rise in the United States and worldwide, more individuals and families are living with the challenge of integrating into their lives a demanding, complex, and lifelong

regimen to control their progressive illness and prevent or delay diabetes complications. Despite the availability of effective treatments and landmark trials establishing the benefits of tight glycemic control (see Hunter, 2016), many struggle with diabetes self-management and fail to achieve levels of risk-marker control that would significantly reduce the likelihood of serious complications and early mortality. In this review, from our vantage point as clinical health psychologists, we (a) provide a brief snapshot of the problem of diabetes self-management, (b) touch on relevant socioeconomic and socio-cultural contexts, and (c) review recent literature on psychosocial factors associated with diabetes self-management, with an emphasis on modifiable targets of psychological intervention. The final section highlights methodological issues for future research and identifies important roles for psychologists in improving the health of those living with diabetes.

Editor's note. This article is one in a collection of articles published in a special issue of *American Psychologist* titled "Diabetes and Psychology" (October 2016). Deborah J. Wiebe, Vicki S. Helgeson, and Christine M. Hunter provided scholarly lead for the special issue.

Authors' note. Jeffrey S. Gonzalez, Ferkauf Graduate School of Psychology, Yeshiva University, and Diabetes Research Center, Albert Einstein College of Medicine; Molly L. Tanenbaum, Department of Pediatrics, Stanford University School of Medicine; Persis V. Commissariat, Joslin Diabetes Center, Boston, Massachusetts.

Jeffrey S. Gonzalez's efforts on this article were partially supported by grants from the National Institute of Diabetes and Digestive and Kidney Diseases (R18 DK098742 and R01 DK104845). We thank Naomi Kane, Deborah Binko, Lauren Klayman, Joslyn Kenowitz, Gabrielle Frackman, and Amit Shapira for their contributions to the literature search for this review.

Correspondence concerning this article should be addressed to Jeffrey S. Gonzalez, Ferkauf Graduate School of Psychology, Yeshiva University, 1300 Morris Park Avenue, Bronx, NY 10461. E-mail: jeffrey.gonzalez@einstein.yu.edu

The Challenge of Diabetes Self-Management

Many individuals living with diabetes have difficulty achieving recommended standards for diabetes management. Long-term medication use and lifestyle change are necessary for the successful management of both T1D and



**Jeffrey S.
Gonzalez**

T2D. In insulin-dependent T1D, exogenous insulin self-administration, via multiple daily injections or an insulin pump, is the focus of treatment. Individuals with T2D are increasingly treated with multiple-medication regimens, including oral medications and/or exogenous insulin. Weight loss and control of cardiovascular risk factors, such as blood pressure and cholesterol levels are often additional areas of focus in T2D management (see Hunter, 2016, for more details). Although self-management demands differ significantly, problems meeting treatment goals in real-world practice are common for both groups of patients. We review literature on T1D and T2D here, but note that samples of “adults with diabetes” often include far more individuals with T2D because of differences in prevalence.

About half of U.S. adults diagnosed with diabetes achieve targets for glycemic control, and similar proportions meet recommended blood pressure and cholesterol targets. Less than one in five adults reach recommended targets for all three risk biomarkers, the foundational “ABCs” of diabetes care (Stark Casagrande, Fradkin, Saydah, Rust, & Cowie, 2013). Although less stringent targets are recommended for youth with T1D, most still do not reach them. A registry-based study showed that 64% of children under 6 years of age, 43% of those 6 to 12, and only 21% of those 13 to 19, reached targets for glycemic control (Wood et al., 2013). This suggests that many patients will not reap the full health benefits of treatment and points to suboptimal diabetes self-management as a target for behavioral interventions that can improve treatment outcomes.

The diverse behavioral requirements for diabetes self-management—including regular medication taking, self-monitoring of blood glucose (SMBG), changes in diet and

physical activity, foot self-care, and visits with health care providers—are detailed elsewhere in this issue (Hunter, 2016). Here, we use medication *adherence* to refer to the extent to which patients follow medication prescriptions from a health care provider. Measurement of adherence may involve comparing, for example, actual medication-taking frequency with prescribed dosing frequency over a defined period. *Self-management* refers to regular performance of the broader set of diabetes self-care behaviors, including medication adherence. Successful self-management thus requires clear communication about the standards for self-care from providers and patient agreement with these goals. Nonadherence to recommendations can be unintentional or purposeful, reflecting patient decision making that weighs anticipated treatment-related benefits and costs to health and quality of life (DiMatteo, 2004). Suboptimal diabetes self-management is clearly associated with worse glycemic control (e.g., Feldman et al., 2014; Hood, Peterson, Rohan, & Drotar, 2009; Schectman, Nadkarni, & Voss, 2002) and increased risk for hospitalization, complications, and mortality (e.g., Cho et al., 2011; Currie et al., 2012; Ho et al., 2006; Kuo et al., 2003). Thus, the consequences of poor diabetes self-management can be severe.

Available evidence suggests that suboptimal diabetes self-management is common, especially for regimen aspects involving lifestyle change. A large multinational survey of adults with diabetes found that successful self-management rates were often suboptimal—best for medication and lowest for physical activity, SMBG, and regular self-examination of the feet (Nicolucci et al., 2013). This is consistent with evidence across chronic illnesses. A meta-analysis of 569 studies placed diabetes among the illnesses with the lowest levels of adequate self-management, likely reflecting the greater focus on lifestyle changes in diabetes care, relative to many other chronic illnesses (DiMatteo, 2004).

Lower rates of successful diabetes self-management also partially reflect the demands of a burdensome treatment regimen. Living with diabetes is best conceptualized as a chronic stressor for patients and families, affecting various life domains (Nicolucci et al., 2013). Self-management is time consuming: Implementation of recommended behaviors has been estimated to take up 2 hr per day for an average adult with T2D (Russell, Suh, & Safford, 2005). These underappreciated time costs are accompanied by direct financial costs related to health care visits, medications, and supplies. Medication regimens are often complex and cause side effects. Intensive treatment regimens raise the risk of hypoglycemia (Yudkin, Richter, & Gale, 2010), which, when severe, can result in physical injury, car accidents, and death. Particular aspects of self-management, such as insulin injections for adults with T2D, are often appraised negatively by patients (Rubin, Peyrot, Kruger, & Travis, 2009) and



**Molly L.
Tanenbaum**

are associated with increased patient distress (Baek, Tanenbaum, & Gonzalez, 2014). Thus, the demands and burdens of diabetes treatment provide an important situational context for any understanding of person-level psychosocial factors associated with suboptimal diabetes self-management.

Psychosocial Factors Related to Diabetes Self-Management

A large body of research has sought to identify psychosocial factors that could be targeted by interventions to improve diabetes self-management and treatment outcomes. In the following sections, we briefly highlight the role of social disadvantage in diabetes self-management before reviewing relevant patient-level modifiable factors across three broad domains: (a) knowledge, beliefs, and related cognitive constructs; (b) emotional distress and well-being; and (c) behavioral skills and coping. Developmental, family context, and social relationship factors are covered elsewhere in this issue (see Hilliard, Powell, & Anderson, 2016; Wiebe, Helgeson, & Berg, 2016). We also limit our examples of successful interventions to those tested in adults and refer readers to a comprehensive review of pediatric interventions in this special issue (see Hilliard et al., 2016).

The Socioeconomic and Cultural Context of Diabetes Self-Management

Evidence clearly shows that socially disadvantaged individuals with diabetes benefit the least from available treatments, both in terms of glycemic control and risk for com-

plications (Brown et al., 2004; Naranjo, Hessler, Deol, & Chesla, 2012). Costs are a common patient-reported barrier to self-management (e.g., Daly et al., 2009) and have been shown to reduce adherence (Roblin et al., 2005) and preventive care (Karter et al., 2003). For example, socially disadvantaged children with T1D are far less likely to be on insulin pump therapy (Lin et al., 2013), an alternative to multiple daily self-injections associated with improved control and less hypoglycemia (e.g., S. R. Johnson, Cooper, Jones, & Davis, 2013). A survey of 14,357 adults with T2D found that 16% of low-income adults experienced severe hypoglycemia in the prior year, compared with 8.8% of high-income adults (Berkowitz et al., 2014). Similar differences were found between those with low (11.9%) and high (8.9%) levels of education (Berkowitz et al., 2014). Thus, socioeconomic determinants of health outside the scope of medical practice have robust effects on diabetes treatment outcomes.

Ethnic minorities achieve relatively worse diabetes treatment outcomes, experience more emotional distress, and report lower quality of life than Whites (Naranjo et al., 2012). However, although some studies report ethnic differences in self-management, population-based survey data suggest few differences; when they exist, they sometimes suggest better self-management among ethnic minorities (P. J. Johnson, Ghildayal, Rockwood, & Everson-Rose, 2014). Data suggest that racial and ethnic minorities and those with lower incomes and less education spend more time on diabetes self-management than less disadvantaged groups (Ettner et al., 2009). Other data suggest that disparities may be most robust for SMBG and medication adherence, for which socioeconomic status (SES) likely plays an important role. For example, ethnic minorities, those with lower SES, those with difficulty communicating in English, and those with higher costs for test strips were significantly less likely to adequately perform SMBG (Karter, Ferrara, Darbinian, Ackerson, & Selby, 2000). Black adults with T2D were just as likely as Whites to fill their medication prescription, but almost twice as likely to discontinue over 10 years (Trinacty et al., 2009). SES often overlaps with ethnic minority status and may partially explain ethnic disparities. For example, epidemiological data suggest that about half of Latinos with diabetes do not have health insurance and link insurance status and SES to diabetes prevalence (Schneiderman et al., 2014).

Cultural beliefs and norms may also play a role in ethnic disparities. Ethnic minorities with T2D may be more likely to worry about drug side effects and medication dependency; they may also be more reluctant to accept treatment intensification when clinically indicated (Huang et al., 2009). Qualitative data suggest that low-income Blacks and Latinos with T2D can have doubts about the necessity of medication and may take less than prescribed to guard against negative effects (Lynch, Fernandez, Lighthouse,



**Persis V.
Commissariat**

Mendenhall, & Jacobs, 2012). A number of small studies suggest that Latinos with T2D can have particularly negative views about insulin therapy, including fears that insulin causes blindness (Gutiérrez, Ferro, & Caballero, 2015). Although cultural factors may influence diabetes self-management and treatment outcomes among ethnic minorities (Naranjo et al., 2012), research suggests that the widest cultural gap is between patients and their diabetes care providers. Patients often endorse explanatory models for diabetes and its treatment that differ significantly from those underlying their physicians' approach to care—differences that persist even when national culture and language are shared (Weller, Baer, Garcia de Alba Garcia, & Salcedo Rocha, 2012). Thus, SES and sociocultural factors should provide important context for the following patient-level constructs.

Patient Knowledge, Beliefs, and Related Cognitive Factors

Knowledge, literacy, and numeracy. Successful diabetes self-management requires considerable knowledge on the part of patients and families, including understanding of the effects of diabetes on the body, the goals of treatment, and the effects of various behaviors on glucose regulation. General intelligence consistently and robustly predicts a variety of health outcomes, seemingly independent of SES, and has been conceptualized as an important causal factor that influences health indirectly, through better health behavior and disease self-management (Gottfredson & Deary, 2004). In one study, social class and reading scores, an indicator of intelligence, from recently diagnosed adults

with T1D independently predicted diabetes knowledge over time (Taylor, Frier, Gold, Deary, & Edinburgh Prospective Diabetes Study, 2003). Although it is difficult to separate intelligence effects from crudely measured indicators of SES, it seems quite plausible that one's general ability to learn, reason, and solve problems would play a role in diabetes self-management. Specific aspects of neurocognitive functioning have been associated with better glycemic control, although at least part of this association likely reflects neurocognitive deficits resulting from poor control (see Ryan, van Duinkerken, & Rosano, 2016).

Health literacy skills, including reading comprehension, writing, communicating, and making decisions about health care, are often low among individuals treated for diabetes and could interfere with successful self-management. A review of 23 studies investigating the relation between health literacy and treatment adherence across chronic illnesses found mixed results: Only five studies found clear evidence of a relation with adherence (Ostini & Kairuz, 2014). A diabetes-specific review, focused primarily on adults with T1D and T2D, came to similar conclusions: Although there is strong, consistent evidence that health literacy is related to diabetes knowledge, associations with self-management and treatment outcomes are inconsistent (Al Sayah, Majumdar, Williams, Robertson, & Johnson, 2013). In youth with T1D, higher caregiver literacy, but not child literacy, has been linked to improved glycemic control (Pulgarón et al., 2014; Sanders, Federico, Klass, Abrams, & Dreyer, 2009).

Numeracy skills include the ability to interpret and respond to numerical feedback (e.g., count carbohydrates in food, recognize out-of-range blood glucose numbers, understand medication dosages). Proficiency in these skills has been weakly related to glycemic control in adults with T1D or T2D (Cavanaugh et al., 2008). In one study of adolescents with T1D, parents' numeracy skills were not related to adolescent self-management (Janisse, Naar-King, & Ellis, 2010); another study found a relationship between parental numeracy and glycemic control (Pulgarón et al., 2014). Results from two randomized trials aiming to improve numeracy and literacy in adults demonstrated initial glycemic improvement that diminished by 6 months, with no observed group differences in diabetes self-management (Cavanaugh et al., 2009). Taken together, findings on literacy and numeracy skills have not shown clear, consistent associations with diabetes self-management and control. Relationships may depend on other factors that vary across available studies.

National survey data also highlight basic knowledge gaps among many adults living with diabetes—less than 50% know their level of glycemic control, 63% know their blood pressure, and only 22% know their cholesterol level; this basic knowledge is even less common among Mexican Americans and those with lower SES

(Stark Casagrande et al., 2012). Guidelines emphasize that all patients should receive formal diabetes self-management education and ongoing support (American Diabetes Association, 2015), but *less than 7%* of privately insured adults receive self-management education and training within the first year after diagnosis (Li et al., 2014). Although these data highlight the need for better education, knowledge appears to be necessary, but not sufficient, for successful diabetes self-management. A review of 72 self-management trials concluded that didactic interventions had positive effects on knowledge but were inconsistent on other outcomes; the authors concluded that factors other than knowledge are needed to achieve long-term change in self-management (Norris, Engelgau, & Narayan, 2001). Similarly, a review of 24 interventions aimed at improving self-management and glycemic control in youth with T1D found little evidence to support the efficacy of educational programs (Murphy, Rayman, & Skinner, 2006). The importance of behavioral interventions that go beyond education to incorporate training in behavioral skills and strategies, with opportunities for practice, feedback, reinforcement, and support, has long been recognized (e.g., Rubin, Peyrot, & Saudek, 1993). Effective T2D interventions tend to be longer, be delivered by teams of providers, and reinforce information over time (Loveman, Frampton, & Clegg, 2008). Characteristics of successful educational youth interventions include parental involvement, self-efficacy promotion, and integration into routine clinical care (Murphy et al., 2006).

Beliefs about illness and treatment. Beyond knowledge, patient beliefs about illness and treatment have been a focus of various theoretical models for understanding health behavior. A recent review of health beliefs in diabetes concluded that Leventhal's common sense model (CSM) of self-regulation (Leventhal, Diefenbach, & Leventhal, 1992) has a unique advantage among various health belief theoretical models, in that it considers both cognitive and emotional processes involved in illness self-management (Harvey & Lawson, 2009). The CSM is a self-regulatory model that views patients as agents acting in a sociocultural context, their efforts guided by beliefs about illness—its *identity* (symptoms and label), perceived *cause*, expected *time-line*, *consequences*, and anticipated *controllability*—and further shaped by appraised feedback about the effects of their coping behaviors. Subsequent work has focused on beliefs about medication; a meta-analysis of 94 of these studies in various illnesses found consistent evidence linking better adherence to greater perceptions of necessity of treatment and fewer concerns about adverse effects (Horne et al., 2013). Adults with diabetes who are younger, African American, or who have low health literacy may have more concerns about their medications

(Aikens & Piette, 2009), perhaps contributing to medication nonadherence over time. Impoverished Latinos and African Americans may have commonsense, but inaccurate, beliefs, such as believing that medications only need to be taken when blood glucose is high—and these beliefs are associated with nonadherence (Mann, Ponieman, Leventhal, & Halm, 2009).

CSM beliefs about diabetes are consistently associated with self-management, both concurrently and prospectively, in adults with T1D or T2D and adolescents with T1D (e.g., Hampson, Glasgow, & Foster, 1995; Nouwen, Urquhart Law, Hussain, McGovern, & Napier, 2009; T. C. Skinner & Hampson, 2001; Watkins et al., 2000). A meta-analysis showed that stronger identity (more symptoms and disease burden), perceptions of greater consequences of diabetes, lower personal control, a more unpredictable course, as well as more diabetes-related emotional distress, were consistently associated with poorer glycemic control in adults with T1D or T2D, with results suggesting 2% to 7% shared variance (Mc Sharry, Moss-Morris, & Kendrick, 2011). Whether these associations indicate that inaccurate beliefs cause poorer self-management, or whether they reflect interpretations of the often confusing lived experience of poorly controlled diabetes, is difficult to ascertain from existing studies.

Evidence from CSM-informed interventions reviewed by Mc Sharry and colleagues (2011) is limited but partially supportive. Of four trials that targeted glycemic control and measured CSM beliefs, two did not achieve changes in beliefs (French et al., 2008; George et al., 2008). Favorable changes, including better understanding of and greater perceived control over diabetes, were found in two trials that specifically targeted CSM beliefs (Davies et al., 2008; Keogh et al., 2011). One of these trials demonstrated an impact on glycemic control (Keogh et al., 2011), and the other on smoking, weight, and depression (Davies et al., 2008). These outcomes are promising, as is the pattern suggesting that studies that were most focused on changing CSM beliefs had the best effects. At its most basic level, this evidence suggests that intuitive, but often inaccurate, beliefs about diabetes and its treatment are common, and are likely implicated in problematic diabetes self-management and malleable to intervention.

Self-efficacy and perceived control. Patients may be fully confident in their ability to implement their diabetes self-management regimen (high self-efficacy), but if they do not believe that these efforts will affect outcomes of interest (low perceived control), they may be less motivated or consistent in their implementation. According to social-cognitive theory, each of these beliefs influences task performance (Bandura & Wood, 1989), and a large literature has developed to study them as well as a variety of related constructs (E. A. Skinner, 1996). We focus on self-efficacy

and perceived control as the most extensively researched constructs in diabetes.

Observational studies of T1D and T2D in adults and adolescents have demonstrated associations between self-efficacy for diabetes self-management and some aspects of diabetes self-management (e.g., King et al., 2010; Nouwen et al., 2009; Sarkar, Fisher, & Schillinger, 2006). Beyond the CSM studies reviewed in the previous section others also find that greater perceived control over diabetes is associated with better self-management in adults with T2D (e.g., Macrodimitris & Endler, 2001; Surgenor, Horn, Hudson, Lunt, & Tennent, 2000; Walker et al., 2012). In adolescents with T1D, self-efficacy and perceived control were consistent correlates of all aspects of diabetes self-management, as well as glycemic control; only perceived control was an independent predictor of diabetes self-management (Griva, Myers, & Newman, 2000). Among distressed adults with T2D, both perceived control and self-efficacy were significantly associated with better adherence, but only perceived control was independently significant. Furthermore, perceived control was independently related to glycemic control and accounted for a substantial portion of the variance shared by emotional distress and these outcomes (Gonzalez, Shreck, Psaros, & Safren, 2015). Perceived competence for diabetes tasks, a construct overlapping with self-efficacy and perceived control, also mediated the relationship between state affect and glycemic control in a study of adolescents with T1D (Fortenberry et al., 2009).

Along with preliminary evidence for intervention effects on perceived control, discussed in the previous section, there is strong evidence that self-efficacy can be increased through a variety of interventions. For example, a large trial evaluated the impact of a telemedicine approach to diabetes education, support, and collaborative goal setting on glycemic control in 1,665 Medicare beneficiaries with T2D. Results showed the intervention improved glycemic control and self-efficacy; changes in self-efficacy and glycemic control were correlated; and self-efficacy was a partial mediator of intervention effects on glycemic control (Trief, Teresi, Eimicke, Shea, & Weinstock, 2009). Influential work has demonstrated enduring effects of a chronic disease self-management program on self-efficacy and health outcomes in various chronic illnesses, including adults with T2D (e.g., Lorig, Ritter, Villa, & Piette, 2008). Consistent with the CSM, these findings suggest that successful diabetes self-management requires confidence in ability, as well as evidence of impact: Patients must be able to observe changes in relevant outcomes (feedback). These beliefs may be fruitful targets for psychological intervention that is sensitive to the context of diabetes, as well as the broader social context of structural barriers that can often pose significant challenges to sustaining these beliefs over time.

Emotional States and Distress

A large literature has explored emotions, mental health, and psychiatric illness in relation to risk for the development of diabetes and treatment outcomes among those with diabetes (see deGroot, Golden, & Wagner, 2016). Here, we focus on the consistent link between depressive symptoms and poor diabetes self-management, which is significant in adults and children, and more robust when stronger measurement methods are used (Gonzalez et al., 2008). This association does not appear to be primarily explained by effects of comorbid psychopathology. Instead, emotional distress, which overlaps substantially with items on depression screening scales, appears to underlie these relationships. Diagnostic interviews for depression are rarely used, but when they are, they show little evidence for a relation between major depressive disorder (MDD) and diabetes self-management or glycemic control (Fisher, Gonzalez, & Polonsky, 2014; Gonzalez, Fisher, & Polonsky, 2011).

In contrast, elevations in depressive symptoms, even at subclinical levels that would not meet the diagnostic threshold for MDD, and distress related to the burdens of living with diabetes, or diabetes-related distress (Polonsky et al., 2005), are consistently associated with poorer self-management (Fisher et al., 2014). Diabetes-related distress generally shows a closer association to glycemic control than depression (Fisher et al., 2014; Gonzalez et al., 2015; Schmitt et al., 2015), and appears to be more common and chronic than depression in adults with T2D (Fisher et al., 2010). International survey data support this and suggest that emotional well-being is the domain of functioning most negatively affected by diabetes, second only to physical health (Nicolucci et al., 2013). Diabetes also provides a situational context for several related emotional distress constructs implicated in self-management. For example, fear of hypoglycemia, measured via validated self-report, is common among parents, children, and adults with T1D, and can be associated with worse glycemic control (Wild et al., 2007). Worries and concerns about hypoglycemia may also be important among adults with T2D (Polonsky, Fisher, Hessler, & Edelman, 2015). Parents often experience significant emotional distress related to their child's diabetes management (Streisand, Braniecki, Tercyak, & Kazak, 2001), and the perceived burden of treatment may explain the association between parent psychological distress and child glycemic control (Cunningham, Vesco, Dolan, & Hood, 2011).

Although much of the literature on emotions in diabetes has focused on negative affective states and/or emotionally distressing problems, some studies have examined the role of positive emotions and resilience factors, including optimism, self-esteem, subjective well-being, and sense of purpose (Celano, Beale, Moore, Wexler, & Huffman, 2013). However, evidence is generally not supportive of inten-

tion or behavior effects for positive affect manipulations (Cameron, Bertenshaw, & Sheeran, 2014). Similarly, although intervention studies for depression and distress have been mostly successful in affecting these negative affective targets, evidence does not support the hypothesis that their amelioration would, on its own, result in better diabetes self-management (Fisher et al., 2014; Markowitz, Gonzalez, Wilkinson, & Safren, 2011). Thus, rather than directly influencing diabetes outcomes, emotions likely play indirect causal roles, interact with other factors, or are outcomes of advancing illness and/or problematic diabetes self-management, rather than a cause.

Behavioral Skills, Coping, Self-Control, and Self-Regulation

Problem solving and coping. Cognitive and behavioral skills—involving the identification of problems, generating potential strategies for their resolution, selecting the most appropriate strategy, implementing it, and evaluating its effectiveness—represent a foundational component of diabetes self-management, as recognized by the American Association of Diabetes Educators (2015). Given the dynamic nature of diabetes progression and treatment over time, problem-solving skills are largely developed in the course of day-to-day experience, outside of medical encounters. Although studies suggest that problem-solving skill is associated with better self-management in adults with T2D (e.g., King et al., 2010) and adolescents with T1D (e.g., Mulvaney et al., 2014), a review of interventions targeting problem-solving skill in diabetes indicates considerable heterogeneity in content, and weak and inconsistent effects on self-management for adults and adolescents; few trials measure problem solving as an outcome or target these skills with sufficient potency (Fitzpatrick, Schumann, & Hill-Briggs, 2013).

Problem-solving skills share some overlap with problem-focused and approach-oriented coping strategies, which have been linked to better emotional adjustment and glycemic control in adults with diabetes (Duangdao & Roesch, 2008). Coping research in youth with chronic illness, including T1D, emphasizes the central role of perceived control over stressors. Primary control strategies, aimed at changing one's situation, may be adaptive for certain aspects of self-management, but may be less well matched to other stressful aspects of living with T1D, such as feeling different from peers or fear of complications. Secondary control coping, involving efforts to adapt to stress through reappraisal, positive thinking, acceptance, and distraction, is associated with better overall adjustment (Compas, Jaser, Dunn, & Rodriguez, 2012). Various effective interventions targeting coping in youth are reviewed in this issue (Hilliard et al., 2016).

Self-control and self-regulation. Research on both adolescents and adults with diabetes has highlighted the importance of self-control in diabetes self-management. Self-control involves one's ability to regulate emotions, behaviors, and impulses, as well as ability to delay gratification. One early and influential study of adults with T1D and T2D suggested that higher levels of self-control were associated with better glycemic control, an effect partly accounted for by better self-management (Peyrot, McMurry, & Kruger, 1999). In a study of adolescents with T1D, impulse control was significantly associated with better self-management, partially mediated by self-efficacy (Stupiansky, Hanna, Slaven, Weaver, & Fortenberry, 2013). Another study of adolescents with T1D showed that those with low self-control and infrequent use of emotion regulation strategies had the worst glycemic control (Hughes, Berg, & Wiebe, 2012). A further study suggests that self-control may mediate the relation between intelligence and better glycemic control in adolescents with T1D (Berg et al., 2014). In a set of preliminary studies, higher levels of self-control, measured by an objective handgrip test, were positively associated with weight loss, program attendance, and behavior change during a behavioral weight loss intervention for obese adults (Leahey, Xu, Unick, & Wing, 2014).

Self-regulatory models of health behavior encompass problem-solving, coping and self-control, but focus on the *process* of executive management of cognitions, emotions, and behavior toward the attainment and maintenance of personal and interpersonal goals (Lansing & Berg, 2014; Maes & Karoly, 2005). Interventions that incorporate strategies aimed at improving self-regulation hold promise for influencing health behaviors relevant to diabetes self-management. For example, one study compared a single-session intervention of information plus self-regulation strategies—including goal identification, mental contrasting, and recording of implementation intentions—against an information-only condition for effects on fruit and vegetable consumption in adult women without diabetes. Although both groups improved, by 4 months post-intervention, the self-regulation group recorded significantly more fruits and vegetable consumption, a difference that persisted over 2 years of follow-up (Stadler, Oettingen, & Gollwitzer, 2010). Another intervention, based on self-monitoring, positive reinforcement, skills training and support, was successful at maintaining weight loss over 18 months. Self-weighting increased with intervention and was strongly associated with weight loss maintenance, emphasizing the importance of feedback in self-regulation (Wing, Tate, Gorin, Raynor, & Fava, 2006).

Study of the self-regulatory processes involved in diabetes self-management, as they occur over time, should be a priority for future behavioral diabetes research. Problem-solving, coping, and self-control in the service of diabetes

self-management should be reinforced by monitored progress toward goals, or feedback (Harkin et al., 2016). However, diabetes provides multiple sources of sometimes competing feedback (e.g., emotional distress and side effects vs. SMBG and glycemic control), which may complicate evaluations of goal progress and effects of self-management efforts (Tanenbaum et al., 2015). A process-oriented approach that views patients as actively engaged in a challenging and dynamic self-regulatory task provides a better fit with the realities of diabetes self-management than concepts such as “compliance” or “adherence.”

Conclusions and Implications for Science and Practice

Improving diabetes self-management among the growing number of children, adolescents, and adults living with T1D and T2D would have a significant impact on treatment outcomes and public health. Psychologists should have a key role in meeting this challenge. Psychological factors related to patient knowledge and health beliefs, emotional states, and use of problem-solving and self-regulatory skills—core constructs relevant to psychological research and practice—have each been linked to diabetes self-management behavior. Improving our understanding of diabetes self-management could strengthen the broad evidence base for psychology.

It is useful to place the current brief review of such a large literature in the context of prior reviews. One of the earliest we could find focused on pediatric diabetes and was written by a pioneering health psychologist and future leader of the American Psychological Association (S. B. Johnson, 1980). Three years later, a strong argument was made in *American Psychologist* for the study of behavioral aspects of diabetes as an important paradigm for the then-emerging field of health psychology (Surwit, Feinglos, & Scovern, 1983). The contemporary literature had been dominated by the psychosomatic perspective, which sought to identify psychogenic factors—primarily personality traits, family characteristics, and life stress—that could influence diabetes onset and course. Both reviews were equally pessimistic about this approach, based on already available evidence. Studies focused on psychogenic factors were unable to rule out the influence of diabetes and its treatment on their psychosocial measures. They also tended to focus on only one or two self-reported factors, measured on a single occasion, and to assess general constructs rather than those specific to diabetes (S. B. Johnson, 1980). There has been much accomplished in the study of the psychosocial experience of diabetes patients since this time. However, we close by echoing S. B. Johnson’s (1980) still relevant “plea” for better measurement, as “first things are [still] first” (p. 113).

One long noted measurement issue relates to studies that treat glycemic control as a proxy for self-management. This is problematic because evidence suggests that measures of self-management share only 2.5% to 10% variance with concurrently measured glycemic control (Feldman et al., 2014; Hood et al., 2009; Schectman et al., 2002), and likely much less (e.g., 1.5% to 5%) with changes in control over time (Houle et al., 2015; Schectman et al., 2002). Self-management is one of many factors influencing glycemic control. Biological factors are also implicated, as are health care factors, such as access, cost, provider expertise, limited time for visits, increasing complexity of medication regimens, limited use of SMBG data by physicians, and failure to intensify treatment when indicated (Vigersky, 2011). Thus, making inferences about self-management from glycemic control, in research and practice, will mostly involve error. Behavior is the primary outcome of behavioral interventions and should be the focus of behavioral science in diabetes. Furthermore, the value of sustained improvement in self-management goes beyond what is captured by glycemic control at a single time point.

Better measurement of diabetes self-management is needed to advance the field. Accurate measurement can be difficult given the diversity of behaviors involved and their modest intercorrelation (e.g., Toobert, Hampson, & Glasgow, 2000). Furthermore, aspects of self-management not directly related to glycemic control are too often neglected. Foot self-care is particularly understudied considering the health and financial costs of foot complications in diabetes (Armstrong et al., 2013; Vileikyte & Gonzalez, 2014). Cardiovascular disease is the most significant health threat to adults with T2D, and it is clear that efforts aimed at controlling blood pressure and cholesterol will have much larger effects on health outcomes than those focused only on glycemic control (Yudkin, Richter, & Gale, 2010). Technological advances provide a growing set of options for measuring self-management, particularly pill taking, SMBG, insulin pump use, and physical activity (Driscoll & Young-Hyman, 2014). As measurement error is relevant to all available methods, latent variable analytic models that pool shared variance across multiple methods and time points of assessment represent a promising approach (Gonzalez & Schneider, 2011).

Greater attention to the situational context in which self-management occurs is also necessary. Even objective measures of self-management will be vulnerable to the “healthy adherer effect”—as they are often confounded by other health behaviors, SES, health care, and social factors. For example, meta-analysis shows that adherence to placebo predicts reduced mortality, even when measured objectively, with an effect similar in magnitude to that of adherence to beneficial drug therapy (Simpson et al., 2006). Unraveling these “healthy adherer” effects from the causal effects of self-management should be prioritized. Covariate

adjustment for crude indicators (e.g., years since diagnosis, counts of comorbidities, years of education, income) of contextual factors is likely to be insufficient. Despite these context-related measurement challenges, aspects of the work of diabetes self-management (e.g., regular pill taking, SMBG, appointment keeping) can be more accurately and easily quantified than many other behavioral outcomes of interest to psychologists. Thus, diabetes self-management provides an important paradigm for a psychological approach to health behavior change and maintenance.

Health psychology has played a major role in increasing the profile of psychological research in the medical and epidemiological literatures, and has influenced practice standards regarding psychosocial aspects of diabetes care. However, the ghost of the psychogenic approach, focused on the identification of causal effects for single, static characteristics, such as “diabetic personalities” and “psychosomatic families,” on diabetic outcomes, which earlier reviewers all but declared dead more than 30 years ago (S. B. Johnson, 1980; Surwit et al., 1983), continues to haunt many studies that overemphasize blood test results for glycemic control over behavioral outcomes. This narrow focus on glycemic control as a key outcome of diabetes is driven by important evidence on the population-level benefits of intensive treatment as well as funding priorities. But it should be recognized that glycemic control, although an important predictor of diabetes complications, is a surrogate outcome that has different meanings for risk of complications depending on patient age and other factors. When outcomes of intensive treatment are considered in terms of risk for macrovascular (e.g., myocardial infarction and stroke) and microvascular (e.g., blindness and renal failure) complications of diabetes or life expectancy, benefits for patients can be modest and come with significant costs and raised risk of hypoglycemia (Yudkin et al., 2010; Montori & Fernandez-Balsells, 2009). Health psychology researchers should be cautious about evaluating the importance of psychosocial factors or behavioral interventions solely in terms of glycemic control and other more distal health outcomes, especially when the biobehavioral pathways that may explain these effects go unmeasured or fail to account for observed effects. Well-powered trials that go beyond surrogate outcomes to evaluate the impact of psychological and behavioral interventions on actual diabetes health outcomes, such as progression and onset of complications, are needed to advance the field.

Should psychological interventions that improve patient quality of life, well-being, and self-management, but that have negligible or null effects on glycemic control, be considered failures unworthy of further research or dissemination? We think most patients and clinicians would disagree, but we are less sure about editors, reviewers, policymakers, and funders. Focusing on glycemic control as the most important outcome in behavioral research in

diabetes is inconsistent with evidence-based, patient-centered care (Jones, Vallis, & Pouwer, 2015; Yudkin et al., 2010; Montori et al., 2009). At the same time that we build an evidence base for cost-effective benefits of psychological interventions in terms of diabetes treatment outcomes, we must also advocate for the evaluation of psychological factors and quality of life as essential outcomes of diabetes care. Behavioral science interventions that minimize the psychosocial burdens of diabetes and its treatment and improve health behaviors, diabetes self-management, and functioning likely have value that is insufficiently captured by change in glycemic control from pre- to post-intervention.

The growing prevalence of diabetes points to an increasing need for psychologists competent in working with individuals living with diabetes and their families. In addition to improving the care of those with comorbid psychiatric conditions, psychologists should also have a role in expanding services to the many individuals and families struggling with diabetes without inaccurately casting their problems in the language of psychopathology and abnormality. Successful self-management involves making challenging changes in behavior and maintaining them over time, despite often confusing feedback and emotional distress. These changes run against powerful societal forces—limited preventive care, long work hours, sedentary lifestyle, and unhealthful foods—and compete with other life demands and goals for daily prioritization.

To be effective in supporting children, adolescents, adults, and families in their efforts to effectively manage diabetes, clinicians must gain expert knowledge of diabetes and the goals of diabetes care. To accomplish this, training opportunities at doctoral, internship, and postdoctoral levels should be formalized and expanded. Models of intervention delivery that can integrate psychological care with disease management delivered by interdisciplinary teams are likely to be most effective and efficient. Though evidence reviewed here and elsewhere (e.g., Markowitz et al., 2011) suggests psychological interventions can be effectively delivered in the context of diabetes care, targeting psychosocial factors in isolation is unlikely to be sufficient to improve self-management and health outcomes in diabetes. Interventions involving psychoeducation, distress management, problem-solving skills training, and optimization of self-regulatory processes should also directly target the task of self-management to have maximal impact. Successful interventions will be sensitive to the burdens of diabetes treatment, SES, and sociocultural context; supportive of patient-centered goals for diabetes care; and evidence-based in communicating expected costs and benefits of self-management for quality of life and health to patients, providers, and policymakers.

References

- Aikens, J. E., & Piette, J. D. (2009). Diabetic patients' medication underuse, illness outcomes, and beliefs about antihyperglycemic and antihypertensive treatments. *Diabetes Care*, 32, 19–24. <http://dx.doi.org/10.2337/dc08-1533>
- Al Sayah, F., Majumdar, S. R., Williams, B., Robertson, S., & Johnson, J. A. (2013). Health literacy and health outcomes in diabetes: A systematic review. *Journal of General Internal Medicine*, 28, 444–452. <http://dx.doi.org/10.1007/s11606-012-2241-z>
- American Association of Diabetes Educators. (2015). *AADE7 Self-Care Behaviors*. Retrieved from <http://www.diabeteseducator.org/ProfessionalResources/AADE7/>
- American Diabetes Association. (2015). Standards of medical care in diabetes – 2015. *Diabetes Care*, 38 (Suppl. 1), S1–S89.
- Armstrong, D. G., Kanda, V. A., Lavery, L. A., Marston, W., Mills, J. L., Sr., & Boulton, A. J. (2013). Mind the gap: Disparity between research funding and costs of care for diabetic foot ulcers. *Diabetes Care*, 36, 1815–1817. <http://dx.doi.org/10.2337/dc12-2285>
- Back, R. N., Tanenbaum, M. L., & Gonzalez, J. S. (2014). Diabetes burden and diabetes distress: The buffering effect of social support. *Annals of Behavioral Medicine*, 48, 145–155. <http://dx.doi.org/10.1007/s12160-013-9585-4>
- Bandura, A., & Wood, R. (1989). Effect of perceived controllability and performance standards on self-regulation of complex decision making. *Journal of Personality and Social Psychology*, 56, 805–814. <http://dx.doi.org/10.1037/0022-3514.56.5.805>
- Berg, C. A., Hughes, A. E., King, P. S., Korbel, C., Fortenberry, K. T., Donaldson, D., . . . Wiebe, D. J. (2014). Self-control as a mediator of the link between intelligence and HbA1c during adolescence. *Children's Health Care*, 43, 120–131. <http://dx.doi.org/10.1080/02739615.2013.837819>
- Berkowitz, S. A., Karter, A. J., Lyles, C. R., Liu, J. Y., Schillinger, D., Adler, N. E., . . . Sarkar, U. (2014). Low socioeconomic status is associated with increased risk for hypoglycemia in diabetes patients: The Diabetes Study of Northern California (DISTANCE). *Journal of Health Care for the Poor and Underserved*, 25, 478–490. <http://dx.doi.org/10.1353/hpu.2014.0106>
- Brown, A. F., Ettner, S. L., Piette, J., Weinberger, M., Gregg, E., Shapiro, M. F., . . . Beckles, G. L. (2004). Socioeconomic position and health among persons with diabetes mellitus: A conceptual framework and review of the literature. *Epidemiologic Reviews*, 26, 63–77. <http://dx.doi.org/10.1093/epirev/mxh002>
- Cameron, D. S., Bertenshaw, E. J., & Sheeran, P. (2014). The impact of positive affect on health cognitions and behaviours: A meta-analysis of the experimental evidence. *Health Psychology Review*, 9, 345–365. <http://dx.doi.org/10.1080/17437199.2014.923164>
- Cavanaugh, K., Huizinga, M. M., Wallston, K. A., Gebretsadik, T., Shintani, A., Davis, D., . . . Rothman, R. L. (2008). Association of numeracy and diabetes control. *Annals of Internal Medicine*, 148, 737–746. <http://dx.doi.org/10.7326/0003-4819-148-10-200805200-00006>
- Cavanaugh, K., Wallston, K. A., Gebretsadik, T., Shintani, A., Huizinga, M. M., Davis, D., . . . Rothman, R. L. (2009). Addressing literacy and numeracy to improve diabetes care: Two randomized controlled trials. *Diabetes Care*, 32, 2149–2155. <http://dx.doi.org/10.2337/dc09-0563>
- Celano, C. M., Beale, E. E., Moore, S. V., Wexler, D. J., & Huffman, J. C. (2013). Positive psychological characteristics in diabetes: A review. *Current Diabetes Reports*, 13, 917–929. <http://dx.doi.org/10.1007/s11892-013-0430-8>
- Cho, Y. H., Craig, M. E., Hing, S., Gallego, P. H., Poon, M., Chan, A., & Donaghue, K. C. (2011). Microvascular complications assessment in adolescents with 2- to 5-yr duration of type 1 diabetes from 1990 to 2006. *Pediatric Diabetes*, 12, 682–689. <http://dx.doi.org/10.1111/j.1399-5448.2011.00762.x>
- Compas, B. E., Jaser, S. S., Dunn, M. J., & Rodriguez, E. M. (2012). Coping with chronic illness in childhood and adolescence. *Annual Review of Clinical Psychology*, 8, 455–480. <http://dx.doi.org/10.1146/annurev-clinpsy-032511-143108>
- Cunningham, N. R., Vesco, A. T., Dolan, L. M., & Hood, K. K. (2011). From caregiver psychological distress to adolescent glycemic control: The mediating role of perceived burden around diabetes management. *Journal of Pediatric Psychology*, 36, 196–205. <http://dx.doi.org/10.1093/jpepsy/jsq071>
- Currie, C. J., Peyrot, M., Morgan, C. L., Poole, C. D., Jenkins-Jones, S., Rubin, R. R., . . . Evans, M. (2012). The impact of treatment noncompliance on mortality in people with type 2 diabetes. *Diabetes Care*, 35, 1279–1284. <http://dx.doi.org/10.2337/dc11-1277>
- Daly, J. M., Hartz, A. J., Xu, Y., Levy, B. T., James, P. A., Merchant, M. L., & Garrett, R. E. (2009). An assessment of attitudes, behaviors, and outcomes of patients with type 2 diabetes. *Journal of the American Board of Family Medicine*, 22, 280–290. <http://dx.doi.org/10.3122/jabfm.2009.03.080114>
- Davies, M. J., Heller, S., Skinner, T. C., Campbell, M. J., Carey, M. E., Craddock, S., . . . Diabetes Education and Self Management for Ongoing and Newly Diagnosed Collaborative. (2008). Effectiveness of the diabetes education and self management for ongoing and newly diagnosed (DESMOND) programme for people with newly diagnosed type 2 diabetes: Cluster randomised controlled trial. *BMJ: British Medical Journal*, 336, 491–495. <http://dx.doi.org/10.1136/bmj.39474.922025.BE>
- de Groot, M., Golden, S. H., & Wagner, J. (2016). Psychological conditions in adults with diabetes. *American Psychologist*, 71, 552–562. <http://dx.doi.org/10.1037/a0040408>
- DiMatteo, M. R. (2004). Variations in patients' adherence to medical recommendations: A quantitative review of 50 years of research. *Medical Care*, 42, 200–209. <http://dx.doi.org/10.1097/01.mlr.0000114908.90348.f9>
- Driscoll, K. A., & Young-Hyman, D. (2014). Use of technology when assessing adherence to diabetes self-management behaviors. *Current Diabetes Reports*, 14, 521. <http://dx.doi.org/10.1007/s11892-014-0521-1>
- Duangdao, K. M., & Roesch, S. C. (2008). Coping with diabetes in adulthood: A meta-analysis. *Journal of Behavioral Medicine*, 31, 291–300. <http://dx.doi.org/10.1007/s10865-008-9155-6>
- Ettner, S. L., Cadwell, B. L., Russell, L. B., Brown, A., Karter, A. J., Safford, M., . . . TRIAD Study Group. (2009). Investing time in health: Do socioeconomically disadvantaged patients spend more or less extra time on diabetes self-care? *Health Economics*, 18, 645–663. <http://dx.doi.org/10.1002/hec.1394>
- Feldman, B. S., Cohen-Stavi, C. J., Leibowitz, M., Hoshen, M. B., Singer, S. R., Bitterman, H., . . . Balicer, R. D. (2014). Defining the role of medication adherence in poor glycemic control among a general adult population with diabetes. *PLoS ONE*, 9(9), e108145. <http://dx.doi.org/10.1371/journal.pone.0108145>
- Fisher, L., Gonzalez, J. S., & Polonsky, W. H. (2014). The confusing tale of depression and distress in patients with diabetes: A call for greater clarity and precision. *Diabetic Medicine*, 31, 764–772. <http://dx.doi.org/10.1111/dme.12428>
- Fisher, L., Mullan, J. T., Arean, P., Glasgow, R. E., Hessler, D., & Masharani, U. (2010). Diabetes distress but not clinical depression or depressive symptoms is associated with glycemic control in both cross-sectional and longitudinal analyses. *Diabetes Care*, 33, 23–28. <http://dx.doi.org/10.2337/dc09-1238>
- Fitzpatrick, S. L., Schumann, K. P., & Hill-Briggs, F. (2013). Problem solving interventions for diabetes self-management and control: A systematic review of the literature. *Diabetes Research and Clinical Practice*, 100, 145–161. <http://dx.doi.org/10.1016/j.diabres.2012.12.016>
- Fortenberry, K. T., Butler, J. M., Butner, J., Berg, C. A., Upchurch, R., & Wiebe, D. J. (2009). Perceived diabetes task competence mediates the

- relationship of both negative and positive affect with blood glucose in adolescents with type 1 diabetes. *Annals of Behavioral Medicine*, 37, 1–9. <http://dx.doi.org/10.1007/s12160-009-9086-7>
- French, D. P., Wade, A. N., Yudkin, P., Neil, H. A. W., Kinmonth, A. L., & Farmer, A. J. (2008). Self-monitoring of blood glucose changed non-insulin-treated Type 2 diabetes patients' beliefs about diabetes and self-monitoring in a randomized trial. *Diabetic Medicine*, 25, 1218–1228. <http://dx.doi.org/10.1111/j.1464-5491.2008.02569.x>
- George, J. T., Valdovinos, A. P., Russell, I., Dromgoole, P., Lomax, S., Torgerson, D. J., . . . Thow, J. C. (2008). Clinical effectiveness of a brief educational intervention in Type 1 diabetes: Results from the BITES (Brief Intervention in Type 1 diabetes, Education for Self-efficacy) trial. *Diabetic Medicine*, 25, 1447–1453. <http://dx.doi.org/10.1111/j.1464-5491.2008.02607.x>
- Gonzalez, J. S., Fisher, L., & Polonsky, W. H. (2011). Depression in diabetes: Have we been missing something important? *Diabetes Care*, 34, 236–239. <http://dx.doi.org/10.2337/dc10-1970> (Erratum published 2011, *Diabetes Care*, 34, p. 2488. [10.2337/dc11-er11](http://dx.doi.org/10.2337/dc11-er11))
- Gonzalez, J. S., Peyrot, M., McCarl, L. A., Collins, E. M., Serpa, L., Mimiaga, M. J., & Safren, S. A. (2008). Depression and diabetes treatment nonadherence: A meta-analysis. *Diabetes Care*, 31, 2398–2403. <http://dx.doi.org/10.2337/dc08-1341>
- Gonzalez, J. S., & Schneider, H. E. (2011). Methodological issues in the assessment of diabetes treatment adherence. *Current Diabetes Reports*, 11, 472–479. <http://dx.doi.org/10.1007/s11892-011-0229-4>
- Gonzalez, J. S., Shreck, E., Psaros, C., & Safren, S. A. (2015). Distress and type 2 diabetes-treatment adherence: A mediating role for perceived control. *Health Psychology*, 34, 505–513. <http://dx.doi.org/10.1037/hea0000131>
- Gottfredson, L. S., & Deary, I. J. (2004). Intelligence predicts health and longevity, but why? *Current Directions in Psychological Science*, 13, 1–4. <http://dx.doi.org/10.1111/j.0963-7214.2004.01301001.x>
- Griva, K., Myers, L. B., & Newman, S. (2000). Illness perceptions and self efficacy beliefs in adolescents and young adults with insulin dependent diabetes mellitus. *Psychology & Health*, 15, 733–750. <http://dx.doi.org/10.1080/08870440008405578>
- Gutiérrez, R. R., Ferro, A. M., & Caballero, A. E. (2015). Myths and Misconceptions about Insulin Therapy among Latinos/Hispanics with Diabetes: A Fresh Look at an Old Problem. *J Diabetes Metab*, 6(482), 2.
- Hampson, S. E., Glasgow, R. E., & Foster, L. S. (1995). Personal models of diabetes among older adults: Relationship to self-management and other variables. *The Diabetes Educator*, 21, 300–307. <http://dx.doi.org/10.1177/014572179502100407>
- Harkin, B., Webb, T. L., Chang, B. P., Prestwich, A., Conner, M., Kellar, I., . . . Sheeran, P. (2016). Does monitoring goal progress promote goal attainment? A meta-analysis of the experimental evidence. *Psychological bulletin*, 142, 198–228. <http://dx.doi.org/10.1037/bul0000025>
- Harvey, J. N., & Lawson, V. L. (2009). The importance of health belief models in determining self-care behaviour in diabetes. *Diabetic Medicine*, 26, 5–13. <http://dx.doi.org/10.1111/j.1464-5491.2008.02628.x>
- Hilliard, M. E., Powell, P. W., & Anderson, B. J. (2016). Evidence-based behavioral interventions to promote diabetes management in children, adolescents, and families. *American Psychologist*, 71, 590–601. <http://dx.doi.org/10.1037/a0040359>
- Ho, P. M., Rumsfeld, J. S., Masoudi, F. A., McClure, D. L., Plomondon, M. E., Steiner, J. F., & Magid, D. J. (2006). Effect of medication nonadherence on hospitalization and mortality among patients with diabetes mellitus. *Archives of Internal Medicine*, 166, 1836–1841. <http://dx.doi.org/10.1001/archinte.166.17.1836>
- Hood, K. K., Peterson, C. M., Rohan, J. M., & Drotar, D. (2009). Association between adherence and glycemic control in pediatric type 1 diabetes: A meta-analysis. *Pediatrics*, 124(6), e1171–e1179. <http://dx.doi.org/10.1542/peds.2009-0207>
- Horne, R., Chapman, S. C. E., Parham, R., Freemantle, N., Forbes, A., & Cooper, V. (2013). Understanding patients' adherence-related beliefs about medicines prescribed for long-term conditions: A meta-analytic review of the Necessity-Concerns Framework. *PLoS ONE*, 8(12), e80633. <http://dx.doi.org/10.1371/journal.pone.0080633>
- Houle, J., Beaulieu, M. D., Chiasson, J. L., Lespérance, F., Côté, J., Strychar, I., . . . Lambert, J. (2015). Glycaemic control and self-management behaviours in Type 2 diabetes: Results from a 1-year longitudinal cohort study. *Diabetic Medicine*, 32, 1247–1254. <http://dx.doi.org/10.1111/dme>
- Huang, E. S., Brown, S. E. S., Thakur, N., Carlisle, L., Foley, E., Ewigman, B., & Meltzer, D. O. (2009). Racial/ethnic differences in concerns about current and future medications among patients with type 2 diabetes. *Diabetes Care*, 32, 311–316. <http://dx.doi.org/10.2337/dc08-1307>
- Hughes, A. E., Berg, C. A., & Wiebe, D. J. (2012). Emotional processing and self-control in adolescents with type 1 diabetes. *Journal of Pediatric Psychology*, 37, 925–934. <http://dx.doi.org/10.1093/jpepsy/jss062>
- Hunter, C. M. (2016). Understanding diabetes and the role of psychology in its prevention and treatment. *American Psychologist*, 71, 515–525. <http://dx.doi.org/10.1037/a0040344>
- Janisse, H. C., Naar-King, S., & Ellis, D. (2010). Brief report: Parent's health literacy among high-risk adolescents with insulin dependent diabetes. *Journal of Pediatric Psychology*, 35, 436–440. <http://dx.doi.org/10.1093/jpepsy/jsp077>
- Johnson, P. J., Ghildayal, N., Rockwood, T., & Everson-Rose, S. A. (2014). Differences in diabetes self-care activities by race/ethnicity and insulin use. *The Diabetes Educator*, 40, 767–777. <http://dx.doi.org/10.1177/0145721714552501>
- Johnson, S. B. (1980). Psychosocial factors in juvenile diabetes: A review. *Journal of Behavioral Medicine*, 3, 95–116. <http://dx.doi.org/10.1007/BF00844916>
- Johnson, S. R., Cooper, M. N., Jones, T. W., & Davis, E. A. (2013). Long-term outcome of insulin pump therapy in children with type 1 diabetes assessed in a large population-based case-control study. *Diabetologia*, 56, 2392–2400. <http://dx.doi.org/10.1007/s00125-013-3007-9>
- Jones, A., Vallis, M., & Pouwer, F. (2015). If it does not significantly change HbA1c levels why should we waste time on it? A plea for the prioritization of psychological well-being in people with diabetes. *Diabetic Medicine*, 32, 155–163. <http://dx.doi.org/10.1111/dme.12620>
- Karter, A. J., Ferrara, A., Darbinian, J. A., Ackerson, L. M., & Selby, J. V. (2000). Self-monitoring of blood glucose: Language and financial barriers in a managed care population with diabetes. *Diabetes Care*, 23, 477–483. <http://dx.doi.org/10.2337/diacare.23.4.477>
- Karter, A. J., Stevens, M. R., Herman, W. H., Ettner, S., Marrero, D. G., Safford, M. M., . . . Translating Research Into Action for Diabetes Study Group. (2003). Out-of-pocket costs and diabetes preventive services: The Translating Research Into Action for Diabetes (TRIAD) study. *Diabetes Care*, 26, 2294–2299. <http://dx.doi.org/10.2337/diacare.26.8.2294>
- Keogh, K. M., Smith, S. M., White, P., McGilloway, S., Kelly, A., Gibney, J., & O'Dowd, T. (2011). Psychological family intervention for poorly controlled type 2 diabetes. *The American Journal of Managed Care*, 17, 105–113.
- King, D. K., Glasgow, R. E., Toobert, D. J., Strycker, L. A., Estabrooks, P. A., Osuna, D., & Faber, A. J. (2010). Self-efficacy, problem solving, and social-environmental support are associated with diabetes self-management behaviors. *Diabetes Care*, 33, 751–753. <http://dx.doi.org/10.2337/dc09-1746>
- Kuo, Y. F., Raji, M. A., Markides, K. S., Ray, L. A., Espino, D. V., & Goodwin, J. S. (2003). Inconsistent use of diabetes medications, diabetes complications, and mortality in older Mexican Americans over a 7-year period: Data from the Hispanic established population for the epidemiologic study of the elderly. *Diabetes Care*, 26, 3054–3060. <http://dx.doi.org/10.2337/diacare.26.11.3054>

- Lansing, A. H., & Berg, C. A. (2014). Adolescent self-regulation as a foundation for chronic illness self-management. *Journal of Pediatric Psychology*, 39, 1091–1096. <http://dx.doi.org/10.1093/jpepsy/jsu067>
- Leahey, T. M., Xu, X., Unick, J. L., & Wing, R. R. (2014). A preliminary investigation of the role of self-control in behavioral weight loss treatment. *Obesity Research & Clinical Practice*, 8(2), e149–e153. <http://dx.doi.org/10.1016/j.orcp.2012.12.005>
- Leventhal, H., Diefenbach, M., & Leventhal, E. A. (1992). Illness cognition: Using common sense to understand treatment adherence and affect cognition interactions. *Cognitive Therapy and Research*, 16, 143–163. <http://dx.doi.org/10.1007/BF01173486>
- Li, R., Shrestha, S. S., Lipman, R., Burrows, N. R., Kolb, L. E., Rutledge, S., & Centers for Disease Control and Prevention (CDC). (2014). Diabetes self-management education and training among privately insured persons with newly diagnosed diabetes—United States, 2011–2012. *Morbidity and Mortality Weekly Report*, 63, 1045–1049.
- Lin, M. H., Connor, C. G., Ruedy, K. J., Beck, R. W., Kollman, C., Buckingham, B., . . . Pediatric Diabetes Consortium. (2013). Race, socioeconomic status, and treatment center are associated with insulin pump therapy in youth in the first year following diagnosis of type 1 diabetes. *Diabetes Technology & Therapeutics*, 15, 929–934. <http://dx.doi.org/10.1089/dia.2013.0132>
- Lorig, K., Ritter, P. L., Villa, F., & Piette, J. D. (2008). Spanish diabetes self-management with and without automated telephone reinforcement: Two randomized trials. *Diabetes Care*, 31, 408–414. <http://dx.doi.org/10.2337/dc07-1313>
- Loveman, E., Frampton, G. K., & Clegg, A. J. (2008). The clinical effectiveness of diabetes education models for Type 2 diabetes: A systematic review. *Health Technology Assessment*, 12, 1–116, iii. <http://dx.doi.org/10.3310/hta12090>
- Lynch, E. B., Fernandez, A., Lighthouse, N., Mendenhall, E., & Jacobs, E. (2012). Concepts of diabetes self-management in Mexican American and African American low-income patients with diabetes. *Health Education Research*, 27, 814–824. <http://dx.doi.org/10.1093/her/cys058>
- Macrodimitris, S. D., & Endler, N. S. (2001). Coping, control, and adjustment in Type 2 diabetes. *Health Psychology*, 20, 208–216.
- Maes, S., & Karoly, P. (2005). Self-regulation assessment and intervention in physical health and illness: A review. *Applied Psychology: An International Review*, 54, 267–299. <http://dx.doi.org/10.1111/j.1464-0597.2005.00210.x>
- Mann, D. M., Ponienman, D., Leventhal, H., & Halm, E. A. (2009). Predictors of adherence to diabetes medications: The role of disease and medication beliefs. *Journal of Behavioral Medicine*, 32, 278–284. <http://dx.doi.org/10.1007/s10865-009-9202-y>
- Markowitz, S. M., Gonzalez, J. S., Wilkinson, J. L., & Safren, S. A. (2011). A review of treating depression in diabetes: Emerging findings. *Psychosomatics: Journal of Consultation and Liaison Psychiatry*, 52, 1–18. <http://dx.doi.org/10.1016/j.psych.2010.11.007>
- Mc Sharry, J., Moss-Morris, R., & Kendrick, T. (2011). Illness perceptions and glycaemic control in diabetes: A systematic review with meta-analysis. *Diabetic Medicine*, 28, 1300–1310. <http://dx.doi.org/10.1111/j.1464-5491.2011.03298.x>
- Montori, V. M., & Fernández-Balsells, M. (2009). Glycemic control in type 2 diabetes: Time for an evidence-based about-face? *Annals of Internal Medicine*, 150, 803–808. <http://dx.doi.org/10.7326/0003-4819-150-11-200906020-00008>
- Mulvaney, S. A., Jaser, S. S., Rothman, R. L., Russell, W. E., Pittel, E. J., Lybarger, C., & Wallston, K. A. (2014). Development and validation of the diabetes adolescent problem solving questionnaire. *Patient Education and Counseling*, 97, 96–100. <http://dx.doi.org/10.1016/j.pec.2014.07.005>
- Murphy, H. R., Rayman, G., & Skinner, T. C. (2006). Psycho-educational interventions for children and young people with Type 1 diabetes. *Diabetic Medicine*, 23, 935–943. <http://dx.doi.org/10.1111/j.1464-5491.2006.01816.x>
- Naranjo, D., Hessler, D. M., Deol, R., & Chesla, C. A. (2012). Health and psychosocial outcomes in U.S. adult patients with diabetes from diverse ethnicities. *Current Diabetes Reports*, 12, 729–738. <http://dx.doi.org/10.1007/s11892-012-0319-y>
- Nicolucci, A., Kovacs Burns, K., Holt, R. I. G., Comaschi, M., Hermanns, N., Ishii, H., . . . DAWN2 Study Group. (2013). Diabetes Attitudes, Wishes and Needs second study (DAWN2): Cross-national benchmarking of diabetes-related psychosocial outcomes for people with diabetes. *Diabetic Medicine*, 30, 767–777. <http://dx.doi.org/10.1111/dme.12245>
- Norris, S. L., Engelgau, M. M., & Narayan, K. M. V. (2001). Effectiveness of self-management training in type 2 diabetes: A systematic review of randomized controlled trials. *Diabetes Care*, 24, 561–587. <http://dx.doi.org/10.2337/diacare.24.3.561>
- Nouwen, A., Urquhart Law, G., Hussain, S., McGovern, S., & Napier, H. (2009). Comparison of the role of self-efficacy and illness representations in relation to dietary self-care and diabetes distress in adolescents with type 1 diabetes. *Psychology & Health*, 24, 1071–1084. <http://dx.doi.org/10.1080/08870440802254597>
- Ostini, R., & Kairuz, T. (2014). Investigating the association between health literacy and non-adherence. *International Journal of Clinical Pharmacy*, 36, 36–44. <http://dx.doi.org/10.1007/s11096-013-9895-4>
- Peyrot, M., McMurry, J. F., Jr., & Kruger, D. F. (1999). A biopsychosocial model of glycemic control in diabetes: Stress, coping and regimen adherence. *Journal of Health and Social Behavior*, 40, 141–158. <http://dx.doi.org/10.2307/2676370>
- Polonsky, W. H., Fisher, L., Earles, J., Dudl, R. J., Lees, J., Mullan, J., & Jackson, R. A. (2005). Assessing psychosocial distress in diabetes: Development of the diabetes distress scale. *Diabetes Care*, 28, 626–631. <http://dx.doi.org/10.2337/diacare.28.3.626>
- Polonsky, W. H., Fisher, L., Hessler, D., & Edelman, S. V. (2015). Identifying the worries and concerns about hypoglycemia in adults with type 2 diabetes. *Journal of diabetes and its complications*, 29, 1171–1176.
- Pulgarón, E. R., Sanders, L. M., Patiño-Fernandez, A. M., Wile, D., Sanchez, J., Rothman, R. L., & Delamater, A. M. (2014). Glycemic control in young children with diabetes: The role of parental health literacy. *Patient Education and Counseling*, 94, 67–70. <http://dx.doi.org/10.1016/j.pec.2013.09.002>
- Roblin, D. W., Platt, R., Goodman, M. J., Hsu, J., Nelson, W. W., Smith, D. H., . . . Soumerai, S. B. (2005). Effect of increased cost-sharing on oral hypoglycemic use in five managed care organizations: How much is too much? *Medical Care*, 43, 951–959. <http://dx.doi.org/10.1097/01.mlr.0000178216.23514.b7>
- Rubin, R. R., Peyrot, M., Kruger, D. F., & Travis, L. B. (2009). Barriers to insulin injection therapy: Patient and health care provider perspectives. *The Diabetes Educator*, 35, 1014–1022. <http://dx.doi.org/10.1177/0145721709345773>
- Rubin, R. R., Peyrot, M., & Saudek, C. D. (1993). The effect of a diabetes education program incorporating coping skills training on emotional well-being and diabetes self-efficacy. *The Diabetes Educator*, 19, 210–214. <http://dx.doi.org/10.1177/014572179301900308>
- Russell, L. B., Suh, D. C., & Safford, M. A. (2005). Time requirements for diabetes self-management: Too much for many? *The Journal of Family Practice*, 54, 52–56.
- Ryan, C. M., van Duinkerken, E., & Rosano, C. (2016). Neurocognitive consequences of diabetes. *American Psychologist*, 71, 563–576. <http://dx.doi.org/10.1037/a0040455>
- Sanders, L. M., Federico, S., Klass, P., Abrams, M. A., & Dreyer, B. (2009). Literacy and child health: A systematic review. *Archives of Pediatrics & Adolescent Medicine*, 163, 131–140. <http://dx.doi.org/10.1001/archpediatrics.2008.539>

- Sarkar, U., Fisher, L., & Schillinger, D. (2006). Is self-efficacy associated with diabetes self-management across race/ethnicity and health literacy? *Diabetes Care*, 29, 823–829. <http://dx.doi.org/10.2337/diacare.29.04.06.dc05-1615>
- Schectman, J. M., Nadkarni, M. M., & Voss, J. D. (2002). The association between diabetes metabolic control and drug adherence in an indigent population. *Diabetes Care*, 25, 1015–1021. <http://dx.doi.org/10.2337/diacare.25.6.1015>
- Schmitt, A., Reimer, A., Kulzer, B., Haak, T., Gahr, A., & Hermanns, N. (2015). Negative association between depression and diabetes control only when accompanied by diabetes-specific distress. *Journal of Behavioral Medicine*, 38, 556–564.
- Schneiderman, N., Llabre, M., Cowie, C. C., Barnhart, J., Carnethon, M., Gallo, L. C., . . . Teng, Y. (2014). Prevalence of diabetes among hispanics/latinos from diverse backgrounds: The hispanic community health study/study of latinos (HCHS/SOL). *Diabetes Care*, 37, 2233–2239.
- Simpson, S. H., Eurich, D. T., Majumdar, S. R., Padwal, R. S., Tsuyuki, R. T., Varney, J., & Johnson, J. A. (2006). A meta-analysis of the association between adherence to drug therapy and mortality. *BMJ: British Medical Journal*, 333, 15. <http://dx.doi.org/10.1136/bmj.33875.675486.55>
- Skinner, E. A. (1996). A guide to constructs of control. *Journal of Personality and Social Psychology*, 71, 549–570. <http://dx.doi.org/10.1037/0022-3514.71.3.549>
- Skinner, T. C., & Hampson, S. E. (2001). Personal models of diabetes in relation to self-care, well-being, and glycemic control. A prospective study in adolescence. *Diabetes Care*, 24, 828–833. <http://dx.doi.org/10.2337/diacare.24.5.828>
- Stadler, G., Oettingen, G., & Gollwitzer, P. M. (2010). Intervention effects of information and self-regulation on eating fruits and vegetables over two years. *Health Psychology*, 29, 274–283. <http://dx.doi.org/10.1037/a0018644>
- Stark Casagrande, S., Fradkin, J. E., Saydah, S. H., Rust, K. F., & Cowie, C. C. (2013). The prevalence of meeting A1C, blood pressure, and LDL goals among people with diabetes, 1988–2010. *Diabetes Care*, 36, 2271–2279. <http://dx.doi.org/10.2337/dc12-2258>
- Stark Casagrande, S., Ríos Burrows, N., Geiss, L. S., Bainbridge, K. E., Fradkin, J. E., & Cowie, C. C. (2012). Diabetes knowledge and its relationship with achieving treatment recommendations in a national sample of people with type 2 diabetes. *Diabetes Care*, 35, 1556–1565. <http://dx.doi.org/10.2337/dc11-1943>
- Streisand, R., Branicki, S., Tercyak, K. P., & Kazak, A. E. (2001). Childhood illness-related parenting stress: The pediatric inventory for parents. *Journal of Pediatric Psychology*, 26, 155–162. <http://dx.doi.org/10.1093/jpepsy/26.3.155>
- Stupiansky, N. W., Hanna, K. M., Slaven, J. E., Weaver, M. T., & Fortenberry, J. D. (2013). Impulse control, diabetes-specific self-efficacy, and diabetes management among emerging adults with type 1 diabetes. *Journal of Pediatric Psychology*, 38, 247–254. <http://dx.doi.org/10.1093/jpepsy/jss110>
- Surgenor, L. J., Horn, J., Hudson, S. M., Lunt, H., & Tennent, J. (2000). Metabolic control and psychological sense of control in women with diabetes mellitus. Alternative considerations of the relationship. *Journal of Psychosomatic Research*, 49, 267–273. [http://dx.doi.org/10.1016/S0022-3999\(00\)00181-1](http://dx.doi.org/10.1016/S0022-3999(00)00181-1)
- Surwit, R. S., Feinglos, M. N., & Scovern, A. W. (1983). Diabetes and behavior. A paradigm for health psychology. *American Psychologist*, 38, 255–262. <http://dx.doi.org/10.1037/0003-066X.38.3.255>
- Tanenbaum, M. L., Leventhal, H., Breland, J. Y., Yu, J., Walker, E. A., & Gonzalez, J. S. (2015). Successful self-management among non-insulin-treated adults with Type 2 diabetes: A self-regulation perspective. *Diabetic Medicine*, 32, 1504–1512.
- Taylor, M. D., Frier, B. M., Gold, A. E., Deary, I. J., & Edinburgh Prospective Diabetes Study. (2003). Psychosocial factors and diabetes-related outcomes following diagnosis of Type 1 diabetes in adults: The Edinburgh Prospective Diabetes Study. *Diabetic Medicine*, 20, 135–146. <http://dx.doi.org/10.1046/j.1464-5491.2003.00887.x>
- Toobert, D. J., Hampson, S. E., & Glasgow, R. E. (2000). The summary of diabetes self-care activities measure: Results from 7 studies and a revised scale. *Diabetes Care*, 23, 943–950. <http://dx.doi.org/10.2337/diacare.23.7.943>
- Trief, P. M., Teresi, J. A., Eimicke, J. P., Shea, S., & Weinstock, R. S. (2009). Improvement in diabetes self-efficacy and glycaemic control using telemedicine in a sample of older, ethnically diverse individuals who have diabetes: The IDEATel project. *Age and Ageing*, 38, 219–225. <http://dx.doi.org/10.1093/ageing/afn299>
- Trinacty, C. M., Adams, A. S., Soumerai, S. B., Zhang, F., Meigs, J. B., Piette, J. D., & Ross-Degnan, D. (2009). Racial differences in long-term adherence to oral antidiabetic drug therapy: A longitudinal cohort study. *BMC Health Services Research*, 9, 24. <http://dx.doi.org/10.1186/1472-6963-9-24>
- Vigersky, R. A. (2011). An overview of management issues in adult patients with type 2 diabetes mellitus. *Journal of Diabetes Science and Technology*, 5, 245–250. <http://dx.doi.org/10.1177/193229681100500207>
- Vileikyte, L., & Gonzalez, J. S. (2014). Recognition and management of psychosocial issues in diabetic neuropathy. *Handbook of clinical neurology*, 126, 195–209.
- Walker, R. J., Smalls, B. L., Hernandez-Tejada, M. A., Campbell, J. A., Davis, K. S., & Egede, L. E. (2012). Effect of diabetes fatalism on medication adherence and self-care behaviors in adults with diabetes. *General Hospital Psychiatry*, 34, 598–603. <http://dx.doi.org/10.1016/j.genhosppsych.2012.07.005>
- Watkins, K. W., Connell, C. M., Fitzgerald, J. T., Klem, L., Hickey, T., & Ingersoll-Dayton, B. (2000). Effect of adults' self-regulation of diabetes on quality-of-life outcomes. *Diabetes Care*, 23, 1511–1515. <http://dx.doi.org/10.2337/diacare.23.10.1511>
- Weller, S. C., Baer, R. D., Garcia de Alba Garcia, J., & Salcedo Rocha, A. L. (2012). Explanatory models of diabetes in the U.S. and Mexico: The patient-provider gap and cultural competence. *Social Science & Medicine*, 75, 1088–1096. <http://dx.doi.org/10.1016/j.socscimed.2012.05.003>
- Wiebe, D. J., Helgeson, V., & Berg, C. A. (2016). The social context of managing diabetes across the life span. *American Psychologist*, 71, 526–538. <http://dx.doi.org/10.1037/a0040355>
- Wild, D., von Maltzahn, R., Brohan, E., Christensen, T., Clauson, P., & Gonder-Frederick, L. (2007). A critical review of the literature on fear of hypoglycemia in diabetes: Implications for diabetes management and patient education. *Patient Education and Counseling*, 68, 10–15. <http://dx.doi.org/10.1016/j.pec.2007.05.003>
- Wing, R. R., Tate, D. F., Gorin, A. A., Raynor, H. A., & Fava, J. L. (2006). A self-regulation program for maintenance of weight loss. *The New England Journal of Medicine*, 355, 1563–1571. <http://dx.doi.org/10.1056/NEJMoa061883>
- Wood, J. R., Miller, K. M., Maahs, D. M., Beck, R. W., DiMeglio, L. A., Libman, I. M., . . . Woerner, S. E. (2013). Most youth with type 1 diabetes in the T1D exchange clinic registry do not meet American Diabetes Association or International Society for Pediatric and Adolescent Diabetes clinical guidelines. *Diabetes Care*, 36, 2035–2037.
- Yudkin, J. S., Richter, B., & Gale, E. A. M. (2010). Intensified glucose lowering in type 2 diabetes: Time for a reappraisal. *Diabetologia*, 53, 2079–2085.

Received March 7, 2015

Revision received December 18, 2015

Accepted March 22, 2016 ■